



# The Brazilian Journal of INFECTIOUS DISEASES

[www.elsevier.com/locate/bjid](http://www.elsevier.com/locate/bjid)



## Original article

# Infections after shoulder arthroplasty are correlated with higher anesthetic risk score: a case-control study in Brazil



Leonardo Hideto Nagaya<sup>a</sup>, Mauro José Costa Salles<sup>b,\*</sup>,  
Lucas Sadawo Chagas Takikawa<sup>a</sup>, Marcelo Fregoneze<sup>a</sup>, Pedro Doneux<sup>a</sup>,  
Luciana Andrade da Silva<sup>a</sup>, Guilherme do Val Sella<sup>a</sup>, Alberto Naoki Miyazaki<sup>a</sup>,  
Sergio Luiz Checchia<sup>a</sup>

<sup>a</sup> Faculdade de Ciências Médicas da Santa Casa de São Paulo, Departamento de Ortopedia e Traumatologia, São Paulo, SP, Brazil

<sup>b</sup> Faculdade de Ciências Médicas da Santa Casa de São Paulo, Departamento de Medicina Interna, Seção de Doenças Infecciosas, São Paulo, SP, Brazil

## ARTICLE INFO

### Article history:

Received 11 March 2017

Accepted 11 June 2017

Available online 10 July 2017

### Keywords:

Orthopedic infection

Hemiarthroplasty

Total shoulder arthroplasty

American Society of

Anesthesiologists score

Risk factor

*Pseudomonas aeruginosa*

## ABSTRACT

**Purposes:** Shoulder arthroplasty (SA) has been performed by many years for the treatment of several conditions, including osteoarthritis and proximal humeral fractures following trauma. Surgical site infection (SSI) following Shoulder arthroplasty remains a challenge, contributing to increased morbidity and costs. Identification of risk factors may help implementing adequate strategies to prevent infection. We aimed to identify pre- and intra-operative risk factors associated with deep infections after Shoulder arthroplasty.

**Methods:** An unmatched case-control study was conducted to describe the prevalence, clinical and microbiological findings, and to evaluate patient and surgical risk factors for prosthetic shoulder infection (PSI), among 158 patients who underwent SA due to any reason, at a tertiary public university institution. Risk factors for PSI was assessed by uni- and multivariate analyses using multiple logistic regression.

**Results:** 168 SA from 158 patients were analyzed, with an overall infection rate of 9.5% (16/168 cases). Subjects undergoing SA with American Society of Anesthesiologists (ASA) grade III or higher (odds ratio [OR] = 5.30, 95% confidence interval [CI] = 1.58–17.79,  $p < 0.013$ ) and presenting local hematoma after surgery (odds ratio [OR] = 7.10, 95% confidence interval [CI] = 1.09–46.09,  $p = 0.04$ ) had higher risk for PSI on univariate analysis. However, only ASA score grade III or higher remained significant on multivariate analysis (OR = 4.74, 95% CI = 1.33–16.92,  $p = 0.016$ ). Gram-positive cocci and Gram-negative bacilli were equally isolated in 50% of cases; however, the most commonly detected bacterium was *Pseudomonas aeruginosa* (18.7%).

\* Corresponding author.

E-mail address: [salles.infecto@gmail.com](mailto:salles.infecto@gmail.com) (M.J. Salles).

<http://dx.doi.org/10.1016/j.bjid.2017.06.003>

1413-8670/© 2017 Sociedade Brasileira de Infectologia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

**Conclusion:** This study provides evidence suggesting that patient-related known factors such as higher ASA score predisposes to shoulder arthroplasty-associated infection. Furthermore, unusual pathogens associated with PSI were identified.

© 2017 Sociedade Brasileira de Infectologia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Introduction

Hemiarthroplasty (HA) and total shoulder arthroplasty (TSA) have been increasingly performed for the treatment of several osteoarticular conditions such as osteoarthritis, humeral head avascular necrosis and proximal humeral fractures following traumas.<sup>1,2</sup> Indeed, the annual number of HA and TSA performed in the United States is growing faster, reaching more than 50,000 shoulder replacements per year.<sup>3</sup> The incidence of periprosthetic shoulder infections (PSI) appears to be less than that of periprosthetic hip (THA) and knee (TKA) infections, though there have been reports of higher rates in the surgery literature.<sup>4,5</sup> Periprosthetic shoulder infections, nonetheless, remain challenging as they increase morbidity as well as raise costs.<sup>6</sup> The rate of infectious complication following primary shoulder arthroplasties appears to be as lower as the rates associated with primary hip and knees arthroplasties, while few single-center studies have demonstrated higher rates of PSI.<sup>1,4–6</sup>

There have been few previous studies analyzing risk factors for PSI.<sup>7–9</sup> Patient related-factors and comorbidities such as male gender, younger age, diabetes, morbid obesity, rheumatoid arthritis, and other types of immunosuppression including tumors, have already been implicated.<sup>7,10</sup> Surgical-related factors such as peri-operative hematomas and prior surgeries, especially in the trauma setting, increase the incidence of infection after primary shoulder arthroplasty.<sup>1,7,11</sup>

In addition, the role of biofilm-forming bacteria, such as *Propionibacterium acnes*, but also *Staphylococcus*, and *Streptococcus* in the pathogenesis of orthopedic implant-associated infections, particularly PSI has been widely accepted.<sup>12–16</sup> Although uncommonly implicated in PSI, *Pseudomonas* is an aerobic Gram-negative bacillus expressing the ability to form complex biofilm structures on the surfaces of orthopedic implants.<sup>17,18</sup> In one large study addressing Gram-negative prosthetic joint infections, *Pseudomonas aeruginosa* was by far the most frequently (40%) isolated pathogen.<sup>19</sup>

We herein sought to identify pre- and intra-operative risk factors predisposing subjects to develop deep infections following HA and TSA, and evaluate causative microorganisms associated with PSI.

## Methods

### Study population

We performed a single-center case-control study with 183 subjects submitted to either HA or TSA that were performed at the Department of Orthopedics and Traumatology of a large urban public teaching hospital, between January 1987 and November 2012. The study included subjects with at least 24 months of follow-up after the surgical implantation of shoulder arthroplasty. Exclusion criteria were age less than 18 years, follow up of less than 24 months, arthroplasties performed for malignant etiologies, prior radiation of the operative site, and patients whose medical records were unavailable. The study was reviewed and approved by the local Institutional Review Board.

### Diagnosis of periprosthetic shoulder infection (PSI)

PSI was diagnosed according to IDSA guidelines by the presence of a sinus tract communicating with the prosthesis, histopathological analyses with the presence of inflammatory cells, visible purulence surrounding the prosthesis, and/or identical microorganisms isolated from two or more cultures. PSI was categorized as early when diagnosed before three months after surgery; intermediate when diagnosed between three and 24 months after surgery; and late infection when diagnosed after two years of prosthesis implantation.<sup>20</sup> Subjects who fulfilled the diagnostic criteria for PSI were considered cases for this study. Relapses at the same joint prosthesis were not considered for analysis. Controls were subjects who had undergone shoulder arthroplasty due to any indication except joint infection and did not develop PSI during follow-up.

### Potential risk factors

To identify potential risk factors associated with PSI, several variables (patient, microbiological findings, and surgery associated variables) were assessed by reviewing medical, intra-operative, and microbiological records. We searched for demographic variables, comorbidities, American Society of Anesthesiologists (ASA) classification, preoperative diagnosis,

primary or revision arthroplasty, previous history of local or systemic infection, use of systemic antibiotic 14 days prior to surgery, or any hospital admission at least three months before surgery. Surgical-related variables including type of implant, need for blood transfusion, duration of surgery, and surgeon's experience level were recorded. The level of experience of the surgeon was categorized as high for senior doctors and head of the group; average for new graduates; and low for medical residents.

### Statistical analysis

For statistical analysis qualitative variables were described as mean and percentage, and quantitative variables were described as median and standard deviation. The association between qualitative variables were assessed by chi-square test and Fisher's exact test. For associations between quantitative variables we used univariate logistic regression. Risk estimates were calculated for the variables associated with PSI and reported as odds ratios with 95% confidence interval (CI). Variables that in univariate analysis had a *p*-value less than 0.20 were included in the multiple logistic regression model to assess their independent association with PSI. Epi-Info<sup>®</sup> Version 3.22 software was used for data tabulation and SPSS version 20.0 (SPSS, Chicago, IL, USA) for statistical analysis.

## Results

### Study population

During the study period of 1987–2012, a total of 183 subjects were included. Twenty-five subjects (13.6%) were excluded from further analysis: 10 due to lack of complete medical information, 12 had less than 24 months of follow up, and three had a diagnosis of shoulder tumor. One-hundred and sixty-eight shoulder implants in 158 subjects (10 subjects had bilateral shoulder prosthesis implanted at different time intervals) were analyzed. Eighty-one percent of these implants were shoulder hemiarthroplasties. After 24 months of follow up, PSI were observed in 9.5% (16/168) of analyzed cases. The clinical characteristics and comorbidities were similar in infected (PSI) and non-infected (NISA) groups (Table 1). Of note, among the 16 subjects presenting PSI, 14 (87.5%) had hemiarthroplasties and two (12.5%) total arthroplasties; in 13 out of the 16 (81.2%) PSI occurred during the first half of the study period while only three (18.8%) occurred in the second half, though this did not reach statistical significance (*p* = 0.103). Table 2 summarizes the 16 clinical cases of PSI concerning demographics, ASA score, diagnosis prior to arthroplasty, type of surgery, presence of hematoma, time elapsed between surgery and PSI, microorganism identified, and the selected medical treatment.

### Risk factors for PSI

The results of the uni- and multivariate analysis of risk factors for PSI are presented in Table 1. Demographic variables, comorbidities, underlying diagnosis, and all other pre-operative characteristics were not significantly associated with PSI. On the other hand, subjects undergoing shoulder arthroplasty

with ASA grade III or higher (odds ratio [OR] = 5.30, 95% confidence interval [CI] = 1.58–17.79, *p* < 0.013) and presenting local hematoma after surgery (odds ratio [OR] = 7.10, 95% confidence interval [CI] = 1.09–46.09, *p* = 0.04) had higher risk for PSI on univariate analysis. Intra-operative aspects such as HA or TSA, blood transfusion (yes/no), duration of surgery longer than 2 h, antibiotic prophylaxis, and surgeon's experience were not significantly associated with PSI. On multivariable analysis, only ASA score III or higher remained statistically associated with PSI (odds ratio [OR] = 4.74, 95% confidence interval [CI] = 1.33–16.92, *p* = 0.016), though hematoma formation showed a trend toward significance (odds ratio [OR] = 6.51, 95% confidence interval [CI] = 0.84–50.70, *p* = 0.074).

### Microbiological findings

Periprosthetic tissue and synovial fluid cultures identified no pathogens in 43.7% (7/16) of PSI subjects, all of which were using systemic empirical antibiotics prior to surgery. Sonication of retrieved shoulder arthroplasty was performed in only one infected shoulder implant, yielding *P. aeruginosa*, *Staphylococcus aureus*, and *Proteus mirabilis* (patient number 14). The infection was polymicrobial in two cases (12.5%). Gram-negative bacilli (GNB) were detected on 50% of cultures, mostly *P. aeruginosa* (18.7%). Interestingly, *P. acnes* was not found in this case series. Table 3 shows the microbiological findings of the 16 patients with PSI.

## Discussion

There have been few studies describing the epidemiology of periprosthetic joint infection (PJI) after shoulder arthroplasty, including microbial etiology, incidence, and predisposing factors. We aimed to investigate the risks of PJI as well as to understand the underlying microbiological aspects in a cohort of subjects who underwent HA and PSA over 25-year period, 1987–2012. While the rates of PJI following TKA and THA seems to be increasing,<sup>10</sup> PSI remains relatively stable according to a large US database survey.<sup>6</sup> Interestingly, in our study the rate of PSI reached 12.4% during the first 12-year period of 1987–1999, but dropped to 4.8% in the subsequent 13-year period of 2000–2012, although this difference was not statistically significant. Many reasons may have been associated with the reduction of PSI rate over time, but we argue that an improvement in the pre- and intra-operative management of trauma patients in our urban tertiary teaching center may have played a role.

We sought to identify potential risk factors for PSI, and when examining specifically pre-operative variables, such as demographic data and underlying medical comorbidities, no significant association with PSI was found. Conversely, previous studies have associated younger male patients and PSI, possibly due to local trauma and surgical procedures performed for fractures,<sup>6,7</sup> while others studies showed PSI to be significantly associated with chronic diseases, including diabetes and rheumatoid arthritis.<sup>1</sup> We identified higher rates of proximal humeral acute fractures as the main reason for shoulder arthroplasty; however, we could not find an association with infection.

**Table 1 – Demographics, comorbidities, and operative characteristics of PSI and NISA study groups. Univariate and multivariate analysis of risk factors associated to develop PSI.**

Variables	Prosthetic shoulder surgeries		Univariate analysis				Multivariate analysis			
	NISA (n = 152)	PSI (n = 16)	OR	IC (95%)		p-value	OR	IC (95%)		p-value
				Lower	Upper			Lower	Upper	
<b>Demographic data</b>										
Male	37 (24.3)	7 (43.7)	2.42	0.84	6.94	0.131 <sup>a</sup>	0.68	0.84	50.7	0.174
Age										
Average	58.4 (14.1)	55.8 (15.7)	0.99	0.95	1.02	0.486 <sup>c</sup>	1.01	0.96	1.06	0.693
Median (range min- max)	60.5 (22-80)	55 (22-77)								
<b>Comorbidities</b>										
Hypertension	58 (38.1)	8 (50)	1.62	0.58	4.55	0.356				
DM	18 (11.8)	2 (12.5)	1.06	0.22	5.07	>0.999				
Rheumatoid arthritis	7 (21.7)	1 (6.3)	1.38	0.16	11.99	0.559				
Pulmonary comorbidity	1 (0.6)	0 (0)	a			>0.999				
HIV	2 (1.3)	0 (0)	a			>0.999				
Smoker	30 (19.7)	4 (25)	1.36	0.41	4.50	0.743				
Alcohol dependency	12 (7.9)	3 (18.7)	2.69	0.67	10.78	0.158				
Sickle cell disease	3 (2)	0 (0)	a			>0.999				
Hypothyroidism	8 (5.2)	0 (0)	a			>0.999				
Hepatitis	3 (2)	1 (6.3)	3.31	0.32	33.85	0.332				
Epilepsy	11 (7.2)	1 (6.3)	0.86	0.10	7.09	>0.999				
Corticoid use	1 (0.6)	0 (0)	a			>0.999				
Chemotherapy	2 (1.3)	0 (0)	a			>0.999				
<b>Preoperative characteristics</b>										
Diagnosis						0.148 <sup>b</sup>				
Acute fracture PH	59 (38.8)	10 (62.5)	1.00							
LPTN	48 (31.5)	4 (25)	0.49	0.15	1.67					
AAHN	45 (29.6)	2 (12.5)	0.26	0.06	1.26					
Reoperation	22 (14.5)	2 (12.5)	0.84	0.18	3.97	>0.999				
Previous Antibiotic Use <sup>A</sup>	13 (8.6)	1 (6.3)	0.71	0.09	5.84	>0.999				
Previous Infection <sup>A</sup>	6 (3.9)	0 (0)	a			>0.999				
Hospitalization <sup>A</sup>	8 (5.3)	2 (12.5)	2.57	0.50	13.31	0.244				
<b>Preoperative hospitalization</b>										
>7 days	51 (33.6)	6 (37.5)	1.17	0.40	3.39	0.779				
<b>Intraoperative characteristics</b>										
ASA class										
I/II	140 (92.1)	11 (68.8)	1.00				1.00			
III	12 (7.9)	5 (31.2)	5.3	1.58	17.79	0.013	4.74	1.33	16.92	0.016
Surgery						0.739				
Hemiarthroplasty	30 (19.7)	2 (12.5)	1.00							
TSA	122 (80.3)	14 (87.5)	1.72	0.37	7.99					
Intraoperative BT	2 (1.3)	0 (0)	a			>0.999				
Operative length of time										
>2h	135 (88.9)	16 (100)	A			0.369				
Antibiotic prophylaxis						0.523				
Cefazoline	142 (93.4)	15 (93.8)	0.53	0.06	4.82					
Surgeon experience						0.393				
High	135 (88.8)	15 (93.8)	1.00							
Median	9 (5.9)	0 (0)	a							
Low	8 (5.3)	1 (6.3)	1.13	0.13	9.62					
<b>Postoperative characteristics</b>										
Hematoma requiring drainage	3 (2)	2 (12.5)	7.10	1.09	46.09	0.04 <sup>c</sup>	6.51	0.84	50.70	0.074
Postoperative BT	2 (1.3)	0 (0)	a			>0.999				

Chi-square test; a, OR, odds ratio; IC (95%), 95% of confidence interval; DM, diabetes mellitus; HIV, human immunodeficiency virus; min, minimum; max, maximum; PSI, periprosthetic shoulder infection; NISA, non infected shoulder arthroplasty; Acute fracture PH, acute fracture of the proximal humerus; TSA, total shoulder arthroplasty; A, last three months; AANU, arthritis, arthropathy and humeral necrosis; LPTN, luxations and post-traumatic humeral necrosis; BT, blood transfusion.

<sup>b</sup> Likelihood ratio test.

<sup>c</sup> Logistic regression analysis.

**Table 2 – Summary of 16 cases of infection after shoulder arthroplasty.**

Case	Gender	Age	ASA	Reoperation	Diagnosis	Surgery	Hematoma	Period of infection	Etiological agent
1	F	69	I	No	Glenohumeral arthrosis	H	No	Early	<i>Enterococcus sp</i>
2	M	75	III	No	Posterior shoulder dislocation	H	No	Late	<i>S. aureus</i>
3	F	72	I	No	PHF	H	No	Delayed	<i>P. aeruginosa</i>
4	F	80	III	No	PHF	H	No	Early	<i>Enterobacter sp</i>
5	M	78	II	No	PHF	H	No	Delayed	<i>S. epidermidis</i>
6	F	60	I	No	Glenohumeral arthrosis	TSA	No	Late	Unidentified
7	M	65	III	No	PHF	H	No	Late	<i>P. aeruginosa</i>
8	M	48	II	No	PHF	H	No	Early	<i>Enterococcus sp</i>
9	F	77	II	No	PHF	H	Yes	Early	Unidentified
10	F	98	III	No	PHF	H	No	Delayed	Unidentified
11	M	62	III	Yes	PHF	H	Yes	Early	<i>S. epidermidis, M. morgani</i>
12	F	86	II	No	PHF	H	No	Early	Unidentified
13	M	86	II	No	Post-traumatic humeral necrosis	H	No	Delayed	Unidentified
14	F	81	II	No	PHF	H	No	Early	<i>P. mirabilis, S. aureus, P. aeruginosa</i>
15	M	59	I	No	PHF	H	No	Late	Unidentified
16	F	80	II	Yes	Post-traumatic humeral necrosis	TSA	No	Early	Unidentified

Source: Medical files from DOTSCMSP.

*S. aureus*, *Staphylococcus aureus*; *S. epidermidis*, *Staphylococcus epidermidis*; *M. morgani*, *Morganela morgani*; *P. mirabilis*, *Proteus mirabilis*; *P. aeruginosa*, *Pseudomonas aeruginosa*; F, female; M, male; PHF, proximal humeral fracture; H, hemiarthroplasty; TSA, total shoulder arthroplasty; ASA, American Society of Anesthesiologists.



**Table 3 – Microbiological findings of 16 patients with periprosthetic shoulder-associated infection.**

Microorganisms	n (%)
<i>Pseudomonas aeruginosa</i>	3 (18.7%)
<i>Enterococcus faecalis</i>	2 (12.5%)
<i>Staphylococcus aureus</i>	2 (12.5%)
<i>Staphylococcus epidermidis</i>	2 (6.2%)
<i>Proteus mirabilis</i>	1 (6.2%)
<i>Morganella morganii</i>	1 (6.2%)
<i>Enterobacter sp</i>	1 (6.2%)
Unidentified	7 (43.7%)

Source: Department of Orthopedics e Traumatology of Santa Casa de São Paulo School of Medical sciences medical files.

Compared to HA, TSA is associated with increased difficulty, increased need for blood transfusions, and longer operative times, though recently published studies show superior outcomes with equivalent postoperative complications, including PSI.<sup>1,2,7</sup> In the present study, 6.2% of TSA became infected, even though TSA was not statistically associated with PSI. In contrast, we identified a statistically significant association between PSI and patients presenting ASA grade III or higher, which was confirmed on multivariate analysis. In fact, patients presenting ASA grade three or higher had 4.74-fold higher risk of PSI. To the best of our knowledge, no previous studies have identified significant association of ASA grade particularly with shoulder arthroplasty infection. A retrospective cohort study of patients undergoing TSA or reverse TSA with an ASA score grade III or higher found greater risk of surgical complications, including nerve damage and perioperative fracture, but not higher incidence of surgical site infections.<sup>21</sup> Composite risk scores aggregate different factors into one and ASA is a six-point scale frequently applied pre-operatively to assess patient's global health and risks for postoperative complications.<sup>22</sup> Furthermore, an elevated ASA grade has been shown to increase the rate of infectious complications after orthopedic surgeries, including PJI following THA and TKA and spinal surgeries.<sup>10,22</sup> Additionally, in the current study the presence of hematoma formation soon after shoulder arthroplasty was found to be a potential risk of PSI, showing a 7.10-fold higher risk of PSI on univariate analysis, though it did not reach statistical significance likely secondary to the low incidence of PSI in our study. Cheung et al.<sup>23</sup> have previously showed an association of postoperative large hematoma formation requiring reoperation and PJI after primary or revision shoulder arthroplasty, in which hematoma formation after PSA was frequently associated with positive intraoperative cultures.

The presence of hair follicles and sebaceous glands do not increase the adherence of bacteria; this is an inherent property of the bacterium. Rather, they likely harbor bacteria that cannot be adequately sterilized with routine peri-operative sterilization procedures.

Studies have associated PSI with higher frequency of chest bacterial colonization, especially the adherence of anaerobic-aerotolerant Gram-positive bacilli such as *P. acnes* and other colonizers of human skin like *Staphylococcus*, *Streptococcus*, and *Pseudomonas* spp.<sup>12-14</sup> Indeed, an increasingly amount of recent data have demonstrated the predominance of *P. acnes*

in PSI.<sup>7,10,12-16</sup> Importantly, our microbiological findings added new information to the epidemiology of PSI as GNB were identified in half of periprosthetic cultures. Notwithstanding, because no *P. acnes* was recovered from our PSI cases, we have modified the microbiological protocol for the diagnosis of PJI in our Institution. Since 2009 we included sonication technique of the retrieved implants, tissue, and synovial fluid culture incubation on thioglycolate broth for up to 14 days. These findings may reflect the fact that in the majority of our PSI cases presented as early-onset, and local trauma had been the main reason for primary shoulder arthroplasty. Hsieh et al., identified 15% of GNB-PJI after THA and TKA, occurring mainly in the early postoperative period, in which *P. aeruginosa* was the principal pathogen.<sup>19</sup> Reasons for the high frequency of GNB-PSI in our study may include: (1) previous use of broad-spectrum systemic antibiotics such as cephalosporin in trauma patients may have selected more naturally resistant pathogens; (2) since 2011 we started applying sonication culture method for the microbial diagnosis of PJI, thus increasing microbial identification of biofilm-forming sessile pathogens. The presence of *S. aureus* and *S. epidermidis* in a quarter of our PSI patients was not surprising, in contrast to the 12.5% detection of *Enterococcus faecalis*. Much like GNB, enterococci are more commonly associated with early postoperative PJI and as part of polymicrobial infection.<sup>10</sup> Since previous studies have associated the *P. acnes* with PSI,<sup>13,24</sup> we have then applied sonication cultures for the microbial diagnosis of different orthopedic implant-associated infections, and prolonged the culture incubation period to 14 days. So far, we have isolated many fastidious and difficult-to-grow bacteria, but not *P. acnes*.<sup>25</sup> Unfortunately, we have been unable to understand why *P. acnes* shows no growth after plating sonication fluid and tissue samples onto solid culture media and liquid broth under anaerobic condition, and incubating for at least 14 days. Perhaps, its growth may be hampered by the concomitant presence of others pathogens on polymicrobial infections.<sup>12</sup> As previously speculated by Singh et al., we also believe that many of the culture-negative results on clinically diagnosed PSI cases would have been caused by *P. acnes*.<sup>7</sup>

We are aware that our study has several important limitations, of course due to its retrospective design and a small number of PSI cases (type II error), thus limiting the performance of multivariate analysis to many important associations with PSI such as presence of hematoma and type of fracture. Interpretation by readers of odds ratio calculation using only binominal logistic regression should be made with caution, because it may overestimate the true effect if the variables analyzed are common. The study was carried out in a large urban public teaching center located in a developing country crowded mega-city which offers a specialized orthopedic care for the local population. Studies addressing PSI risk factors among different populations may not reproduce these results, thus the generalizability of our conclusions should be seen with caution. On the other hand, one important strength of our study is that we performed an active surveillance among all subjects submitted to HA and TSA for at least two years after surgery. Another strength was the large number of trauma patients included in the analysis, thus generating a good epidemiological data among this frequent reason for shoulder arthroplasty.

Despite these limitations, our study revealed higher rates of PSI after HA and TSA, estimated at 9.5% over a 25-year period. PSI was mainly associated with an underlying diagnosis of local trauma, and on multivariate analysis, only ASA grade three or higher was associated with increased risk of PSI. Conversely, controlling for individual patient comorbidity, we found no increased association with PSI. Gram-negative bacilli, especially *P. aeruginosa* was an important cause of infection and *P. acnes* was neither detected in periprosthetic tissues nor on sonication fluid cultures. Since 2013, PSI rates after shoulder arthroplasty have reduced dramatically in our center, by applying infection control measures especially on trauma patients requiring HA or PSA, which improved the quality of care. Prospective studies including larger number of patients are needed to examine potential risk factor for PSI after HA and TSA.

### Conflicts of interest

The authors declare no conflicts of interest.

### Acknowledgments

We thank Rogério Ruscitto do Prado for his support in statistical analysis.

### REFERENCES

- Franceschini V, Chillemi C. Periprosthetic shoulder infection. *Open Orthop J.* 2013;28:243–9.
- Shields E, Iannuzzi JC, Thorsness R, Noyes K, Voloshin I. Perioperative complications after hemiarthroplasty and total shoulder arthroplasty are equivalent. *J Shoulder Elbow Surg.* 2014;23:1449–53.
- Kim SH, Wise BL, Zhang Y, Szabo RM. Increasing incidence of shoulder arthroplasty in the United States. *J Bone Joint Surg Am.* 2011;93:2249–54.
- Fehringer EV, Mikuls TR, Michaud KD, Henderson WG, O'Dell JR. Shoulder arthroplasties have fewer complications than hip or knee arthroplasties in US veterans. *Clin Orthop Relat Res.* 2010;468:717–22.
- Farmer KW, Hammond JW, Queale WS, Keyurapan E, McFarland EG. Shoulder arthroplasty versus hip and knee arthroplasties: a comparison of outcomes. *Clin Orthop Relat Res.* 2007;455:183–9.
- Padegimas EM, Maltenfort M, Ramsey ML, Williams GR, Parvizi J, Namdari S. Periprosthetic shoulder infection in the United States: incidence and economic burden. *J Shoulder Elbow Surg.* 2015;24:741–6.
- Singh JA, Sperling JW, Schleck C, Harmsen W, Cofield RH. Periprosthetic infections after shoulder total shoulder arthroplasty: a 33-year perspective. *J Shoulder Elbow Surg.* 2012;21:1534–41.
- Sperling JW, Kozak TK, Hanssen AD, Cofield RH. Infection after shoulder arthroplasty. *Clin Orthop Relat Res.* 2001;382:206–16.
- Weber P, Utzschneider S, Sadoghi P, Andress HJ, Jansson V, Müller PE. Management of the infected shoulder prosthesis: a retrospective analysis and review of the literature. *Int Orthop.* 2011;35:365–73.
- Tande AJ, Patel R. Prosthetic joint infection. *Clin Microbiol Rev.* 2014;27:302–45.
- Coste JS, Reig S, Trojani C, Berg M, Walch G, Boileau P. The management of infection in arthroplasty of the shoulder. *J Bone Joint Surg Br.* 2004;86:65–9.
- Achermann Y, Goldstein EJ, Coenye T, Shirtliff ME. *Propionibacterium acnes*: from commensal to opportunistic biofilm-associated implant pathogen. *Clin Microbiol Rev.* 2014;27:419–40.
- Portillo ME, Corvec S, Borens O, Trampuz A. *Propionibacterium acnes*: an underestimated pathogen in implant-associated infections. *Biomed Res Int.* 2013:804391.
- Piper KE, Jacobson MJ, Cofield RH, et al. Microbiologic diagnosis of prosthetic shoulder infection by use of implant sonication. *J Clin Microbiol.* 2009;47:1878–84.
- Bayston R, Ashraf W, Barker-Davies R, et al. Biofilm formation by *Propionibacterium acnes* on biomaterials in vitro and in vivo: impact on diagnosis and treatment. *J Biomed Mater Res.* 2007;81:705–9.
- Ramage G, Tunney MM, Patrick S, Gorman SP, Nixon JR. Formation of *Propionibacterium acnes* biofilms on orthopaedic biomaterials and their susceptibility to antimicrobials. *Biomaterials.* 2003;24:3221–7.
- Neut D, Hendriks JG, van Horn JR, van der Mei HC, Busscher HJ. *Pseudomonas aeruginosa* biofilm formation and slime excretion on antibiotic-loaded bone cement. *Acta Orthop.* 2005;76:109–14.
- Rodríguez-Pardo D, Pigrau C, Lora-Tamayo J, et al. REIPI Group for the Study of Prosthetic Infection. Gram-negative prosthetic joint infection: outcome of a debridement, antibiotics and implant retention approach. A large multicentre study. *Clin Microbiol Infect.* 2014;20:911–9.
- Hsieh PH, Lee MS, Hsu KY, Chang YH, Shih HN, Ueng SW. Gram-negative prosthetic joint infections: risk factors and outcome of treatment. *Clin Infect Dis.* 2009;49:1036–43.
- Osmon DR, Berbari EF, Berendt AR, et al. Diagnosis and management of prosthetic joint infection: clinical practice guidelines by the Infectious Diseases Society of America. *Clin Infect Dis.* 2013;56:e1–25.
- Johnson CC, Sodha S, Garzon-Muvdi J, Petersen SA, McFarland EG. Does preoperative American Society of Anesthesiologists score relate to complications after total shoulder arthroplasty? *Clin Orthop Relat Res.* 2014;472:1589–96.
- Whitmore RG, Stephen JH, Vernick C, et al. ASA grade and Comorbidity Index of spinal surgery patients: correlation with complications and societal costs. *Spine J.* 2014;14:31–8.
- Cheung EV, Sperling JW, Cofield RH. Infection associated with hematoma formation after shoulder arthroplasty. *Clin Orthop Relat Res.* 2008;466:1363–7.
- Achermann Y, Sahin F, Schwyzer HK, Kolling C, Wust J, Vogt M. Characteristics and outcome of 16 periprosthetic shoulder joint infections. *Infection.* 2013;41:613–20.
- Yano MH, Klautau GB, da Silva CB, et al. Improved diagnosis of infection associated with osteosynthesis by use of sonication of fracture fixation implants. *J Clin Microbiol.* 2013;52:4176–82.