




# ACTAS Derma-Sifiliográficas

Full English text available at  
[www.actasdermo.org](http://www.actasdermo.org)



## IMAGES IN DERMATOLOGY

### Dissecting cellulitis of the scalp

### Celulitis disecante del cuero cabelludo

A. Ortiz-Prieto\*, A. Pulpillo-Ruiz, J. Conejo-Mir



*Servicio de Dermatología, Hospital Universitario Virgen del Rocío, Sevilla, Spain*

A 38-year-old Caucasian male presented to our department with 8 months of progressive painful subcutaneous nodules and abscesses in occipital scalp that caused numerous areas of alopecia (Figure 1). He had received treatment with isotretinoin due to acne conglobata two years back. Skin-biopsy specimen revealed a deep, chronic inflammatory infiltrate affecting the inferior portion of hair follicles. PAS, Grocott and Ziehl-Nielsen stains were negative. Bacterial, mycobacterial and mycotic biopsy cultures were also negative. Thus, dissecting cellulitis of the scalp was diagnosed. This pathology is defined as a chronic suppurative dermatosis of the scalp; it is more frequent in young afro-descendant males; originating from follicular occlusion that causes progressive communicating abscesses that dissect the scalp, resulting in areas of scarring alopecia. Trichoscopy is also an useful diagnostic tool in dissecting cellulitis, showing yellow, structureless areas and yellow dots with "3D" structure imposed over dystrophic hair shafts as most characteristic findings. The patient began treatment with oral isotretinoin 0.7 mg/kg/day and therapy was maintained for 6 months with adequate control of the disease after 2 months of treatment.



Figure 1

\* Corresponding author.

E-mail address: [ale.ortizprieto@hotmail.com](mailto:ale.ortizprieto@hotmail.com) (A. Ortiz-Prieto).