



ACTAS Derma-Sifiliográficas

Full English text available at
www.actasdermo.org



ORIGINAL ARTICLE

Impact on Quality of Life in Dermatology Patients Attending an Emergency Department



A. Alegre-Sánchez^{a,*}, D. de Perosanz-Lobo^a, A. Pascual-Sánchez^b,
C. Pindado-Ortega^a, P. Fonda-Pascual^a, Ó.M. Moreno-Arrones^a, P. Jaén-Olasolo^a

^a Dermatology Service, Ramon y Cajal Hospital, Madrid, Spain

^b Psychiatry Service, Ramon y Cajal Hospital, Madrid, Spain

Received 11 April 2017; accepted 2 July 2017

Available online 28 October 2017

KEYWORDS

Dermatological emergencies;
Quality of life;
Health Related Quality of Life;
DLQI;
Emergencies

Abstract

Introduction: Dermatological complaints have been estimated to represent up to 5–10% of all the visits to emergency departments. The main objective of our study was to determine how affected is the Health related Quality of Life (HRQL) in a series of patients attending an emergency department due to skin symptoms.

Patients and methods: A prospective study during one month (July 2016) was conducted in a hospital with full-time on-call dermatologists. The Short-Form SF-12v2 Health Survey and the Dermatology Life Quality Index (DLQI) were offered to all the patients over 18 years old attending the emergency department with cutaneous complaints. Clinical and epidemiological characteristics were also collected.

Results: In total 108 patients completed the study. Mean age found was 45.1 ± 16.1 years. Mean DLQI score found was 10.56 ± 6.12 . Fifty-three patients (49%) had a score of 11 or higher in the DLQI questionnaire. Most affected subscales were "Symptoms and Feelings" in DLQI scale and "Overall Health" and "Vitality" for the SF-12. A very significant difference ($p < 0.0001$) was found between women's (12.4 ± 5.7) and men's (7.5 ± 5.6) DLQI mean score (mean difference of 4.9; 95% confidence interval of the difference: 2.7–7.1).

Conclusions: Patients visiting emergency units with cutaneous complaints seem to feel a moderate-large impact on their quality of life which is mainly related to the symptoms and feelings that they are experiencing. This impact is significantly higher among women.

© 2017 AEDV. Published by Elsevier España, S.L.U. All rights reserved.

* Corresponding author.

E-mail address: adrian.alegresanchez@gmail.com (A. Alegre-Sánchez).

PALABRAS CLAVE

Urgencias
 dermatológicas;
 Calidad de vida;
 Calidad de vida
 relacionada con la
 salud;
 DLQI;
 Urgencias

Calidad de vida en pacientes dermatológicos valorados en urgencias**Resumen**

Introducción: Los problemas dermatológicos suponen hasta un 5-10% de todas las visitas a los servicios de urgencias. El principal objetivo de nuestro estudio fue determinar la calidad de vida relacionada con la salud en una serie amplia de pacientes que acudieron a un servicio de urgencias por sintomatología cutánea.

Pacientes y métodos: Se realizó un estudio prospectivo de un mes de duración (julio de 2016) en un hospital terciario con dermatólogo de guardia a tiempo completo. La versión corta del Cuestionario de Salud SF-12v2 y del cuestionario DLQI fueron entregados a todos los pacientes mayores de edad que acudieron al servicio de urgencias con problemas dermatológicos. Se recogió información clínica y epidemiológica de todos los pacientes.

Resultados: En total, 108 pacientes completaron el estudio. La edad media de nuestra muestra fue de $45,1 \pm 16,1$ años. La puntuación media del DLQI fue $10,56 \pm 6,12$. Del total, 33 pacientes (49%) tuvieron una puntuación de 11 o más en el DLQI. Las subescalas más afectadas fueron la de «síntomas y sentimientos» para el DLQI y las de «salud general» y «vitalidad» para el SF-12. Se encontró una diferencia muy significativa ($p < 0,0001$) entre las puntuaciones totales del DLQI en mujeres ($12,4 \pm 5,7$) y hombres ($7,5 \pm 5,6$), con una diferencia media de 4,9 (intervalo de confianza al 95%: 2,7-7,1).

Conclusiones: Los pacientes que acuden a los servicios de urgencias con problemas dermatológicos parecen tener una afectación moderada-grave en su calidad de vida, debido principalmente a los síntomas que están experimentando. Este impacto es significativamente mayor en las mujeres.

© 2017 AEDV. Publicado por Elsevier España, S.L.U. Todos los derechos reservados.

Introduction

Dermatological complaints have been estimated to represent up to 5–10% of all the visits to emergency departments.¹ However, the exact incidence is very heterogeneous due to the differences that can be found amongst the methodology of previous studies in the literature regarding their duration, type of patients included or setting (hospital in-patient settings, ambulant settings or emergency departments).¹⁻⁵

Cutaneous diseases might not represent a real threat to patient's life in most cases, but they are frequently felt as very disturbing due to their symptoms, such as itching or pain, or due to the fact that cutaneous lesions are easily observed by the patients. The patient's subjective sensation of alarm is what makes them visit emergency departments seeking for a specialized attention.

Although there are previous studies describing clinical and demographical features of dermatological patients attending emergency departments,¹⁻⁵ none of them has measured the Health Related Quality of Life (HRQL) impact in this population.

The main objective of our study was to determine how affected is the HRQL in patients with dermatological complaints evaluated in an emergency department. Also, we wanted to detect if there were differences in these scores associated with any epidemiological or clinical variable.

Patients and methods

A prospective study was performed during one month (July 2016) in the emergency department of a tertiary hospital

with full-time (24h) on-call dermatologist. In total, nine different dermatology residents attended patients at the emergency department during this period. A standardized form was used to collect epidemiological and clinical information such as age, sex, marital status, level of education, diagnosis and group of diagnosis, previous visits, need of supplementary tests and discharge destination.

After every consultation, patient's self-assessment of QoL was measured using the Spanish validated versions of two self-administered questionnaires: one generic (Short-Form SF-12 Health Survey) and one specific for dermatologic patients (Dermatology Life Quality Index – DLQI). The SF-12v2 is a multidimensional 12-item instrument that includes eight domains: overall health, physical functioning, physical role, emotional role, corporal pain, mental health, vitality and social functioning. The scores in the SF-12 are standardized of 0–100 in which higher scores represent better QoL.⁶ The DLQI measures the impact on the QoL in a patient with a skin problem in the past 7 days.⁷ It has 10 questions categorized into six dimensions: symptoms and feelings (SYF), daily activities (DA), leisure (L), work or school (WS), personal relationships (PR) and treatment (T). Its scores are measured in a range of 0–30 with higher scores meaning higher impairment. This scale has been widely used in the past years in different dermatological diseases.⁶

Both questionnaires were offered to all adult patients visiting the emergency department with any dermatological complaint. Patients with reading or understanding difficulties, such as elderly patients, were excluded from the study. Underage patients (<18 years-old) were also excluded.

All the data were analyzed statistically using the IBM SPSS Statistics software for Windows V22.0. (Armonk,




Figure 1 Flow-chart of the study.

NY: IBM Corp). Kolmogorov–Smirnov test demonstrated that the scores obtained in both DLQI and SF-12v2 tests had a normal distribution. Therefore, the parametric Student's *t*-test was used to analyze differences between the questionnaire scores depending on variables with two categories (i.e. sex) and the analysis of variance (ANOVA) was used for those variables with more than two categories (i.e. marital status). In those cases with groups smaller than 30 patients, non-parametric Mann–Whitney *U* test and Kruskal–Wallis test were used. A *p*-value of less than 0.05 was considered as statistically significant.

Results

In total, 245 patients were attended by dermatologists at the emergency department during the period of the study (mean of 8.17 patients per day). This figure represents the 3.7% of the total number of patients attended at the emergency department of the hospital on that period (6621 patients). In all, 108 patients (42.5%) met the inclusion criteria and completed both questionnaires (flow-chart of the study is represented in Fig. 1). Mean age of the studied population was 45.1 years (standard deviation: 16.1; range 18–75 years). Overall, the most common diagnoses made were urticaria (17 patients, 16%), eczema (16 patients, 14%) and infectious cellulitis (7 patients, 6%). Attending to groups of diagnoses, 69 patients (64%) had an inflammatory dermatosis; 33 patients (31%) had an infectious dermatosis and the rest of the patients (6 patients, 5%) had either a tumoral

dermatosis or a post-surgical complication. The rest of the demographical and clinical features of the studied population can be found in Table 1.

The mean DLQI score found was 10.56 (standard deviation 6.12; range 0–25). All DLQI total scores and subscale scores attending to clinical and demographical variables are detailed in Table 2. Fifty-three patients (49%) had a score of 11 or higher in the DLQI questionnaire, what is categorized as a “very large/extremely large effect on HRQL”. The number of patients on each category of the DLQI is represented in Table 3. A very significant difference ($p < 0.0001$) was found between women's (12.4 ± 5.7) and men's (7.5 ± 5.6) DLQI mean score (mean difference of 4.9; 95% confidence interval of the difference: 2.7–7.1). This difference, nearly 5, is also greater to what is considered as the Minimum Clinical Important score Difference (4 points) for the DLQI. Attending to the different subscales of the DLQI depending on sex, all of the subscales except the “Treatment” item showed a statistically significant higher impairment among female patients.

No significant differences were found in total DLQI scores associated with age, marital status, education level, diagnosis or any other variable. Significant differences were found on the “Personal relationship” subscale depending on age ($p = 0.04$) and “Treatment” subscale depending on marital status ($p = 0.04$).

Regarding SF-12 scores, global subscale scores can be found in Table 4. The most affected dimensions in SF-12 scale were “Overall Health” (mean score of 53.47 out of 100) and “Vitality” (mean score of 52.78). Significant differences were only found in the “Social Functioning” SF-12

Table 1 Clinical and demographical features of the patients included in the study.

Characteristic	Patients (n = 108); n (%)
Sex	
Male	41 (38%)
Female	67 (62%)
Age (years) 45.1 ± 16.1, Range: 18–75 y.	
<31	24 (22%)
31–55	54 (50%)
>55	30 (28%)
Educational level	
Elementary/secondary	26 (24%)
Professional/Universitary	82 (76%)
Marital status	
Married or living with partner	50 (46%)
Living without partner	58 (54%)
Previous visits due to similar complaints	
Yes	25 (23%)
No	83 (77%)
Supplementary tests (blood test, biopsy, imaging, etc.)	
Yes	35 (32%)
No	73 (68%)
Discharge destination	
Home	70 (65%)
General physician/dermatology clinics	38 (35%)
Diagnosis	
Inflammatory disease	69 (64%)
Infectious disease	33 (31%)
Others (tumoral, external agents, etc.)	6 (5%)

subscale depending on age, with lower scores in patients over 60 years old ($p=0.032$). There were also significantly lower scores in ‘‘Physical Functioning’’ and ‘‘Vitality’’ items among patients with lower education levels ($p=0.01$, $p=0.04$ respectively).

Discussion

Dermatologic diseases are rarely real life-threatening medical emergencies. However, they represent a significant percentage of the visits to emergency departments. Understanding how affected is the QoL in these patients due to the symptoms that they are experiencing is a necessary duty in order to better assess their medical problems.

The presence of an on-call dermatologist is not a constant fact in most of the hospitals in our country and general doctors attend most of the urgent cutaneous complaints. This is probably due to the general feeling among many non-dermatologist doctors that consider the skin diseases as

‘‘minor’’ complaints compared to other diseases attended at emergency departments.

Interestingly, the score found in our study suggests that patients visiting emergency departments with dermatological complaints have a very significant impact in their HRQL, with a mean score in the DLQI of 10.56, considered between the categories of ‘‘moderate effect (5–10)’’ and ‘‘very large effect (11–20)’’. In fact, 49% of the patients had a DLQI score higher than 11. This might reflect the fact that patients usually have a lot of concern when they are experiencing cutaneous acute diseases, although most of them are not considered as real medical emergencies.

Compared to the scores in usual cutaneous diseases, the mean DLQI total score in our study would be similar to that found in other studies in severe psoriasis,⁷ atopic dermatitis or pemphigus vulgaris⁸ and higher than what have been estimated for pathological scars,⁸ acne⁶ or vitiligo.⁹ Even though our study includes patients with different diseases, this comparison can help us to understand how intense might be the impairment that they feel.

Another interesting point is the higher impact found in female patients compared to males, of almost 5 points in total DLQI scores. The only subscale that showed no significant difference depending on sex was the ‘‘Treatment’’ item, probably because most of the patients included were not using any treatment at the moment of the visit. There are previous studies assessing HRQL in Dermatology that have found greater impact among women, specially those studies studying diseases with important esthetical impairment such as alopecias.¹⁰ The differences found in our study are high enough to be considered as clinically relevant. Therefore, we can assume that overall, women experience dermatological acute symptoms with a much greater concern than males. We hypothesize whether these results are due to the more importance that women usually give to the appearance of their skin. Other possibility might be that women are more concerned about their own health. In any case, dermatologists should have this in consideration when attending this population.

Overall, the most affected subscale of the DLQI is the ‘‘Symptoms and Feeling’’ (SYF) dimension, with a total mean score of 3.28 (± 1.71). Therefore, we can assume that the main triggers that make these patients attend emergency departments are the symptoms and the subjective concern that they feel.

A limitation of our study was that a relatively high proportion of the patients were excluded due to their age or because of reading or understanding disabilities (87 patients, 35.5% of the total population). Also a percentage of the patients included initially did not complete the questionnaires (31.6%). Another possible limitation might have been the possible impact on the HRQL due to medical or psychological comorbidities of the patients included which are sometimes associated with dermatological diseases (i.e. depression, anxiety, etc.). Also, we suggest that further studies comparing HRQL in patients with dermatological emergencies and other type of medical emergencies could provide interesting information.

Table 2 Association between DLQI scores (global and subscales) and clinical variables of dermatology patients attending the emergency department.

Characteristic	No	DLQI						
		SYF	DA	L	WS	PR	T	Total
Sex								
Male	41	2.41 ± 1.60	1.27 ± 1.48	1.73 ± 1.64	0.80 ± 0.98	0.90 ± 1.32	0.39 ± 0.70	7.52 ± 5.61
Female	67	3.81 ± 1.57	2.64 ± 1.81	2.46 ± 1.81	1.28 ± 1.07	1.54 ± 1.49	0.70 ± 0.97	12.41 ± 5.72
<i>p</i> -Value		<0.0001	<0.0001	0.04	0.02	0.03	0.08	<0.0001
Age								
<31	24	3.33 ± 1.63	2.21 ± 1.96	2.17 ± 1.97	1.67 ± 1.01	1.08 ± 1.28	0.54 ± 0.83	10.52 ± 5.53
31–55	54	3.39 ± 1.74	2.26 ± 1.92	2.39 ± 1.75	1.22 ± 1.09	1.67 ± 1.58	0.69 ± 0.97	11.61 ± 6.51
>55	30	3.03 ± 1.75	1.80 ± 1.47	1.83 ± 1.66	0.83 ± 1.02	0.80 ± 1.86	0.43 ± 0.77	8.71 ± 5.63
<i>p</i> -Value		0.59	0.57	0.29	0.24	0.04	0.60	0.67
Educational level								
Elementary/high school	26	3.04 ± 1.73	2.46 ± 1.73	2.04 ± 1.56	1.08 ± 0.93	1.77 ± 1.68	0.61 ± 0.85	11.01 ± 6.02
Professional/University	82	3.35 ± 1.71	2.01 ± 1.84	2.23 ± 1.85	1.11 ± 1.10	1.15 ± 1.35	0.57 ± 0.90	10.42 ± 6.23
<i>p</i> -Value		0.38	0.23	0.81	0.96	0.12	0.67	0.30
Marital status								
Married	50	3.16 ± 1.81	2.27 ± 1.89	2.00 ± 1.77	0.90 ± 1.10	1.14 ± 1.41	0.41 ± 0.76	10.71 ± 5.72
Single/divorced/other	58	3.31 ± 1.61	1.82 ± 1.67	2.33 ± 1.78	1.22 ± 1.05	1.37 ± 1.44	0.76 ± 0.99	10.39 ± 6.48
<i>p</i> -Value		0.65	0.19	0.89	0.12	0.41	0.04	0.80
Previous visits								
Yes	25	3.32 ± 1.82	2.56 ± 1.76	2.68 ± 1.86	1.04 ± 1.06	1.28 ± 1.24	0.72 ± 1.06	11.63 ± 5.41
No	83	3.27 ± 1.69	1.99 ± 1.82	2.04 ± 1.74	1.12 ± 1.07	1.31 ± 1.52	0.54 ± 0.83	10.19 ± 6.31
<i>p</i> -Value		0.90	0.17	0.12	0.64	0.93	0.06	0.32
Supplementary tests								
Yes	35	3.00 ± 1.93	1.86 ± 1.93	2.03 ± 1.95	1.00 ± 1.00	1.00 ± 1.31	0.69 ± 0.99	9.71 ± 6.59
No	73	3.41 ± 1.60	2.25 ± 1.75	2.26 ± 1.70	1.15 ± 1.09	1.44 ± 1.51	0.53 ± 0.83	11.1 ± 5.89
<i>p</i> -Value		0.22	0.29	0.53	0.49	0.14	0.38	0.30
Discharge destination								
Home	70	3.14 ± 1.81	2.08 ± 1.82	2.21 ± 1.85	0.95 ± 1.01	1.22 ± 1.35	0.67 ± 0.97	10.59 ± 5.78
General physician/dermatology clinics	38	3.47 ± 1.56	2.18 ± 1.82	2.16 ± 1.69	1.31 ± 1.10	1.40 ± 1.60	0.47 ± 0.76	11.11 ± 6.09
<i>p</i> -Value		0.34	0.78	0.89	0.09	0.54	0.27	0.41
Type of diagnosis								
Inflammatory disease	69	3.55 ± 1.73	2.06 ± 1.84	2.21 ± 1.79	1.15 ± 1.02	1.19 ± 1.36	0.57 ± 0.86	11.12 ± 6.12
Infectious	33	2.2 ± 1.53	2.32 ± 1.81	2.27 ± 1.74	1.11 ± 1.15	1.54 ± 1.61	0.57 ± 0.90	9.81 ± 6.21
<i>p</i> -Value		0.07	0.50	0.87	0.86	0.25	1.00	0.60
Total		3.28 ± 1.71	2.12 ± 1.81	2.18 ± 1.77	1.10 ± 1.06	1.30 ± 1.45	0.58 ± 0.89	10.56 ± 6.12

SYF, symptoms and feelings; DA, daily activities; L, leisure; WS, work and school; PR, personal relationship; T, treatment; S, symptoms; E, emotions; SF, social functioning. Bold figures serve to highlight statistically significant values.

Patients visiting emergency units with cutaneous complaints seem to experience a great impact on their quality of life related to the symptoms and feelings that they are living. This situation reflects the importance of attending these patients in an expeditious way, giving a correct diagnosis and a prompt treatment; a duty that can only be optimally assessed by a dermatologist.

We consider that the results of this study firmly support the presence of on-call dermatologists to better attend dermatological patients at emergency departments. We think that it is necessary to take into account the real impact of the diseases on each patient's quality of life in order to consider them as "urgent" or "non-urgent" problems.

Table 3 DLQI score groups and number of patients in each group.

DLQI scores groups and interpretation	Number of patients in our study in each group
0–1 = no effect at all on patient's life	8 (7%)
2–5 = small effect on patient's life	17 (16%)
6–10 = moderate effect on patient's life	30 (28%)
11–20 = very large effect on patient's life	47 (44%)
21–30 = extremely large effect on patient's life	6 (5%)

Table 4 SF 12 mean scores in each dimension of the test.

SF-12 dimensions	Mean \pm SD (0–100)
Overall health	53.47 \pm 22.01
Physical functioning	73.15 \pm 29.54
Physical role	63.43 \pm 42.47
Emotional role	76.67 \pm 38.02
Corporal pain	65.74 \pm 30.87
Mental health	67.69 \pm 20.89
Vitality	52.78 \pm 27.13
Social functioning	84.03 \pm 22.27

Ethical disclosures

Protection of human and animal subjects. The authors declare that no experiments were performed on humans or animals for this study.

Confidentiality of data. The authors declare that no patient data appear in this article.

Right to privacy and informed consent. The authors declare that no patient data appear in this article.

Conflict of interests

The authors declare no conflict of interests.

References

1. Grillo E, Vaño-Galván S, Jiménez-Gómez N, Ballester A, Muñoz-Zato E, Jaén P. Dermatologic emergencies: descriptive analysis of 861 patients in a tertiary care teaching hospital. *Actas Dermosifiliogr.* 2013;104:316–24.
2. Bancalari-Díaz D, Gimeno-Mateos LI, Cañueto J, Andrés-Ramos I, Fernández-López E, Román-Curto C. Dermatologic emergencies in a tertiary hospital: a descriptive study. *Actas Dermosifiliogr.* 2016;107:666–73, <http://dx.doi.org/10.1016/j.ad.2016.05.001>
3. Mirkamali A, Ingen-Housz-Oro S, Valeyrie Allanore L, Bouvrese S, Duong TA, Chosidow O, et al. Dermatological emergencies: a comparative study of activity in 2000 and 2010. *J. Eur. Acad. Dermatology Venereol.* 2013;27:916–8.
4. Wakosa A, Roussel A, Delaplace M, Le Bidre E, Binois R, Valéry A, et al. Intérêt d'une consultation dermatologique d'urgence dans un centre hospitalier régional. *Presse Med.* 2013;42:e409–15.
5. Ruzza N, Itin PH, Beltraminelli H. Urgent consultations at the dermatology department of Basel University Hospital, Switzerland: characterisation of patients and setting – a 12-month study with 2,222 patients data and review of the literature. *Dermatology.* 2014;228:177–82.
6. Ware JJ, Kosinski MM, Keller SSD. A 12-Item Short-Form Health Survey: construction of scales and preliminary tests of reliability and validity. *Med Care.* 1996;34:220–33.
7. Finlay AY, Khan GK. Dermatology Life Quality Index (DLQI) – a simple practical measure for routine clinical use. *Clin Exp Dermatol.* 1994;19:210–6.
8. Basra MKA, Fenech R, Gatt RM, Salek MS, Finlay AY. The Dermatology Life Quality Index 1994–2007: a comprehensive review of validation data and clinical results. *Br J Dermatol.* 2008;159:997–1035.
9. Amer AAA, Gao XH. Quality of life in patients with vitiligo: an analysis of the dermatology life quality index outcome over the past two decades. *Int J Dermatol.* 2016;55:608–14.
10. Janković S, Perić J, Maksimović N, Ćirković A, Marinković J, Janković J, et al. Quality of life in patients with alopecia areata: a hospital-based cross-sectional study. *J Eur Acad Dermatol Venereol.* 2016;30:840–6.