




ACTAS Derma-Sifiliográficas

Full English text available at
www.actasdermo.org



CASE AND RESEARCH LETTERS

Nodule on the External Ear[☆]



Nódulo en el pabellón auricular

Medical History

A 30-year-old man with no relevant past medical history presented with a lesion on the lower helix of the right ear that had first appeared several months earlier. The lesion was asymptomatic and its growth had remained stable since it first appeared. The patient reported no history of trauma at the site and had no similar lesions at other sites.

Physical Examination

Physical examination revealed a soft, raised, well-defined, erythematous-violaceous tumor approximately 1 cm in diameter that had a smooth surface and was not painful to the touch (Fig. 1).

Histopathology

Hematoxylin-eosin staining (Figs. 2A and B) revealed a dermal lymphoid infiltrate composed of small to medium-sized lymphocytes with little atypia, no mitotic figures, and no phenomena of avascular necrosis or epidermotropism. In addition, a grenz zone separating the lesion from the dermis—which was completely spared—was also observed.



Figure 1




Figure 2 A, Hematoxylin-eosin, original magnification $\times 40$. B, Hematoxylin-eosin, original magnification $\times 100$.




Figure 3

Immunohistochemical staining was clearly positive for the CD3 and CD8 markers, partially positive for CD4, and negative for CD20 and CD30 (Fig. 3).

Additional Tests

A blood workup revealed no alterations and a computed tomography study of the abdominal and pelvic region was entirely normal.

What Is Your Diagnosis?

[☆] Please cite this article as: Heras-González S, Aspe-Unanue L, González-Pérez R. Nódulo en el pabellón auricular. Actas Dermosifiliogr. 2017;108:581–582.

Diagnosis

Indolent CD8+ lymphoid proliferation of the ear.

Clinical Course and Treatment

Complete excision of the lesion was performed, followed by closure by plasty. No recurrence was detected during 2 years of follow-up.

Comment

Indolent CD8+ lymphoid proliferation of the ear, first described by Petrella et al. in 2007, is a rare lymphoproliferative process that is not considered to be a cutaneous T-cell lymphoma according to the 2005 classification of the World Health Organization–European Organization for Research and Treatment of Cancer. Although it presents immunophenotypic characteristics also seen in aggressive CD8+ T-cell lymphomas, this new indolent lymphoproliferative process is probably a phenotypic variant of primary cutaneous CD4+ small-medium pleomorphic T-cell lymphoma, as first suggested by Beltraminelli et al. in 2009.^{1,2} Clinically, the entity is characterized by slow-growing lesions confined to a single cutaneous site, usually the ears or acral sites. In the cases reported to date, the entity has exhibited indolent clinical behavior and shown no signs of systemic involvement.¹⁻³

Cutaneous CD8+ T-cell lymphomas were initially considered to be aggressive lymphomas with frequent extracutaneous involvement and poor prognosis. Subsequently, CD8+ lymphomas with a favorable prognosis—which did not match the original descriptions—were reported. In 2007, Petrella et al.³ reported for the first time 4 cases of a nonaggressive, nonepidermotropic, monoclonal, CD8+ lymphoid proliferation on the ear that did not appear to belong to any of the recognized types of cutaneous T-cell lymphoma. Other cases with the same clinical, histologic, and immunophenotypic characteristics were subsequently reported, prompting various authors to suggest that this new entity might be a new phenotypic variant of primary cutaneous CD4+ small-medium pleomorphic T-cell lymphoma.^{1,2,4-6}

Physical examination of these indolent cases usually reveals an asymptomatic, slow-growing, erythematous nodule on the helix of the ear measuring approximately 1 cm in diameter.^{1,2} Clinical differential diagnosis of this entity includes conditions such as cutaneous pseudolymphoma (benign lymphocytoma cutis) caused by *Borrelia burgdorferi* infection or by ear piercing. Histopathologic differential diagnosis includes benign lymphocytoma cutis, reactions to insect bites, discoid lupus erythematosus, panniculitis-like subcutaneous lymphoma, polymorphous light eruption, and angiolymphoid hyperplasia with eosinophilia, and less frequently primary cutaneous B-cell lymphoma. The CD8+/CD4-immunophenotype can be seen in CD8+ mycosis fungoides,

pagetoid reticulosis, aggressive CD8+ primary lymphoma, prolymphocytic variants of leukemia, CD8+ lymphomatoid papulosis, and other conditions.¹

Histologically, the entity is characterized by a dense dermal infiltrate of CD8+ T cells with a lymphoblast-like appearance, T-cell clonality, and no epidermotropism. The lymphoid infiltrate is composed of medium-sized pleomorphic cells with irregular nuclei and scant cytoplasm. There is no evidence of necrosis, mitosis, ulceration, or angiodestruction.⁶ Immunohistochemical staining is consistently positive for the CD3 and CD8 markers, while there is little expression of CD20, CD4, CD30, and CD56.⁴⁻⁶

Although in some cases the histopathologic appearance suggests a high-grade lymphoma, the clinical behavior of the entity is benign and therefore it is essential to differentiate it from other cutaneous CD8+ lymphomas that are associated with a poorer prognosis in order to avoid aggressive and unnecessary treatments.^{2,3}

Conflicts of Interest

The authors declare that they have no conflicts of interest.

References

- Swick BL, Baum CL, Venkat AP, Liu V. Indolent CD8+ lymphoid proliferation of the ear: Report of two cases and review of the literature. *J Cutan Pathol*. 2011;38:209–15.
- Beltraminelli H, Müllegger R, Cerroni L. Indolent CD8+ lymphoid proliferation of the ear: A phenotypic variant of the small-medium pleomorphic cutaneous T-cell lymphoma? *J Cutan Pathol*. 2010;37:81–4.
- Petrella T, Maubec E, Cornillet-Lefebvre P, Willemze R, Pluot M, Durlach A, et al. Indolent CD8-positive lymphoid proliferation of the ear: a distinct primary cutaneous T-cell lymphoma. *Am J Surg Pathol*. 2007;31:1887–92.
- Zeng W, Nava VE, Cohen P, Ozdemirli M. Indolent CD8-positive T-cell lymphoid proliferation of the ear: A report of two cases. *J Cutan Pathol*. 2012;39:696–700.
- Suchak R, O'Connor S, McNamara C, Robson A. Indolent CD8-positive lymphoid proliferation on the face: part of the spectrum of primary cutaneous small-/medium-sized pleomorphic T-cell lymphoma or a distinct entity. *J Cutan Pathol*. 2010;37:977–81.
- Greenblatt D, Ally M, Child F, Scarisbrick J, Whittaker S, Morris S, et al. Indolent CD8+ lymphoid proliferation of acral sites: A clinicopathologic study of six patients with some atypical features. *J Cutan Pathol*. 2013;40:248–58.

S. Heras-González,* L. Aspe-Unanue, R. González-Pérez

Servicio de Dermatología, Hospital Universitario Araba, Vitoria-Gasteiz, Spain

* Corresponding author.

E-mail address: soniaherasgonzalez@gmail.com

(S. Heras-González).