

## Mycosis Fungoides in Pediatric Patients: A Diagnostic Challenge<sup>☆</sup>



### Micosis fungoide en pacientes pediátricos: un reto diagnóstico

Fortunately, few cases of primary cutaneous lymphoma and mycosis fungoides (MF) are seen in pediatric patients. In my experience, as head of the Lymphoma Unit at Hospital Universitario Basurto, in Bilbao, Spain, I have seen just 2 cases of MF, both classic, with an onset before the age of 18 years. The course of disease was more aggressive than usual in both patients. One of the patients died at the age of 33 years following allogeneic hematopoietic progenitor cell transplantation and the other developed tumor-stage MF (IIB).

<sup>☆</sup> Please cite this article as: Beloso RI. Micosis fungoide en pacientes pediátricos: un reto diagnóstico. *Actas Dermosifiliogr.* 2017;108:499.

In this issue of *Actas Dermo-Sifiliográficas*, Bettina Cervini A et al. present 14 cases of MF, most of which were hypopigmented, from a pediatric hospital. I recently had the opportunity to see a similar case in an immigrant girl who had been diagnosed in Columbia. Hypopigmented MF is more common in patients with dark skin (Fitzpatrick skin types IV-V), as evidenced by the literature, and we may therefore see an increasing number of cases in our setting. The main entity that should be contemplated in the differential diagnosis is pityriasis alba. This is a very common condition in pediatric settings, but the possibility of hypopigmented MF needs to be considered in cases that are refractory to treatment or that do not follow a typical course.

Reading this article will help us to do this.

Rosa Izu Beloso\*

*Servicio de Dermatología, Hospital Universitario de Bilbao, Bilbao, Vizcaya, Spain*

\* Corresponding author.

E-mail address: [ROSAMARIA.IZUBELLOSO@osakidetza.eus](mailto:ROSAMARIA.IZUBELLOSO@osakidetza.eus)  
1578-2190/

© 2017 Elsevier España, S.L.U. and AEDV. All rights reserved.

## Eosinophils as a Warning Sign<sup>☆</sup>



### El eosinófilo como alerta

Eosinophils are effector cells of the innate immune system that play a key role in the body's defense against parasites and tumors. Their main function is to eliminate biological targets, such as helminths, protozoa, bacteria, and tumor cells. The pathological circumstances that trigger an increase in the number of eosinophils in tissues or peripheral blood are very varied. The literature includes very few review articles dealing specifically with eosinophilic dermatoses, and attempts to draw up a preliminary classification are even rarer.<sup>1</sup>

In 2001, Byrd et al.<sup>2</sup> coined the term *eosinophilic dermatosis of myeloproliferative disease*, for which they proposed the following clinical and pathological diagnostic criteria: a pruritic papular or vesiculobullous eruption refractory to standard treatment; a concurrent diagnosis of blood dyscrasia or hematologic malignancy; an eosinophil-rich lymphohistiocytic infiltrate in the superficial and deep dermis; and the exclusion of other causes of tissue eosinophilia. This description corresponds to the dermatosis originally described in 1965 by Weed<sup>3</sup> as a *mosquito bite-like reaction*. More recently, a new nomenclature has been proposed: *eosinophilic dermatosis of hematologic malignancy*.<sup>4</sup> It is a reactive process without specific hematologic cell infiltration. The etiology and pathogenesis is poorly understood,

but it is thought to be associated with the immune dysregulation found in conditions such as chronic lymphocytic leukemia.

Our journal has already published a small case series of this skin condition.<sup>5</sup> In this issue of *ACTAS DERMOSIFILIOGRÁFICAS*, Lucas-Truyols et al.<sup>6</sup> review 4 new typical examples and, for the first time, 1 case associated with mycosis fungoides. The great interest of this condition, and all skin disorders linked to systemic diseases, is that its prompt recognition may have diagnostic and, in some cases, prognostic value with regard to the underlying hematologic disease.

## References

- Rodríguez-Díaz E, Álvarez-Cuesta C, Blanco Barrios S, Galache Osuna C, Requena Caballero C. Dermatitis eosinofílicas (y II). *Actas Dermosifiliogr.* 2003;94:131–43.
- Byrd JA, Scherschun L, Chaffins ML, Fivenson DP. Eosinophilic dermatosis of myeloproliferative disease: Characterization of a unique eruption in patients with hematologic disorders. *Arch Dermatol.* 2001;137:1368–80.
- Weed RI. Exaggerated delayed hypersensitivity to mosquito bites in chronic lymphocytic leukemia. *Blood.* 1965;26:257–68.
- Farber MJ, Forgia SL, Sahu J, Lee JB. Eosinophilic dermatosis of hematologic malignancy. *J Cutan Pathol.* 2012;39:690–5.
- Mangas C, Bielsa I, Mate JL, Fernández-Figueras M<sup>ª</sup>T, Ribera M, Ferrándiz C. Dermatitis eosinofílica asociada a leucemia linfática crónica: estudio clínico, microscópico e inmunohistoquímico de 5 casos. *Actas Dermosifiliogr.* 2004;95:165–70.
- Lucas-Truyols S, Rodrigo-Nicolás B, Lloret-Ruiz C, Quecedo-Estébanez E. Cuatro nuevos casos de dermatosis eosinofílicas asociadas a procesos hematológicos. *Actas Dermosifiliogr.* 2017. Doi: <http://dx.doi.org/10.1016/j.ad.2016.10.021>

<sup>☆</sup> Please cite this article as: Rodríguez E. El eosinófilo como alerta. *Actas Dermosifiliogr.* 2017;108:499–500.