



Revista Brasileira de Hematologia e Hemoterapia
Brazilian Journal of Hematology and Hemotherapy

www.rbhh.org



Original article

Telomere length correlates with disease severity
and inflammation in sickle cell disease



Marina Pereira Colella^a, Barbara A. Santana^{b,c}, Nicola Conran^a, Vinicius Tomazini^a,
Fernando F. Costa^a, Rodrigo T. Calado^{b,c}, Sara T. Olalla Saad^{a,*}

^a Universidade Estadual de Campinas/Hemocentro (UNICAMP), Campinas, SP, Brazil

^b Universidade de São Paulo (USP), Ribeirão Preto, SP, Brazil

^c Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP), Ribeirão Preto, São Paulo, SP, Brazil

ARTICLE INFO

Article history:

Received 17 January 2017

Accepted 15 February 2017

Available online 11 March 2017

Keywords:

Sickle cell disease

Telomere length

Inflammation

ABSTRACT

Background: Telomeres, the ends of linear chromosomes, shorten during mitotic cell division and erosion may be aggravated by inflammation or proliferative and oxidative stress. As the bone marrow is under hyperproliferative pressure in sickle cell disease and several tissues are submitted to chronic inflammation, this study sought to determine the telomere length of patients with sickle cell disease.

Methods: The mean telomere length was measured in peripheral blood leukocytes by quantitative polymerase chain reaction. The age-adjusted telomere to single copy gene ratio was compared between 91 adult sickle cell disease patients and 188 controls.

Results: Sickle cell disease patients had significantly shorter telomeres than the controls (p -value < 0.0001). Moreover, among sickle cell disease genotypes, Hb SS patients had significantly shorter telomeres compared to Hb SC and Hb S β patients (p -value < 0.0001). Patients on hydroxyurea also had shorter telomeres in comparison to those off the drug (p -value = 0.02). A positive correlation was observed between telomere length and hemoglobin level (r = 0.3; p -value = 0.004), whereas negative correlations were detected between telomere length and lymphocyte count (r = -0.3; p -value = 0.005) and interleukin-8 serum levels (r = -0.4; p -value = 0.02).

Conclusions: The findings of this study indicate that telomeres are short in sickle cell disease patients and that telomere erosion directly correlates with disease genotype, inflammation markers, and the use of hydroxyurea.

© 2017 Associação Brasileira de Hematologia, Hemoterapia e Terapia Celular. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

* Corresponding author at: Hematology and Hemotherapy Center, Universidade Estadual de Campinas (UNICAMP), Rua Carlos Chagas, 480, 13083-878 Campinas, SP, Brazil.

E-mail address: sara@unicamp.br (S.T. Saad).

<http://dx.doi.org/10.1016/j.bjhh.2017.02.007>

1516-8484/© 2017 Associação Brasileira de Hematologia, Hemoterapia e Terapia Celular. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Table 1 – Clinical characteristics of sickle cell disease patients.

	Hb SS	Hb SC/Hb S β ⁺
Number of patients – n	51	40
Mean age – years (range)	37 (18–58)	41 (19–66)
Gender: male/female – n (%)	18/33 (35/65)	14/26 (35/65)
α -Thalassemia trait – n (%)	11 (23)	11 (32)
Use of hydroxyurea – n (%)	37 (73)	3 (8)
Mean age-adjusted T/S – ratio (range)	–0.34 (–0.61 to 0.21)	–0.21 (–0.51 to 0.68)
T/S: telomere to single copy gene ratio.		
* α -Thalassemia trait: –3.7 kb deletion (– α 3.7).		
The results were available for 47 patients in the HbSS group and 34 patients in the HbSC/HbS β group.		

variations (CV) of triplicate measurements were 2% and 1% or less, for telomere and single gene reactions, respectively. The considered inter-assay CV was up to 5%.

Leukocyte telomere length by Southern blot analysis

To validate the qPCR terminal restriction fragment (TRF), analysis was performed by Southern blot of 17 samples according to the manufacturer's instructions with minor changes (TeloTAGGG Telomere Length Assay – Roche Applied Science, Mannheim, Germany). Briefly, 800 ng of genomic DNA was digested by an optimized mixture of *HinfI* and *RsaI* FastDigest restriction enzymes (Thermo Scientific, Waltham, MA, USA) at 37 °C for 2 h. Following DNA digestion, DNA fragments were separated by electrophoresis in 0.8% agarose gel during four hours at 80V. Gel was denatured and neutralized, samples were transferred to a nylon membrane by Southern blotting and probed, and the terminal restriction fragments were detected by chemiluminescence. Mean TRF length was determined according to the formula $TRF = \Sigma(ODi)/\Sigma(ODi/Li)$, where ODi is the chemiluminescent signal and Li is the length of the fragment at a given position.

Inflammation markers

Tumor necrosis factor-alpha (TNF- α) and IL-8 were assessed as pro-inflammatory markers. TNF- α and IL-8 levels were measured, in duplicate, in serum samples using ultra-sensitive enzyme-linked immunosorbent assay (ELISA) kits according to the manufacturer's instructions (TNF- α US and IL-8 US, Invitrogen, Camarillo, CA, USA).

Statistical analyses

The relationship between telomere length and age was estimated using linear regression. Estimated regression coefficients were used to calculate the observed minus expected (O-E), or age-adjusted telomere length for each subject. Age-adjusted T/S ratios were used for all the study analyses except for the correlation with age. Spearman's rank correlation coefficient was used to analyze bivariate associations between telomere lengths, hemolysis and inflammation markers. T/S ratios were compared according to the diagnosis, use of hydroxyurea and gender using Wilcoxon rank sum test. Linear regression was used to analyze the correlation between

telomere length measured by qPCR and TRF by Southern blot. All *p*-values ≤ 0.5 were considered significant. Statistical analyses were performed using the R statistics program version 3.1.3.

Results

Patients' characteristics

Ninety-one adult SCD patients were included in the study: 51 Hb SS, 38 Hb SC, and 2 Hb S β ⁺. Forty were on hydroxyurea with a median dose of 1 g/day (range: 500–1750 mg/day). The dose was titrated according to clinical improvement or to the maximum tolerated dose. The clinical indications for hydroxyurea were frequent painful crises (*n*=14; 35%), acute chest syndrome (*n*=5; 13%), stroke (*n*=6; 16%), and severe hemolytic anemia (*n*=15; 37%). Patients' clinical characteristics are described in Table 1. The control group consisted of 188 healthy individuals (Hb AA), with a mean age of 38 years (range: 18–88 years), including 97 women and 91 men.

Telomeres are short in sickle cell disease patients

Telomeres were significantly shorter in SCD patients compared to age-matched healthy controls [T/S ratio: –0.28 in SCD vs. –0.01 in controls; standard deviation (SD)=0.20 vs. 0.23; *p*-value<0.0001 – Figure 1A]. When genotypes were compared, the telomeres were significantly shorter in Hb SS patients than in the Hb SC and Hb S β genotypes (T/S ratio: –0.34 vs. –0.21, respectively; SD=0.17 vs. 0.21; *p*-value<0.0001 – Figure 1B). Telomeres were also shorter in patients on hydroxyurea (T/S ratio: –0.34 vs. –0.25; SD=0.20 vs. 0.20; *p*-value=0.02) (Figure 1C). Although telomeres shortened with age in healthy controls (*r*=–0.5; *p*-value<0.0001), age did not affect telomere length in SCD patients (*r*=–0.02; *p*-value=0.8). Telomere length was not influenced by gender in either SCD patients (*p*-value=0.2) or controls (*p*-value=0.9).

In order to validate these findings, telomere lengths were also measured by Southern blot for 17 patients showing that T/S ratios and TRFs were highly correlated ($R^2=0.86$; *p*-value<0.0001 – Figure 2) supporting the accuracy of the qPCR results.




Figure 1 – Age-adjusted T/S ratio in controls and sickle cell disease (SCD) patients. (A) SCD patients presented shortened telomere length compared to normal controls. (B) Hb SS patients presented shortened telomere length compared to Hb SC and Hb S β patients. (C) Hb SS and Hb SC patients treated with hydroxyurea (Hb SS/Hb SC HU) presented shortened telomere lengths when compared to patients not treated with hydroxyurea (Hb SS/Hb SC/Hb S β *). T/S ratios were compared according to the diagnosis and use of hydroxyurea using Wilcoxon rank sum test. Mean telomere length was measured in peripheral blood leukocytes by quantitative polymerase chain reaction (qPCR).

Telomere length correlates with hemolysis and inflammation in sickle cell disease

The association between telomere length and marker levels for hemolysis (hemoglobin, hematocrit, lactate dehydrogenase, indirect bilirubin, reticulocyte counts) and inflammation (IL-8, TNF- α , total leukocyte, neutrophil, lymphocyte and monocyte counts) were analyzed (Table 2). There was a weak positive correlation between telomere length and hemoglobin concentration ($r = 0.3$; p -value = 0.004), but no correlation with other hemolysis markers. Among inflammatory markers, there was a weak negative correlation of telomere length with lymphocyte counts ($r = -0.3$; p -value = 0.005) and with IL-8 serum levels ($r = -0.4$; p -value = 0.02; Table 2). Of note, overall, SCD patients had higher lymphocyte counts (SCD: $2.9 \times 10^9/L$

vs. controls: $1.9 \times 10^9/L$; SD = 1.4 vs. 0.51; p -value < 0.0001) and IL-8 levels compared to controls (SCD: 3.3 pg/mL vs. controls: 2.3 pg/mL; SD = 1.9 vs. 0.8; p -value = 0.009). Telomere length was not associated with the plasma levels of other inflammatory markers.

Discussion

In the present study, we found that telomeres of peripheral blood leukocytes of patients with SCD are short independent of age and telomere attrition was more pronounced in patients with Hb SS compared to Hb SC or Hb S β ⁺ genotypes. Telomere length correlated with hemoglobin concentration and inversely correlated with IL-8 level and absolute lymphocyte count. Taken together, these findings suggest that telomere length is associated with disease severity and chronic inflammation in SCD.

Our results are in sharp contrast with one single previous report by Drasar et al., who found SCD patients had longer telomeres in comparison to age-matched controls.¹⁴ However, their study presented some technical issues precluding appropriate interpretation of their findings. First, it is not clear in their work whether they used fresh or frozen samples for DNA extraction. Second, telomere length was highly heterogeneous among their patients, and a significant proportion of older patients had telomeres longer than expected for healthy cord bloods. Finally, their qPCR results were not validated with a second method. Taken together, these concerns might suggest that the DNA used for analysis could have been degraded. During qPCR, DNA degradation causes T/S ratios to be higher and consequently telomeres to be erroneously interpreted as longer.¹⁰ This effect is explained by the significantly lower abundance (seven to eight log change) of the housekeeping gene (single gene) in comparison to telomere sequences. This difference makes the housekeeping gene more susceptible to DNA degradation, artificially augmenting the T/S ratio. In contrast, degraded DNA produces shorter telomeres in Southern blot due to DNA fragmentation. Thus, DNA degradation results in divergent results between qPCR




Figure 2 – Correlation between telomere length measured by quantitative polymerase chain reaction (qPCR) and terminal restriction fragment (TRF) by Southern Blot. Analysis of 17 samples was performed to validate qPCR TRF. Telomere lengths measured by qPCR presented an adequate correlation with measurements of TRF by Southern blot, analyzed by linear regression ($R^2 = 0.86$; p -value < 0.0001).

Table 2 – Correlations of telomere lengths with hemolysis and inflammation markers in sickle cell disease patients.

	Mean; median (range)	Correlation with telomere length (T/S ratio) <i>r</i> (<i>p</i> -value) ^a
Hemoglobin (g/dL)	9.8; 9.6 (6.0–16.0)	0.3 (0.004)
Hematocrit (%)	29.2; 29.1 (16.7–48.4)	0.2 (0.2)
Lactate dehydrogenase (U/L)	716; 615 (257–1680)	–0.2 (0.08)
Indirect bilirubin (mg/dL)	2.2; 1.3 (0.6–9.0)	–0.2 (0.3)
Absolute reticulocyte count ($\times 10^9/L$)	274; 261 (80–667)	–0.1 (0.3)
Absolute leukocyte count ($\times 10^9/L$)	8.7; 8.6 (3.6–16.5)	–0.1 (0.3)
Absolute neutrophil count ($\times 10^9/L$)	4.5; 4.6 (1.5–9.5)	0.04 (0.7)
Absolute monocyte count ($\times 10^9/L$)	0.5; 0.4 (0.1–1.4)	–0.03 (0.7)
Absolute lymphocyte count ($\times 10^9/L$)	3.1; 2.9 (0.7–7.3)	–0.3 (0.005)
Absolute platelet count ($\times 10^9/L$)	406; 407 (81–1164)	–0.2 (0.1)
TNF- α (pg/mL)	2.6; 2.6 (0–6.9)	0.07 (0.7)
IL-8 (pg/mL)	3.5; 3.3 (0.8–11.6)	–0.4 (0.02)

IL-8: interleukin 8; TNF- α : tumor necrosis factor-alpha.

^a Spearman's rank correlation coefficient (*r*) was used to analyze bivariate associations and all *p*-values are two-tailed. *p*-Values < 0.05 are considered significant.

and Southern blotting. To avoid this problem, all samples in our study were collected and processed within 16 h from blood drawing for the purpose of telomere length measurement over a three-month period under rigorous quality control. To validate the qPCR findings of this study, telomere length was also performed by Southern blotting for 17 patients, with high correlation between both methods ($R^2 = 0.86$). Analyzing the distribution of our patients simultaneously with ones from the English study, the ages of our patients were distributed more between 20 and 60 years old whereas the patients of the English study were mainly concentrated between 20 and 40 years old. In fact, the median age of our sample is almost ten years higher than the English sample (median age of Campinas vs. England – Hb SS: 39 vs. 32 years; Hb SC/Hb S β^+ : 43 vs. 34 years). Thus, we believe that divergences in sample number, disease severity, co-morbidities, socioeconomic status, ethnicity and age distribution of the patients and possibly DNA collection might be responsible for the conflicting results and encourage further studies to clarify this issue.

The involvement of white blood cells in the severity of clinical manifestations of SCD patients is well known.^{15–18} Indeed, the number of blood cells increases in SCD for many reasons, including high levels of circulating granulocyte macrophage colony-stimulating factor and increased cell survival.^{19–22}

This study also observed that telomeres were shorter in patients using hydroxyurea. Hydroxyurea has a cytostatic effect and is capable of reducing the number of high turnover cells such as platelets, neutrophils, and reticulocytes. Hydroxyurea may have modulated blood counts, producing more prominent lymphocyte counts. In addition, most patients with more severe disease were on hydroxyurea, which may explain the association.

Telomere length did not correlate with age in SCD patients. As inflammation is chronic and starts at a young age, it may provoke excessive telomere shortening early in life, inducing early “aging” of the hematopoietic tissue in SCD patients. This finding may contribute to early multiple organ failure in SCD and may explain the shorter life expectancy related to

the disease. This hypothesis should be addressed in future studies.

Our study has some limitations. More significantly, we recruited a small number of patients with the Hb S β^+ genotype. In addition, we did not prospectively evaluate the effects of telomere length on disease complication events, such as acute chest syndrome and stroke. Finally, hydroxyurea may have interfered with telomere length by modulating leukocyte subsets.

In conclusion, our data confirm that telomere lengths are greatly influenced by inflammation and reactive oxygen species in SCD, a well-known phenomena of this disorder.

Authorship

Contribution: M.P.C. participated in the selection of patients, clinical follow up of patients, performed experimental work, analyzed the results and wrote the manuscript; B.A.S. participated in the selection of controls, performed experimental work and wrote the manuscript; N.C. and F.F.C. participated in the analysis of the results and writing of the manuscript; R.T.C. conceived the study, designed experiments, analyzed results and wrote the manuscript; S.T.O.S. conceived the study, participated in the selection of patients, clinical follow up of patients, analyzed the results and wrote the manuscript. All authors approved the final version of the manuscript.

Conflict of interest

The authors declare no conflicts of interest.

Acknowledgements

The authors would like to thank Roberto Zulli for his help with the statistical analyses.

REFERENCES

1. Colella MP, De Paula EV, Conran N, Machado-Neto JA, Annicchino-Bizzacchi JM, Costa FF, et al. Hydroxyurea is associated with reductions in hypercoagulability markers in sickle cell anemia. *J Thromb Haemost*. 2012;10(9):1967-70.
2. Almeida CB, Souza LE, Leonardo FC, Costa FT, Werneck CC, Covas DT, et al. Acute hemolytic vascular inflammatory processes are prevented by nitric oxide replacement or a single dose of hydroxyurea. *Blood*. 2015;126(6):711-20.
3. Conran N, Franco-Penteado CF, Costa FF. Newer aspects of the pathophysiology of sickle cell disease vaso-occlusion. *Hemoglobin*. 2009;33(1):1-16.
4. Dutra FF, Bozza MT. Heme on innate immunity and inflammation. *Front Pharmacol*. 2014;5:115.
5. Figueiredo RT, Fernandez PL, Mourao-Sa DS, Porto BN, Dutra FF, Alves LS, et al. Characterization of heme as activator of Toll-like receptor 4. *J Biol Chem*. 2007;282(28):20221-9.
6. Blackburn EH. Switching and signaling at the telomere. *Cell*. 2001;106(6):661-73.
7. Calado RT, Young NS. Telomere diseases. *N Engl J Med*. 2009;361(24):2353-65.
8. Masi S, Salpea KD, Li K, Parkar M, Nibali L, Donos N, et al. Oxidative stress, chronic inflammation, and telomere length in patients with periodontitis. *Free Radic Biol Med*. 2011;50(6):730-5.
9. Wolkowitz OM, Mellon SH, Epel ES, Lin J, Dhabhar FS, Su Y, et al. Leukocyte telomere length in major depression: correlations with chronicity, inflammation and oxidative stress - preliminary findings. Kiechl S, editor. *PLoS One*. 2011;6(3):e17837.
10. Gutierrez-Rodrigues F, Santana-Lemos BA, Scheucher PS, Alves-Paiva RM, Calado RT. Direct comparison of flow-FISH and qPCR as diagnostic tests for telomere length measurement in humans. *PLoS One*. 2014;9(11):e113747.
11. Cawthon RM. Telomere measurement by quantitative PCR. *Nucleic Acids Res*. 2002;30(10):e47.
12. Brouillette SW, Moore JS, McMahon AD, Thompson JR, Ford I, Shepherd J, et al. Telomere length, risk of coronary heart disease, and statin treatment in the West of Scotland Primary Prevention Study: a nested case-control study. *Lancet*. 2007;369(9556):107-14.
13. Calado RT, Cooper JN, Padilla-Nash HM, Sloand EM, Wu CO, Scheinberg P, et al. Short telomeres result in chromosomal instability in hematopoietic cells and precede malignant evolution in human aplastic anemia. *Leukemia*. 2012;26(4):700-7.
14. Drašar ER, Jiang J, Gardner K, Howard J, Vulliamy T, Vasavda N, et al. Leukocyte telomere length in patients with sickle cell disease. *Br J Haematol*. 2014;165(5):725-7.
15. Okpala I. The intriguing contribution of white blood cells to sickle cell disease - a red cell disorder. *Blood Rev*. 2004;18(1):65-73.
16. Litos M, Sarris I, Bewley S, Seed P, Okpala I, Oteng-Ntim E. White blood cell count as a predictor of the severity of sickle cell disease during pregnancy. *Eur J Obstet Gynecol Reprod Biol*. 2007;133(2):169-72.
17. Kinney TR, Sleeper LA, Wang WC, Zimmerman RA, Pegelow CH, Ohene-Frempong K, et al. Silent cerebral infarcts in sickle cell anemia: a risk factor analysis. The Cooperative Study of Sickle Cell Disease. *Pediatrics*. 1999;103(3):640-5.
18. Platt OS, Brambilla DJ, Rosse WF, Milner PF, Castro O, Steinberg MH, et al. Mortality in sickle cell disease. Life expectancy and risk factors for early death. *N Engl J Med*. 1994;330(23):1639-44.
19. Opdenakker G, Fibbe WE, Van Damme J. The molecular basis of leukocytosis. *Immunol Today*. 1998;19(4):182-9.
20. Conran N, Saad ST, Costa FF, Ikuta T. Leukocyte numbers correlate with plasma levels of granulocyte-macrophage colony-stimulating factor in sickle cell disease. *Ann Hematol*. 2007;86(4):255-61.
21. Almeida CB, Favero ME, Pereira-Cunha FG, Lorand-Metze I, Saad ST, Costa FF, et al. Alterations in cell maturity and serum survival factors may modulate neutrophil numbers in sickle cell disease. *Exp Biol Med (Maywood)*. 2011;236(11):1239-46.
22. Djaldetti M, Bergman M, Salman H, Cohen AM, Fibach E, Bessler H. On the mechanism of post-splenectomy leukocytosis in mice. *Eur J Clin Invest*. 2003;33(9):811-7.