



Revista Brasileira de Hematologia e Hemoterapia Brazilian Journal of Hematology and Hemotherapy

www.rbhh.org



Scientific Comment

Human platelet antigens and primary immune thrombocytopenia[☆]



Vagner Castro*

Universidade Estadual de Campinas (UNICAMP), Campinas, SP, Brazil

Primary immune thrombocytopenic purpura (ITP) is an acquired immune-mediated disorder characterized by transient or persistent decreased platelet count ($<100 \times 10^9/L$) that affects children and adults in the absence of other underlying causes.^{1,2} The low platelet count results from platelet destruction by antiplatelet autoantibodies associated to causes such as insufficient platelet production, which is also related to these autoantibodies, and T cell immune dysregulation (Figure 1).²

The presence of the autoantibodies may be demonstrated in the serum of approximately 70% of ITP patients, usually directed against platelet surface glycoproteins (Gp), such as Gp IIb-IIIa, Gp Ib-IX and Gp Ia-IIa.^{2,3} ITP has an estimated incidence of 5.8 cases in 100,000 young individuals with similar distribution between genders, and 1.6 cases in 100,000 middle aged individuals with higher rates among women (1.9 females: 1 male). This condition may be classified according to duration as newly diagnosed (less than 3 months), persistent (3–12 months) and chronic (more than 12 months).^{1,2} The clinical features are usually different in children and adults. In childhood, ITP usually has an abrupt onset, often starting 1–2 weeks after a viral infection or up to six weeks after vaccinations (generally the measles, mumps and rubella – MMR vaccination) with recovery being spontaneous in around 70–80% of the cases within a few weeks regardless of therapy. In adults, the disease has an insidious onset, with no

preceding illness and frequently it has a chronic course.⁴ Clinical manifestations of ITP are usually a consequence of the severe thrombocytopenia leading to bleeding symptoms. Hemorrhagic events are highly heterogeneous but patients frequently present mild mucocutaneous bleeding.^{5,6} Major bleeding risk is low in pediatric patients but in adults, additional modifiers such as existing comorbidities, age, activities and medications may affect the risk of significant bleeding.⁶ The main goal of ITP treatment is to raise the platelet count to control bleeding symptoms and to prevent major hemorrhage as well as to minimize treatment toxicity and maximize quality of life.⁶ The treatment is rarely indicated when the platelet count is above $20 \times 10^9/L$ in the absence of other factors such as major bleeding.⁶ First line therapy is generally based on the use of corticosteroids with an initial response rate of approximately 80% or intravenous immunoglobulin (IgIV or anti-D in Rh positive patients), when a rapid increase in the platelet count is needed, with a transient response in approximately 80% of cases.⁶ When the patient fails to respond or relapses after first-line therapy, the second-line treatment used generally consists of immunosuppressive agents, rituximab, splenectomy and more recently thrombopoietin receptor antagonists (TRA).^{2,6} TRA (romiplostim, eltrombopag) are new agents that have very high efficacy and few side effects but are expensive and still need to be evaluated over a longer follow-up.^{6,7}

DOI of original article: <http://dx.doi.org/10.1016/j.bjhh.2017.01.003>.

[☆] See paper by do Carmo et al. in Rev Bras Hematol Hemoter. 2017;39(2):122–6.

* Correspondence to: Platelet Immunology Laboratory, Hematology and Hemotherapy Center – Instituto Nacional de Ciência e Tecnologia do Sangue – INCTS, Universidade Estadual de Campinas (UNICAMP), Rua Carlos Chagas, 480 – Cidade Universitária “Prof. Zeferino Vaz”, Distrito de Barão Geraldo, CEP: 13083-878 Campinas, SP, Brazil.

E-mail address: vagner@unicamp.br

<http://dx.doi.org/10.1016/j.bjhh.2017.02.008>

1516-8484/© 2017 Associação Brasileira de Hematologia, Hemoterapia e Terapia Celular. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).




Figure 1 – Simplified representation of ITP mechanisms. Immune dysregulation caused by environmental factors, such as viral infections and vaccines, associated to genetic factors (cytokine polymorphisms, HLA, HPA, etc.) results in autoantibody production that mainly recognize surface glycoproteins such as GpIIb-IIIa, GpIb-IX and GpIa-IIa which are present on the surface of platelets and megakaryocytes. These autoantibodies lead to platelet destruction and decreased platelet production, resulting in thrombocytopenia and bleeding symptoms.

Although the comprehension of this complex autoimmune disorder has developed greatly since its first description in the middle of last century,⁸ mechanisms and factors leading to ITP are not yet fully understood.^{2,6} Environmental and genetic factors, such as cytokine gene polymorphisms, human leukocyte antigens (HLAs) and human platelet antigens (HPAs) seem to play a role.⁹⁻¹² HPAs are polymorphisms of platelet surface Gps which result in alloantigen expression, leading to alloimmunization mainly after transfusions and pregnancies.¹³ These polymorphisms have heterogeneous distribution in populations with distinct ethnic backgrounds and especially HPA-2 and HPA-5 have been described to be possible risk factors in the development of ITP.^{10,11,14,15} In this issue, Carmo et al.¹⁶ have evaluated the association of HPAs and chronic ITP (cITP) in a small group of patients from the Amazonian region of Brazil, which has a population with a particular HPA distribution compared to the Brazilian population in general.¹⁷ It is important to note that ITP is not a common disease and it is difficult for a single center study to enroll a large number of patients. The authors found a higher incidence of the HPA-1a, HPA-3b and HPA-5b alleles in cITP compared to healthy blood donors from the same region, but they found no involvement of HPA-2 in this group of patients. These results differ from data obtained in other studies in which the HPA-1 and HPA-3 systems seem to have no association with ITP in the populations studied, but HPA-2 and HPA-5 were associated.^{10-12,15} The mechanisms that explain these associations are not known yet. Although Carmo et al.¹⁶ evaluated a small group of patients from a single center, the results are interesting and further enhance the concept that ITP is a

complex autoimmune disorder with mechanisms that need extensive effort to understand better and provide new and more efficient treatment possibilities.

Conflicts of interest

The authors declare no conflicts of interest.

REFERENCES

1. Rodeghiero F, Stasi R, Gernsheimer T, Michel M, Provan D, Arnold DM, et al. Standardization of terminology, definitions and outcome criteria in immune thrombocytopenic purpura of adults and children: report from an international working group. *Blood*. 2009;113(11):2386-93.
2. Psaila B, Bussell JB. Immune thrombocytopenia (ITP). In: Michelson AD, editor. *Platelets*. 3rd ed. San Diego, CA, USA: Academic Press, Elsevier Science; 2013. p. 819-33.
3. Cines DB, McMillan R. Pathogenesis of chronic immune thrombocytopenic purpura. *Curr Opin Hematol*. 2007;14(5):511-4.
4. Stasi R. Pathophysiology and therapeutic options in primary immune thrombocytopenia. *Blood Transfus*. 2011;9(3):262-73.
5. Neunert C, Lim W, Crowther M, Cohen A, Solberg L Jr, Crowther MA, et al. The American Society of Hematology 2011 evidence-based practice guideline for immune thrombocytopenia. *Blood*. 2011;117(16):4190-207.
6. Neunert CE. Current management of immune thrombocytopenia. *Hematol Am Soc Hematol Educ Program*. 2013;2013:276-82.

7. Grace RF, Neunert C. Second-line therapies in immune thrombocytopenia. *Hematol Am Soc Hematol Educ Program*. 2016;2016(1):698-706.
8. Harrington WJ, Minnich V, Hollingsworth JW, Moore CV. Demonstration of a thrombocytopenic factor in the blood of patients with thrombocytopenic purpura. *J Lab Clin Med*. 1951;38(1):1-10.
9. Consolini R, Legitimo A, Caparello MC. The centenary of immune thrombocytopenia – Part 1: Revising nomenclature and pathogenesis. *Front Pediatr*. 2016;4:102.
10. Kim BS, Song KS. Genetic polymorphism of human platelet specific antigen(HPA) in patients with immune thrombocytopenia. *Thromb Haemost*. 1997:1037.
11. Castro V, Oliveira GB, Origa AF, Annichino-Bizzacchi JM, Arruda VR. The human platelet alloantigen 5 polymorphism as a risk for the development of acute idiopathic thrombocytopenia purpura. *Thromb Haemost*. 2000;84(2):360-1.
12. Pavkovic M, Stojanovic A, Karanfilski O, Cevreska L, Spiroski M. Association of polymorphisms in human platelet antigens with idiopathic thrombocytopenic purpura in Macedonians. *Prilozi*. 2012;33(1):135-46.
13. Kaplan C, Ni H, Freedman J. Alloimmune thrombocytopenia. In: Michelson AD, editor. *Platelets*. 3rd ed. San Diego, CA, USA: Academic Press, Elsevier Science; 2013. p. 953-70.
14. Castro V, Origa AF, Annichino-Bizzacchi JM, Soares M, Menezes RC, Goncalves MS, et al. Frequencies of platelet-specific alloantigen systems 1-5 in three distinct ethnic groups in Brazil. *Eur J Immunogenet*. 1999;26(5):355-60.
15. Thude H, Gatzka E, Anders O, Barz D. Allele frequencies of human platelet antigen 1, 2, 3, and 5 systems in patients with chronic refractory autoimmune thrombocytopenia and in normal persons. *Vox Sang*. 1999;77(3):149-53.
16. do Carmo JC, Klippel PS, Cordeiro SC, Fernandes AM, Pinto RM, Weber SS, et al. Molecular typing of human platelet antigens in immune thrombocytopenia patients on North Brazil. *Rev Bras Hematol Hemoter*. 2017;39(2):122-6.
17. Portela CN, Schriefer A, Albuquerque SR, Perdomo RT, Parente AF, Weber SS. The human platelet alloantigen profile in blood donors from Amazonas, Brazil. *Transfus Med*. 2016;26(6):448-56.