

Case Report

Compound heterozygote of Hb D^{Iran} [HBB: c.67G>C, β 22(B4) Glu>Gln] with β⁰-thalassemia [cfs 41/42 (-CTTT)] from Eastern India



Pradeep Kumar Mohanty^a, Satyabrata Meher^a, Snehadhini Dehury^a,
Subhra Bhattacharya^b, Kishalaya Das^a, Siris Patel^a, Biswanath Sarkar^{b,*}

^a Veer Surendra Sai Institute of Medical Science and Research (VIMSAR), Burla, Sambalpur, Odisha, India

^b Anthropological Survey of India, Kolkata, India

ARTICLE INFO

Article history:

Received 1 September 2017

Accepted 21 September 2017

Available online 1 December 2017

Introduction

Hereditary hemoglobinopathies, the most common monogenic hemoglobin (Hb) disorders, result in a variety of clinical consequences. It has been observed that various Hb variants and thalassemias are found common to specific ethnic groups and regions. Hb D^{Iran} is a structural Hb variant resulting from the substitution of glutamine with glutamate at codon 22 (GAA>CAA, Glu>Gln) of the beta globin gene. This Hb variant was first reported by Rahbar in 1973 in a family from the central part of Iran.¹ A deletion of four bases in codon 41/42 (-CTTT) is a rare β⁰-thalassemia mutation reported in India with a prevalence of 3–15%.² The present report describes a rare combination of these two mutations for the first time in India.

Case report

A 45-year-old Sikh female from Sundergarh district of Odisha, India with a family history of β-thalassemia attended the Sickle Cell Institute, VIMSAR, Burla to screen her status. She was asymptomatic and had no history of blood transfusion or vaso-occlusive crisis. Ultrasonographic examination revealed normal spleen and liver. The various investigations of the proband and her daughter, including a complete blood count and biochemistry, are shown in [Table 1](#). As evident, the index case had features suggestive of microcytic hypochromic anemia (mean corpuscular volume: 58.7 fL and mean corpuscular hemoglobin: 17.8 pg). An iron profile study indicated possible iron overload [iron 5.027 mg/dL (reference range – RR: 0.005–0.175 mg/dL); ferritin: 138.7 μg/L (RR: 20–200 μg/L) and transferrin: 490.05 mg/dL (RR: 212–360 mg/dL)].

* Corresponding author at: Anthropological Survey of India, 4th Floor, Spirit Building, 27 Jawaharlal Nehru Road, Kolkata 700016, India.

E-mails: drbnsarkar@rediffmail.com, drbnsarkar@yahoo.com (B. Sarkar).

<https://doi.org/10.1016/j.bjhh.2017.09.001>

2531-1379/© 2017 Associação Brasileira de Hematologia, Hemoterapia e Terapia Celular. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Table 1 – Hematological and biochemical indices of proband and her daughter.

	Unit (SI)	Proband	Daughter
White blood cell count	$\times 10^9/L$	7.4	6.9
Red blood cell count	$\times 10^{12}/L$	5.67	5.07
Hemoglobin	g/L	10.1	9.9
Hematocrit	%	33.3	33.8
Mean corpuscular volume	fL	58.7	66.7
Mean corpuscular hemoglobin	Pg	17.8	19.5
Mean corpuscular hemoglobin concentration	g/dL	30.3	29.3
Platelet count	$\times 10^9/L$	169	171
Serum creatinine	$\mu\text{mol}/L$	0.05	0.08
Aspartate transaminase	U/L	12.7	15.3
Alanine transaminase	U/L	12.5	9.0
Total bilirubin	$\mu\text{mol}/L$	0.03	0.34
Lactate dehydrogenase	U/L	198	189
Iron	$\mu\text{mol}/L$	5.027	5.258
Transferrin	g/L	490.05	462.09
Ferritin	pmol/L	138.7	111.6

Because of the endemicity of the sickle cell hemoglobinopathy and its combination with β -thalassemia in this region, the sickling test and alkaline agarose gel Hb electrophoresis were performed; the sickling test was negative and a single band in the Hb S/D position was observed by Hb electrophoresis (pH-8.6). Cation exchange high performance liquid chromatography (CE-HPLC) was performed using the VARIANT-II hemoglobin testing system with the CDM 5.1.1TM software and Beta Thal Short Program (Bio-Rad Laboratories, Hercules, CA, USA) which showed a prominent peak in the Hb A₂ window [82.8%; retention time (RT): 3.57 min] with low Hb A₀ and Hb F peaks (4.6% and 1.0%, respectively) (Figure 1). The possibility of homozygous Hb E was ruled out by the absence of a band in the position of Hb A₂/Hb E by Hb electrophoresis. Hence, the case was initially suspected to be a rare finding of Hb Tianshui, which has similar alkaline Hb electrophoresis and CE-HPLC findings as reported earlier.³ However, the peak morphology characteristic of the present case was different from that of Hb Tianshui. Consequently, β -globin gene sequencing using the Big-Dye terminator protocol Ver 3.1 in an automated ABI-3730 DNA Analyzer (Applied Biosystems, USA) confirmed the case to be compound heterozygote of Hb D^{Iran} ($\beta 22$ (B4) Glu>Gln; HBB: c.67G>C, GAA>CAA, rs33959855) (Figure 2A) with a β^0 -thalassemia mutation [4-base pair (bp) deletion at cds 41/42 (-CTTT)] (Figure 2B). An additional investigation for the XmnI polymorphism and deletional alpha thalassemia revealed that the proband was a homozygote for the wild type allele (CC) in the XmnI locus and heterozygous for the 3.7 kb deletional alpha thalassemia. The presence of the homozygous wild allele in the XmnI locus corroborates the observed low Hb F level in this case, while the possible effect of heterozygous deletional alpha thalassemia on the microcytic hypochromic red cell parameters of the case could

be masked by the simultaneous presence of the cds 41/42 (-CTTT) mutation.

Discussion

The Hb D^{Iran} trait and homozygous cases have been reported earlier.^{4,5} However, few studies have reported compound heterozygotes of Hb D^{Iran} with other Hb variants like Hb S and Hb D^{Punjab}, β^+ -thalassemia IVS1–5 (G>C), β^0 -thalassemia (619 bp-deletion) and undefined β -thalassemia from India and Pakistan. Various studies have reported that the quantity of Hb D^{Iran} eluting in the Hb A₂ window in HPLC varies from 36.0 to 47.7% in a heterozygous condition, while in compound heterozygous states, the quantity varies between 47.3 and 94.4% (with Hb D^{Punjab}, Hb S, β -thalassemia with the 619 bp deletion mutation and beta thalassemia with unknown mutation).^{6–10} Almost all these cases were mild in presentation with concomitant anemia.

Codon 22 (GAA), is a mutational hotspot in exon I of the human β globin gene, although it does not take part in α - β or protein-heme interactions, as this is an external residue positioned at the B4 site of the helix. To date, six Hb variants (Hb D^{Iran}, Hb E-Saskatoon, Hb G-Coushatta, Hb D-Granada, Hb G-Taipei and Hb Bury) and one β^0 -thalassemia mutation [Codon 22 (G>T); GAA(Glu)>TAA (stop codon)] have been reported involving this codon. In Hb D^{Iran}, the change of glutamate to glutamine leads to an overall change of charge from negative to positive resulting in a protein that migrates to the position of Hb S in alkaline Hb electrophoresis.^{1,10} This rare variant has heat stability with no effect on oxygen equilibrium, intracellular 2,3-diphosphoglycerate or the Bohr effect.¹⁰ The homozygous state of Hb D^{Iran} reveals a milder phenotype even when Hb D^{Iran} co-inherits with β^0 -thalassemia.^{5,9} The present case agrees with this as evidence from the clinical and hematological investigations show. Although Hb D^{Iran} in combination with β -thalassemia produces a moderate microcytic and hypochromic red cell picture that is not transfusion dependent, the appearance of Hb D^{Iran} in the position of Hb S in alkaline agarose gel electrophoresis can lead to significant confusion and might falsely be reported as a sickle cell hemoglobinopathy unless a sickling test and HPLC are read together with these findings. Hb S can easily be distinguished from Hb D^{Iran} by performing CE-HPLC.

Reportedly in CE-HPLC, nine abnormal Hbs elute in the Hb A₂ window (3.27–3.83 as per the manufacturer's guidelines in the operating software): Hb Deer Lodge, Hb Lepore, Hb D^{Iran}, Hb E, Hb Hamadan, Hb Osu-Christiansborg, Hb Tianshui, Hb G Honolulu and Hb G Copenhagen. Among these, Hb Deer Lodge, Hb Lepore and Hb D^{Iran} elute prior to the standard RT of Hb A₂ (3.6 min) while others have higher RT to that of Hb A₂. Interestingly, Hb Lepore has the lowest average quantity (7–15%) followed by Hb G Honolulu (about 15% of total hemoglobin quantity) and Hb E (about 30% of total hemoglobin in absence of α -thalassemias). All the other variants eluting in the Hb A₂ window have variant hemoglobin quantities higher than 30% on average under heterozygous conditions, making it difficult to distinguish in HPLC. Amongst these, Hb D^{Iran} has been reported to elute in this window at

Peak name	Calibrated area %	Area %	Retention time (min)	Peak area
Unknown	---	0.3	0.96	6736
F	1.0	---	1.09	17882
Unknown	---	0.8	1.60	16390
P3	---	4.0	1.73	78869
A0	---	4.6	2.20	90859
Unknown	---	1.0	2.50	20501
A2	82.8*	---	3.57	1740824

Total area: 1,972,061

F concentration = 1.0 %

A2 concentration = 82.8* %

*Values outside of expected ranges

Analysis comments:




Figure 1 – CE-HPLC showing characteristic peak of HbD^{Iran}/β⁰ thal [c_{ds} 41/42 (-CTTT)].

a RT of 3.49–3.58 min; almost in the middle of the window (3.27–3.83). However, the pattern of mobility of these variants in alkaline electrophoresis is quite interesting. Hb E stands apart as its mobility is at the position of Hb A₂ and can be easily identified. Hb Deer Lodge and Hb G Copenhagen have almost similar alkaline electrophoretic mobility i.e., slightly anodic to the position of Hb S. The rest of the variants are very difficult to distinguish even in alkaline electrophoresis because of their identical mobility to the position of Hb S. Additionally, all of these variants have a negative sickling test result.¹¹

Further, as Hb D^{Iran} elutes in the Hb A₂ window in HPLC masking elevated Hb A₂, it becomes difficult to suspect the presence of β-thalassemia and direct gene sequencing needs to be performed. To the best of our knowledge, this is the first report of Hb D^{Iran} with β⁰-thalassemia [c_{ds} 41/42 (-CTTT)] reported from Odisha, India.

Funding

The study was performed under the Odisha Sickle Cell Project, funded by the National Health Mission of India, Odisha, India.

Conflicts of interest

The authors declare no conflicts of interest.

Acknowledgement

We acknowledge the support from Prof. Bijaya Kumar Dutta, Dean & Principal, Veer Surendra Sai Institute of Medical Science and Research (VIMSAR), Burla, Odisha. The authors are indebted to late Dr. Dilip Kumar Patel, Ex-Associate Professor,




Figure 2 – (A) DNA sequence chromatogram showing HbD^{Iran} mutation on 22 codon (GAA>CAG). (B) DNA sequence chromatogram showing β⁰ thal {4 bp del Cds 41/42 (-CTTT)}

Department of Medicine, Veer Surendra Sai Medical College (now VIMSAR), Burla, Sambalpur, Odisha and Ex-Project Coordinator, Odisha Sickle Cell Project (NHM, Odisha) for his inestimable contribution to this study. The authors acknowledge the support of the Director, Anthropological Survey of India, Ministry of Culture, Government of India for sanctioning the collaboration program (Vide letter No. 18-22/PMI/2011, dated June 15 2013).

REFERENCES

- Rahbar S. Hemoglobin D Iran: β₂22 glutamic acid→ glutamine (B4). *Br J Haematol*. 1973;24(1):31–5.
- Verma IC, Saxena R, Thomas E, Jain PK. Regional distribution of beta-thalassemia mutations in India. *Hum Genet*. 1997;100(1):109–13.
- Meher S, Dehury S, Mohanty PK, Patel S, Pattanayak C, Bhattacharya S, et al. Hb Tianshui (HBB: C.119A > G) in compound heterozygosity with Hb S (HBB: C.20A>T) from Odisha, India. *Hemoglobin*. 2016;40(4):270–2.
- De Marco EV, Crescibene L, Bagalh A, Brancati C, Quatieri A, Bria M. Hb D-Iran [B22(B4)Glu->Gln] In Southern Italy. *Hemoglobin*. 1994;18(1):65–9.
- Thornburg CD, Zimmerman SA, Schultz WH, Ware RE. An infant with homozygous hemoglobin D-Iran. *J Pediatr Hematol Oncol*. 2001;23(1):67–8.
- Serjeant B, Myerscough E, Serjeant GR, Higgs DR, Moo-Penn WF. Sickle cell-hemoglobin D Iran: benign sickle cell syndrome. *Hemoglobin*. 1982;6(1):57–9.
- Gupta A, Saraf A, Dass J, Mehta M, Radhakrishnan N, Saxena R, et al. Compound heterozygous hemoglobin D-Punjab/hemoglobin D-Iran: a novel hemoglobinopathy. *Indian J Hematol Blood Transfus*. 2014;30 Suppl. 1:S409–12.
- Bhat VS, Mandal AK, Mathew B. Co-inheritance of HbD(Iran)/beta thalassemia IVS1-5 (G > C) trait in a Punjabi lady with diabetes. *Ind J Clin Biochem*. 2012;27(2):202–6.
- Agrawal MG, Bhanushali AA, Dedhia P, Jeswani KD, Dayanand S, Dasgupta A, et al. Compound heterozygosity of Hb D^{Iran} (β²² Glu → Gln) and β⁰-thalassemia (619 bp-deletion) in India. *Eur J Haematol*. 2007;79(3):248–50.
- Rohe RA, Sharma V, Ranney HM. Hemoglobin D Iran α₂^Aβ₂²²-Glu → Gln in association with thalassemia. *Blood*. 1973;42(3):455–62.
- Hardison RC, Chui DH, Giardine B, Riemer C, Patrinos GP, Agnagou N, et al. HbVar: a relational database of human hemoglobin variants and thalassemia mutations at the globin gene server. *Hum Mutat*. 2002;19(3):225–33.