



Research Letters

Protecting biodiversity in urbanizing regions: The role of urban reserves for the conservation of Brazilian Atlantic Forest birds

Thayz R. Enedino^a, Alan Loures-Ribeiro^b, Bráulio A. Santos^{b,*}^a Programa de Pós-Graduação em Biodiversidade, Universidade Federal da Paraíba, Rodovia PB 079, CEP 58397-000, Campus II, Areia PB, Brazil^b Departamento de Sistemática e Ecologia, Universidade Federal da Paraíba, Cidade Universitária, Castelo Branco, CEP 58051-900, Campus I, João Pessoa PB, Brazil

ARTICLE INFO

Article history:

Received 1 August 2017

Accepted 1 November 2017

Available online 15 November 2017

Keywords:

Birdwatching
Forest fragmentation
Habitat loss
Protected areas
Species richness
Urbanization

ABSTRACT

Although the Brazilian Atlantic Forest and other global biodiversity hotspots are rapidly urbanizing, there is little information on the potential of metropolitan regions to safeguard biodiversity within their reserves. We sampled bird assemblages in nine protected areas (14.8–1058 ha) of the metropolitan region of João Pessoa, Northeast Brazil, to test the hypothesis that larger urban reserves are more effective than smaller ones in protecting the regional species richness, as more habitat and environmental heterogeneity are available for birds. We recorded 126 species belonging to 41 families, including seven endemics and two introduced species. Thraupidae and Tyrannidae were the most representative families with 16 species each. Larger areas clearly protected greater proportion of the regional species richness, but the smaller areas accounted for at least 35.7% of the regional richness. Differences in size and isolation significantly influenced species similarity, but hunting, catching for illegal trade, presence of exotic predators and noise pollution were likely to affect composition as well. The remarkable bird diversity suggests good potential for legal-based, well-planned birdwatching activities in the studied region. The results highlight that urban reserves of any size are valuable for bird conservation in metropolitan regions, though larger ones should be prioritized.

© 2017 Associação Brasileira de Ciência Ecológica e Conservação. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Although the Brazilian Atlantic Forest is considered one of the world's largest repositories of biodiversity (Myers et al., 2000), only 11.4–16% of the original forest cover were left (Ribeiro et al., 2009). More than 80% of the almost 250,000 forest fragments are smaller than 50 ha, about half of the remaining vegetation is less than 100 m from the nearest edge, and the average distance between the fragments is 1440 m (Ribeiro et al., 2009). During this long process of habitat loss and fragmentation, many metropolitan regions have raised and dominated agricultural landscapes that first replaced the forest (Joly et al., 2014; Stevens, 2014). As in other regions of the world, cities and dense settlements show a clear trend for growth in the Brazilian Atlantic Forest (Seto et al., 2013; United Nations, 2014), which has been somehow counterbalanced by the creation of urban protected areas. However, to date there is little empirical evidence showing the ability of such protected areas to

safeguard biodiversity and ecosystem services that positively affect biodiversity.

With nearly 2000 species of birds, including 277 endemics and 1692 resident species, Brazil has one of the most diverse avifauna in the world (Piacentini et al., 2015). Studies have shown that forest destruction and fragmentation have resulted in species loss at local and regional scales (Anjos et al., 2015). The microclimatic changes, especially the increase in light and decrease in air humidity arising from landscape modification, eliminate most sensitive species (Morante-Filho et al., 2017). The reduction of food and sites for nesting also lead to the decline of many species, especially those that require large areas (Marini and Garcia, 2005). This decline, in turn, results in negative effects on the plants that birds pollinate and disperse, insects and other animals they consume as well as predators and parasitoid they are attacked by (Ford et al., 2001). Birds are also the basis for birdwatching tourism, considered one of the most biodiversity-friendly activities with potential to conciliate economic development with human well-being (Sekercioglu, 2002; Millennium Ecosystem Assessment, 2005). Nonetheless, the provision of this ecosystem service in urban reserves of the Brazilian Atlantic Forest has been poorly explored.

* Corresponding author.

E-mail address: braulio@dse.ufpb.br (B.A. Santos).

In the Northeast of Brazil it is located the Pernambuco Center of Endemism of the Brazilian Atlantic Forest, a ~56,000-km² area that includes forests north of the São Francisco River in the states of Alagoas, Pernambuco, Paraíba and Rio Grande do Norte (Porto et al., 2005). This center homes at least 434 bird species, including 35 endemics and 40 officially recognized as threatened by the Brazilian Ministry of Environment (MMA) (Roda et al., 2011; Lobo-Araújo et al., 2013), as well as four medium-sized cities that encompass metropolitan regions of 1–5 million inhabitants (United Nations, 2014). One of them is the metropolitan region of João Pessoa, state of Paraíba, where this study was conducted. This region has faced intense urbanization since 1970 and a peak of protected area creation in the 2000s, following the promulgation of the federal law 9985/2000 that established the National System of Protected Areas. Currently, 18 protected areas varying from 14.8 ha to 1058 ha safeguard a variety of ecosystems under distinct administrative levels (i.e., municipal, state, federal areas) (Stevens, 2014). These areas are expected to home a large portion of the 395 species, 28 orders and 67 families of birds reported for the whole state (Marinho, 2014), but no estimate has been done so far, precluding the adoption of legal-based, environmental-friendly management of the birds.

In this study we record the avifauna of nine protected areas in the metropolitan region of João Pessoa to test the hypothesis that larger urban reserves are more effective than smaller ones in protecting the regional richness, as more habitat and environmental heterogeneity are available for birds (Sekercioglu et al., 2002). We quantify the relative contribution of each protected area for regional bird species richness and assess how differences in protected area size and isolation affect species composition. We also provide a brief recommendation for the development of legal-based, well-planned birdwatching activities in the region.

Materials and methods

Study area

Our study area has 63,000 ha and encompasses the most densely populated portion of the metropolitan region of João Pessoa, including the municipalities of Cabedelo, Santa Rita, Bayeux, Lucena and João Pessoa (34° 58' 87" W 7° 15' 46" S and 34° 47' 19" W 6° 54' 43" S) (Fig. 1). A recent study indicates that 57% of the original vegetation cover were lost between 1974 and 2010, and the most affected areas were the rain forests, sandbank forests (*restingas*) and *tabuleiro* forests (Stevens, 2014). Currently, the 14,847 ha of native vegetation left are scattered over 236 forest remnants, most smaller than 50 ha and surrounded by urban agglomerations or agricultural fields (mainly sugarcane plantations) under peri-urban conditions. About 20% of the remaining vegetation is protected by law within 18 protected areas, either within the federal law 9985/2000 or other legal instruments at the state and municipal spheres. Nine protected areas showed minimum level of security to permit access and fieldwork inside. They vary in size, internal level of fragmentation, surrounding matrix, predominant vegetation, presence of waterbodies, and administration regime (Table 1). Altogether, they conserve a mosaic of ecosystems composed of distinct types of forests, from rain to drier forests (*restinga* and *tabuleiro* forests), salt marshes, mangrove swamps and estuaries. In this study we consider urbanization as a multidimensional process that manifests itself through rapidly changing human population and changing land cover (*sensu* Seto et al., 2013) and, consequently, assume that all remaining vegetation inside the study area are facing urbanization.

Field surveys

Between June 2014 and May 2015, we used 50 Mackinnon lists of 10 species to sample birds in each of the nine protected areas, starting early in the morning (around 6 am) to maximize species detection (Bibby et al., 1998). In the Mackinnon method (Ribon, 2010), each species is recorded only once in each list until the maximum number of species in the list is reached (10 in our study). After that, a new list is started. We produced 10 lists per day per area during five consecutive days, totaling 450 lists. An advantage of this method is that the list may be generated throughout a day without application time, or on subsequent days. Furthermore, it allows the calculation of species frequency by list and area, uncovering common and rare species (Manhães and Loures-Ribeiro, 2011). Nonetheless, we were conservative and excluded any doubtful record from our lists.

We recorded the birds visually with binoculars Nikon 7 × 50, aurally through species vocalization and confirmed species identification with specialized literature and sound recordings of private collection (Sick, 1997; Erize et al., 2006; Sigrist, 2009). We alternated consecutive field surveys between smaller and larger protected areas to control for possible seasonal effects on bird detection, activity and occurrence along the gradient of protected area size. Furthermore, we restricted the surveys to the forest formations of each protected area (i.e., rain forest, *tabuleiro* forest, *restinga* forest; Table 1) to minimize compositional differences attributed to habitat type. The taxonomic classification of the species followed the Brazilian Ornithological Records Committee (Piacentini et al., 2015). We searched for the conservation status of identified species in the red lists of IUCN and the Brazilian government. Introduced species were also identified.

Data analyses

To quantify the relative contribution of each protected area for the regional bird species richness, we divided the species richness of each area by the regional species richness. Our expectation was that smaller areas showed smaller proportion of the regional richness than larger areas. A simple linear regression was performed in JPM 8 (SAS Institute Inc., Cary, NC, USA) to test this relationship. We used the Jackknife method of first order to estimate the regional species richness following the equation: $S_{\text{jack}1} = S_{\text{obs}} + Q1 (m - 1/m)$, where $S_{\text{jack}1}$ is the estimated species richness; S_{obs} is the observed number of species in all lists; $Q1$ is the number of species recorded in a single list; and m is the number of lists. We obtained the estimates and species accumulation curves with the software Estimates 9.1 with 1000 randomizations (Colwell, 1997).

To assess how differences in protected area size and isolation affected species composition, we constructed a dissimilarity matrix of taxonomic composition based on the Jaccard distance ($1 - J$) and used Mantel tests to correlated it with matrices of geographic distance and difference in size. We expected that less isolated and similar-sized protected areas showed an avifauna more similar to each other than more isolated, different-sized areas. We produced matrices and ran Mantel tests in Primer 6 (Clarke and Gorley, 2006). To identify the most common and rarest species in the region, we calculated the frequency of occurrence of each species by dividing their number of occurrence by the total number of occurrence of all species in the lists.




Fig. 1. Study area in the metropolitan region of João Pessoa, Northeast Brazil. The nine protected areas are shown in black.

Results

Contribution of protected areas to the regional species richness

We recorded a total of 126 species (7 endemics), 17 orders and 41 bird families (Table S1). The richness estimated with Jackknife method of first order was approximately 136 species (Fig. S1), indicating the presence of 10 species more in our study region. Thraupidae and Tyrannidae, with 16 species each, were the most representative families, followed by Trochilidae (8 spp.), Thamnophilidae (6 spp.), Rhynchocyclidae (5 spp.), Picidae (5 spp.) and Columbidae (5 spp.). Eighteen of the 41 families were represented by only one species (Fig. S2).

As expected, the richest area was the largest reserve (Gargaú Private Reserve of Natural Heritage, 1058 ha), with 95 species, accounting for 75.4% of the regional species richness. The Buraquinho Refuge of Wildlife (518 ha) and the Xem-Xem State Park Forest (182 ha), second and third largest areas, followed with 80 and 62 species, respectively (63.5% and 49.2% of the regional richness). Five areas ranging from 22 ha to 155 ha protected between 40.5% and 43.7% of the regional richness: Cuiá Natural Municipal Park (56 species), Arruda Câmara Zoo Botanical

Park (55 species), Cabedelo Sandbank National Forest (54 species), Cabo Branco Park (51 species) and Paraíba Federal University Forest (51 species). The smallest area, the Cidade Verde Municipal Forest Nursery (14.8 ha), protected as much as 35.7% of the regional richness (45 species). When we modeled the proportion of regional richness in each protected area as a function of its size, a clear positive relationship arose ($R^2=0.916$; $F_{1,7}=77.2$; $P<0.0001$; Fig. 2), showing the remarkable contribution of larger areas for bird conservation.

Species similarity across the protected areas


Pairwise similarity averaged 52.6% across the protected areas, varying from 36.5% (Gargaú Private Reserve of Natural Heritage vs. Cidade Verde Municipal Forest Nursery) to 71.4% (Arruda Câmara Zoo Botanical Park vs. Paraíba Federal University Forest). There was a trend for distance-decay in species composition ($Rho=0.379$; $P=0.055$) supported by lower similarity between more distant areas (Fig. 3). Differences in size also influenced species composition ($Rho=0.421$; $P=0.04$), with similar-sized areas being more taxonomically similar than different-sized areas (Fig. 3). Nonetheless, in both cases the strength of the relationship (Rho values) was

Table 1

The nine protected areas studied in the metropolitan region of João Pessoa, Northeast Brazil, sorted by their size (from the largest to the smallest area).

Name of protected area	Declared area (ha)	Belongs to SNUC	Creation act	Administration	Surrounding matrix	Predominant vegetation**	Presence of waterbodies
Gargaú Private Reserve of Natural Heritage (<i>RPPN do Engenho Gargaú</i>)	1058	Yes	IBAMA Ordinance 64-N, 14/June/1994	Federal	Agricultural	Rain forest (290 ind./39 spp.)	Interior and neighborhood
Buraquinho Refuge of Wildlife (<i>Refúgio de Vida Silvestre da Mata do Buraquinho</i>)	517.8	Yes	State Decree no. 35.195, 28/August/2014	State	Urban	Rain forest (366 ind./52 spp.)	Interior
Xem-Xem State Park Forest (<i>Parque Estadual Mata do Xém-Xém</i>)	182	Yes	State Decree no. 21.262, 28/August/2000	State	Urban and agricultural	Rain forest and <i>tabuleiro</i> forest (309 ind./57 spp.)	Interior and neighborhood
Paraíba Federal University Forest (<i>Mata da UFPB</i>) [*]	155	No	State Law 1.366, 2/December/1955	Federal	Urban	Rain forest (277 ind./41 spp.)	Neighborhood
Cabedelo Sandbank National Forest (<i>Floresta Nacional da Restinga de Cabedelo</i>)	116.8	Yes	Federal Decree no. 2/June/2004	Federal	Urban and mangrove	<i>Restinga</i> forest and mangrove (218 ind./44 spp.)	Interior and neighborhood
Cabo Branco Park [*] (<i>Parque Cabo Branco</i>)	73.1	No	Municipal Decree no. 5.343, 29/April/2005	Municipal	Urban	<i>Tabuleiro</i> forest (144 ind./25 spp.)	None
Cuiá Natural Municipal Park (<i>Parque Natural Municipal do Cuiá</i>)	43.2	Yes	Municipal Decree no. 7.517, 29/April/2011	Municipal	Urban	Rain forest (440 ind./42 spp.)	Interior and neighborhood
Arruda Câmara Zoo Botanical Park [*] (<i>Parque Zoobotânico Arruda Câmara</i>)	22.4	No	?	Municipal	Urban	Rain forest (149 ind./34 spp.)	Interior
Cidade Verde Municipal Forest Nursery (<i>Horto Florestal Municipal Cidade Verde</i>)	14.8	No	?	Municipal	Urban	Rain forest (656 ind./43 spp.)	Neighborhood

* Internally fragmented and urbanized.



** We show in parenthesis the number of plant individuals and species with diameter at breast height ≥ 2.5 cm in 0.1 ha of forest formation, following Gentry's protocol of vegetation sampling.**Fig. 2.** Relationship between the size of protected areas (ha) and the proportion of regional bird species richness (126 spp.) in the metropolitan region of João Pessoa, Brazil.

low, suggesting that the size and isolation of the protected areas are not the main drivers of species composition.

Common species and conservation status

The most frequent species was *Pitangus sulphuratus* (Great kiskadee; Tyrannidae), taking place in all areas and in more than half of the 450 lists. Other common species included *Coereba flaveola* (Thraupidae), *Vireo olivaceus* (Vireonidae), *Fluvicola nengeta* (Tyrannidae), and *Columbine talpacoti* (Columbidae), occurring in 30%–37% of the lists (Table S1). In addition to *P. sulphuratus*, other 18 species occurred in all protected areas, but at lower frequencies. Ten species were recorded only once in the entire study.

Only one species, *Picumnus fulvescens* (Picidae), is recognized by the IUCN as near threatened (NT). However, six species appear in the Brazilian Red List of Endangered Species in worrying categories: *Penelope superciliaris* (Cracidae; CR – Critically endangered), *Momotus momota* (Momotidae; EN – Endangered),

**Fig. 3.** Relationship of species dissimilarity (1 – J) with geographic distance and difference in size for nine protected areas in the metropolitan region of João Pessoa, Brazil.

Conopophaga melanops (Conopophagidae; VU – Vulnerable), *Xenopus minutus* (Xenopidae; VU), *Platyrinchus mystaceus* (Platyrinchidae; VU) and *Xiphorhynchus atlanticus* (Dendrocolaptidae; VU). Only two introduced species were detected: *Estrilda astrild* (Estrildidae) and *Passer domesticus* (Passeridae), but *P. domesticus* occurred in 25.8% of the lists.

Discussion

Our findings support the hypothesis that larger urban reserves are more effective than smaller ones in protecting the regional richness, emphasizing the critical role of large reserves (>1000 ha) for bird conservation in the studied metropolitan region. However, even the smallest and more isolated area significantly contribute to the regional avifauna, safeguarding more than one third of the species richness found in the region (see also [Silva et al., 2014](#)). This clearly demonstrates that small urban reserves are not negligible for bird conservation. The results also highlight the great conservation value of the metropolitan region of João Pessoa for bird conservation at broader spatial scales, protecting at least 32% of the 395 species reported for the state of Paraíba ([Marinho, 2014](#)) and 29% of the 434 species reported for the Pernambuco Center of Endemism ([Roda et al., 2011](#); [Lobo-Araújo et al., 2013](#)). This regional richness is actually underestimated according to the species accumulation curve, possibly because migratory, wandering and regionally rare species require greater sampling period to be detected.

The greater richness observed in larger areas is not surprising. Compared with smaller areas, larger areas usually offer a wider variety of food resources, micro-habitats and suitable conditions for reproduction, dispersion and survival ([Sekercioglu et al., 2002](#)). Species with different physiological and ecological requirements are sorted across the micro-habitats available, increasing the turnover within the area and resulting in the co-occurrence of a large number of species. In our study, all but one reserve had waterbodies inside and/or nearby, two were composed of more than one vegetation type and three were internally fragmented and urbanized. These features aggregate environmental heterogeneity to all, but certainly more to the largest areas (i.e. the Gargaú Private Reserve of Natural Heritage, 1058 ha, and the Buraquinho Refuge of Wildlife, 518 ha), allowing them to contribute more than the others to the regional richness.

Although 23 bird species have left the Brazilian list of endangered species (MMA Ordinance 444/2014), 234 species/subspecies are still threatened. In our surveys we recorded six endangered species, demonstrating that the urbanizing region is still able to harbor threatened birds. However, if future urbanization in the region follows the trend of the 1970–2010 period ([Stevens, 2014](#)), a massive erosion in bird diversity is expected ([Sol et al., 2017](#)). During the fieldwork, it was also evident the unauthorized use of the protected areas for illegal hunting, firewood extraction, livestock, fishing, garbage deposition, and other destructive uses, especially in those under state and municipal administration (see also [Silva-Junior and Santos, 2017](#)). These chronic management failures affect the biota as whole and may also jeopardize the long-term bird conservation.

The presence of the introduced species *Estrilda astrild* and *Passer domesticus* is also worrisome, as biological invasion is a powerful driver of biodiversity loss at multiple spatial scales (e.g. [Jones et al., 2008](#)). *Estrilda astrild* entered accidentally in the Brazilian territory, but *P. domesticus* was introduced intentionally to control agricultural pests ([Sick, 1997](#)). The effects of the introduction of exotic species are usually irreparable, especially if the exotics is a superior competitor able to rapidly reproduce, establish and spread over the new areas ([Shaw et al., 2008](#); [MacGregor-Fors et al., 2010](#)). In this

scenario, native species are not able to fight back and the exotics becomes an invasive species. In West Mexico, for example, areas invaded by *P. domesticus* have heavily dominated avian communities with low species richness, while non-invaded areas exhibit highly-even and species-rich bird communities ([MacGregor-Fors et al., 2010](#)). These might be the case for *E. astrild* and *P. domesticus* in our study region, but more studies are needed to quantify their invasiveness, if any.

On the other hand, the high representation of the native Thraupidae and Tyrannidae was expected ([Hilty, 2011](#)), given that both are commonly found in urban parks, squares, gardens and other altered habitats. Tyrannidae is considered the largest family of the Western Hemisphere, with 413 species representing approximately 18% of the passerines known for South America ([Sick, 1997](#); [Fitzpatrick, 2004](#)). Thraupidae is less speciose (240 spp.), but is also very common along the Brazilian Atlantic Forest. In our study region, one in four species belongs to these two families. These birds usually feed fruits, seeds and insects ([Lopes et al., 2005](#)). The frugivorous species usually consume small seeds and fruits (<1.5 cm length), favoring the dispersal and proliferation of plant species that bear such attributes. These plant attributes are generally correlated with soft wood, fast growth, copious production of small seeds, and low capability to store carbon, being typical in pioneer tree species that do not characterize old-growth conserved forests ([Santos et al., 2012](#)). The dominance of Thraupidae and Tyrannidae might therefore help to maintain the flora at early to mid successional stages and boost the regional floristic homogenization driven by the proliferation of native tree species ([Lôbo et al., 2011](#)).

Regardless this dominance of Thraupidae and Tyrannidae, the species similarity varied considerably across the protected areas, following our expectation that less isolated, similar-sized areas showed an avifauna more similar to each other than more isolated, different-sized areas. This suggests certain level of dispersal limitation to more distant sites and major environmental differences between small and large areas. Both possibilities have been reported elsewhere ([Sekercioglu et al., 2002](#); [Uezu et al., 2005](#); [White et al., 2005](#)). However, the weakness of the relationships, though statistically significant, indicates that species composition are also influenced by factors other than size and isolation. Based on field observations, we hypothesize that varying levels of hunting, catching for illegal trade, pressure by exotic predators (e.g. cats and dogs) and noise pollution also play important role in taxonomic composition.

Implications for management and conservation

In addition to more investment in basic infrastructure and full-time staff ([Silva-Junior and Santos, 2017](#)), we strongly recommend the responsible agencies to follow the federal law 9.985/2000 and create the management council and the management plan of the four areas officially recognized as 'conservation units'. Lawfully, these actions should have been taken in the five years after creation, but to date only the Cabedelo Sandbank National Forest is in conformity with legislation. This legal noncompliance threatens the protected areas with downgrading, downsizing and degazettement ([Bernard et al., 2014](#)) and make public use of their ecosystem services almost unfeasible. Three municipal protected areas, which are not yet recognized as conservation units, face even more legal insecurity as their creation act may be revoked more easily. In this sense, we also recommend the official recognition of these areas as conservation units under federal law 9.985/2000.

Despite these challenges, the remarkable bird diversity of small and large urban reserves suggest that birdwatching for contemplative and educational purposes may help to change the course of conservation in the region, provided that it is well planned and based on the current legislation. Birdwatching is the act of

observing and identifying birds in their native habitats; it is becoming the most rapidly growing and most environmentally conscious segment of ecotourism and provides economic hope for many threatened natural areas around the world (Sekercioglu, 2002). Positive impacts of birdwatching include link between avian diversity and local income, financial incentive to conserve wildlife, less impact and more income than typical tourism, increased local control due to unique bird species, visitation of areas outside traditional tourist itineraries, protection of unprotected areas with desired species, valuation of local natural history knowledge, education and employment of local guides, generation of funds for bird conservation and contribution to ornithological knowledge (Sekercioglu, 2002). Negative impacts involve disturbing birds by playing tapes and by approaching, increased nest predation and nest abandonment, increased disturbance of rare and/or threatened birds, visitor-related pollution and habitat destruction, cash leaks from local communities, resentment by excluded locals and cultural degradation associated with tourism (Sekercioglu, 2002).

Following Sekercioglu's recommendations for optimal birdwatching, we suggest that birdwatching initiatives in ours and other urbanizing regions of the Brazilian Atlantic Forest adhere to and insist on ethical birding conduct, avoid nests and young birds as much as possible, show particular care with threatened and rare species, minimize tape use and try to minimize being seen, do not approach further once a bird notices the watcher, stick to established roads/trails/walkways, use scopes for observation and photography, educate locals about birds and their financial benefits, support local and low-impact establishments and contribute to non-governmental organizations active in bird conservation. Providing a comprehensive birdwatching planning extrapolates the limits of this study, but its principles are presented here. We hope conservationists and decision makers find this information useful and give more attention to the urban reserves under their responsibility.

Acknowledgments

This work was supported by the National Council for Scientific and Technological Development (CNPq) (grant numbers 476135/2013-3 and 310340/2016-0). We thank the Higher Education Personnel Improvement Coordination (CAPES) for the scholarship to the TER and the UFPB Graduate Program in Biodiversity for institutional support. We also thank the Chico Mendes Institute for Biodiversity Conservation (ICMBio) and the Paraíba Environment Administration (SUDEMA) for authorization to carry out this work within the protected areas. We are grateful to Pamela Stevens for producing the figure of the study area, Orione Alvares da Silva for discussing ideas and supporting fieldwork, and two anonymous reviewers for their contributions and suggestions on earlier versions of this manuscript.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at [doi:10.1016/j.pecon.2017.11.001](https://doi.org/10.1016/j.pecon.2017.11.001).

References

- Anjos, L., Collins, C.D., Holt, R.D., Volpato, G.H., Lopes, E.V., Bochio, G.M., 2015. Can habitat specialization patterns of Neotropical birds highlight vulnerable areas for conservation in the Atlantic rainforest, Southern Brazil? *Biol. Conserv.* 188, 32–40.
- Bernard, E., Pena, L.A.O., Araújo, E., 2014. Downgrading, downsizing, degazettement, and reclassification of protected areas in Brazil. *Conserv. Biol.* 28, 939–950.
- Bibby, C., Jones, M., Marsden, S., 1998. *Expedition Field Techniques: Bird Surveys*. Expedition Advisory Centre, Royal Geographical Society, London.
- Clarke, K.R., Gorley, R.N., 2006. *PRIMER v6: User Manual/Tutorial*. PRIMER-E, Plymouth.
- Colwell, R.K., 1997. Estimates: Statistical Estimation of Species Richness and Shared Species from Samples, <http://viceroy.eeb.uconn.edu/estimates> (accessed 05.07.2015).
- Erize, F., Mata, J.R.R., Rumboll, M., 2006. *Birds of South America, Non Passerines: Rheas to Woodpeckers*. Princeton University Press, Princeton.
- Fitzpatrick, J.W., 2004. Family Tyrannidae (tyrant-flycatchers). In: Hoyo, J., Elliott, A., Christie, D.A. (Eds.), *Handbook of the Birds of the world: Cotingas to pipits and wagtails*, Vol. 9. Lynx Edicions, Barcelona, pp. 170–462.
- Ford, H.A., Barrett, G.W., Saunders, D.A., Recher, H.F., 2001. Why have birds in the woodlands of Southern Australia declined? *Biol. Conserv.* 97, 71–88.
- Hilty, S.L., 2011. Family Thraupidae (Tanagers). In: Hoyo, J., Elliott, A., Christie, D.A. (Eds.), *Handbook of the Birds of the world: Tanagers to New World Blackbirds*, Vol. 16. Barcelona, Lynx Edicions, pp. 46–329.
- Joly, C.A., Metzger, J.P., Tabarelli, M., 2014. Experiences from the Brazilian Atlantic Forest: ecological findings and conservation initiatives. *New Phytol.* 204, 459–473.
- Jones, H.P., Tershry, B.R., Zavaleta, E.S., Croll, D.A., Keitt, B.S., Finkelstein, M.E., Howald, G.R., 2008. Severity of the effects of invasive rats on seabirds: a global review. *Conserv. Biol.* 22, 16–26.
- Lobo-Araújo, L.W., Toledo, M.T.F., Efe, M.A., Malhado, A.C.M., Vital, M.V.C., Toledo-Lima, G.S., Macario, P., Santos, J.G., Ladle, R.J., 2013. Bird communities in three forest types in the Pernambuco Centre of Endemism, Alagoas, Brazil. *Iheringia. Série Zool.* 103, 85–96.
- Lôbo, D., Leão, T., Melo, F.P.L., Santos, A.M.M., Tabarelli, M., 2011. Forest fragmentation drives Atlantic Forest of northeastern Brazil to biotic homogenization. *Divers. Distrib.* 17, 287–296.
- Lopes, L.E., Fernandes, A.M., Marini, M.A., 2005. Diet of some Atlantic forest birds. *Ararajuba* 13, 95–103.
- MacGregor-Fors, I., Morales-Perez, L., Quesada, J., Schondube, J.E., 2010. Relationship between the presence of house Sparrows (*Passer domesticus*) and Neotropical Bird community structure and diversity. *Biol. Invasions* 12, 87–96.
- Manhães, M.A., Loures-Ribeiro, A., 2011. The avifauna of the Poço D'Anta municipal biological reserve, Juiz de Fora, MG. *Biota Neotrop.* 11, 275–286.
- Marinho, M.F.A., (M. Sc. dissertation) 2014. *Aves da Paraíba: uma revisão de informações históricas e atuais*. Universidade Federal da Paraíba, Areia.
- Marini, M.A., Garcia, F.I., 2005. Conservação de aves no Brasil. *Megadiversidade* 1, 95–102.
- Millennium Ecosystem Assessment, 2005. *Ecosystems and Human Well-Being: Synthesis*. Island Press, Washington.
- Morante-Filho, J.C., Arroyo-Rodríguez, V., Andrade, E., Santos, B.A., Cazetta, E., Faria, D., 2017. Compensatory dynamics maintain bird phylogenetic diversity in fragmented tropical landscapes. *J. Appl. Ecol.*, <http://dx.doi.org/10.1111/1365-2664.12962> (in press).
- Myers, N., Mittermeier, R.A., Mittermeier, C.G., Fonseca, G.A.B., Kent, J., 2000. Biodiversity hotspots for conservation priorities. *Nature* 403, 853–858.
- Piacentini, V., de, Q., Aleixo, A., Agne, C.E., Maurício, G.N., Pacheco, J.F., Bravo, G.A., Brito, G.R.R., Naka, L.N., Olmos, F., Posso, S., Silveira, L.F., Betini, G.S., Carrano, E., Franz, I., Lees, A.C., Lima, L.M., Pioli, D., Schunck, F., Amaral, F.R.do, Bencke, G.A., Cohn-Haft, M., Figueiredo, L.F.A., Straube, F.C., Cesari, E., 2015. Annotated checklist of the birds of Brazil by the Brazilian Ornithological Records Committee/Lista comentada das aves do Brasil pelo Comitê Brasileiro de Registros Ornitológicos. *Rev. Bras. Ornitol.* 23, 91–298.
- Porto, K.C., Almeida-Cortez, J.S., Tabarelli, M., 2005. *Diversidade Biológica e Conservação da Floresta Atlântica ao Norte do Rio São Francisco*. Ministério do Meio Ambiente, Brasília.
- Ribeiro, M.C., Metzger, J.P., Martensen, A.C., Ponzoni, F.J., Hirota, M.M., 2009. The Brazilian Atlantic Forest: how much is left and how is the remaining forest distributed? Implications for conservation. *Biol. Conserv.* 142, 1141–1153.
- Ribon, R.J., 2010. Amostragem de aves pelo método de listas de Mackinnon. In: Von Matter, S., Straube, F., Accordi, I., Piacentini, V., Cândido-Jr, J.F. (Eds.), *Ornitologia e conservação: ciência aplicada, técnicas de pesquisa e levantamento*. Technical Books, Rio de Janeiro, pp. 33–46.
- Roda, S.A., Pereira, G.A., Albano, C., 2011. Conservação de aves endêmicas e ameaçadas do centro de endemismo Pernambuco. Editora Universitária da UFPE, Recife.
- Santos, G.G.A., Santos, B.A., Nascimento, H.E.M., Tabarelli, M., 2012. Contrasting demographic structure of short- and long-lived pioneer tree species on Amazonian Forest Edges. *Biotropica* 44, 771–778.
- Sekercioglu, C.H., 2002. Impacts of birdwatching on human and avian communities. *Environ. Conserv.* 29, 282–289.
- Sekercioglu, C.H., Ehrlich, P.R., Daily, G.C., Aygen, D., Goehring, D., Sandy, R.F., 2002. Disappearance of insectivorous birds from tropical forest fragments. *Proc. Natl. Acad. Sci. U. S. A.* 99, 263–267.
- Seto, K.C., Parnell, S., Elmqvist, T., 2013. A global outlook on urbanization. In: Elmqvist, T., Fragkias, M., Goodness, J., Güneralp, B., Marcotullio, P.J., McDonald, R.I., Parnell, S., Haase, D., Sendstad, M., Seto, K.C., Wilkinson, C. (Eds.), *Urbanization, Biodiversity and Ecosystem Services: Challenges and Opportunities*. Springer Netherlands, Dordrecht, pp. 1–12.
- Shaw, L., Chamberlain, D., Evans, M., 2008. The House Sparrow *Passer domesticus* in urban areas: reviewing a possible link between post-decline distribution and human socio-economic status. *J. Ornithol.* 149, 293–299.
- Sick, H., 1997. *Ornitologia Brasileira*. Nova Fronteira, Rio de Janeiro.
- Sigrist, T., 2009. *Aves do Brasil Oriental*. Avis Brasilis, São Paulo.

- Silva-Junior, V., Santos, B.A., 2017. Using environmental perception and local knowledge to improve the effectiveness of an Urban Park in Northeast Brazil. *Ethnobiol. Conserv.* 6, 2, <http://dx.doi.org/10.15451/ec2017-03-6.2-1-24>.
- Silva, F.C., Silva, G.G., Chagas, M.O., Jung, D.M.H., 2014. *Composição da comunidade de aves em área urbana no sul do Brasil*. *Neotrop. Biol. Conserv.* 9, 78–90.
- Stevens, P.O., (M. Sc. dissertation) 2014. *Dinâmica da Paisagem no Geossistema do Estuário do Rio Paraíba—Extremo Oriental das Américas: Estimativas de Perdas de habitat e cenários de recuperação da Biodiversidade*. Universidade Federal da Paraíba, João Pessoa.
- Sol, D., Bartomeus, I., González-Lagos, C., Pavoine, S., 2017. *Urbanisation and the loss of phylogenetic diversity in birds*. *Ecol. Lett.* 20, 721–729.
- Uezu, A., Metzger, J.P., Vielliard, J.M.E., 2005. *Effects on structural and functional connectivity and patch size on the abundance of seven Atlantic Forest bird species*. *Biol. Conserv.* 123, 507–519.
- United Nations, 2014. *World Urbanization Prospects: the 2014 revision*. Department of Economic and Social Affairs, United Nations, New York.
- White, J., Antos, M., Fitzsimons, J., Palmer, G., 2005. *Non-uniform bird assemblages in urban environments: the influence of streetscape vegetation*. *Landsc. Urban Plan.* 71, 123–135.