



Policy Forums

Can you hear the noise? Environmental licensing of seismic surveys in Brazil faces uncertain future after 18 years protecting biodiversity

Cristiano Vilardo^{a,b,*}, André Favaretto Barbosa^a^a IBAMA – Brazilian Institute of the Environment and Renewable Natural Resources, Rio de Janeiro, Brazil^b Energy Planning Program, Alberto Luiz de Coimbra Engineering Graduate School and Research Center, Federal University of Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil

ARTICLE INFO

Article history:

Received 8 August 2017

Accepted 27 November 2017

Available online 11 January 2018

Keywords:

Environmental licensing

Environmental impact assessment

Biodiversity monitoring

Seismic surveys

Marine biodiversity

ABSTRACT

Marine Seismic Surveys are an important source of concern for marine biodiversity conservation worldwide. In Brazil, Environmental Federal Agency IBAMA has developed a considerably advanced mitigation/monitoring requirements package in 18 years of environmental licensing practice, with standardized guidelines since 2005. Adding to global efforts aiming at filling knowledge gaps over the impacts on biodiversity, IBAMA has been able to foster important marine research through environmental licensing requirements. Better communication of research findings to the international scientific community remains a challenge to be addressed. Nevertheless, current institutional and legal reforming initiatives jeopardize the evolution of environmental control of Marine Seismic Surveys in Brazil.

© 2018 Associação Brasileira de Ciência Ecológica e Conservação. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Marine Seismic Surveys affects marine biodiversity

In the last decades, the marine scientific community has developed an increasing concern over the anthropogenic noise intensification in the oceans as an important threat to biodiversity and animal welfare (e.g., Cummings and Brandon, 2004; IWC SC, 2012, 2013, 2014, 2015, 2016, 2017; Popper and Hawkins, 2016, 2012; Simmonds et al., 2014; Williams et al., 2015). Very far from Jacques Cousteau's 'Silent World', our oceans are a realm of sound – which travels far better underwater than light does. In the oceans, underwater sound becomes a critical resource for marine fauna communication, orientation and overall spatial perception (Jasny et al., 2005; Simmonds et al., 2004). The main anthropogenic sources of noise are related to shipping, oil and gas exploration and production, naval sonar, military operations, fishing and research (Harris et al., 2017; Hildebrand, 2009). Among these activities, Marine Seismic Surveys (MSS) are often cited as one of the largest contributors to this scenario.

Marine Seismic Surveys are a widespread method used in the offshore petroleum industry to locate potential areas of exploratory interest. The seismic techniques use sound to probe the Earth

and help geophysicists develop geological models of the sedimentary basins. In marine exploratory surveys, underwater sound generation is generally accomplished through a quick release of pressurized air from metal cylinders known as airguns (Dragoset, 2000). Seismic specialized vessels tow arrays of airguns that fire simultaneously in regular intervals (10–15 s) to produce high-intensity sound pulses that propagate down the seabed (Caldwell and Dragoset, 2000) and interact with the different geological layers. The reflecting echoes are captured by towed hydrophone arrays and processed to form the underground images that are used to guide the planning of wells drilling.

While there are still significant knowledge gaps on the impacts seismic surveys can have upon marine life, there is a growing body of literature demonstrating its potential harmful effects on different taxa. Reviews can be found summarizing state-of-the-art research for marine mammals (e.g., Nowacek et al., 2015, 2007; Southall et al., 2007; Weilgart, 2007), sea turtles (Nelms et al., 2016), fishes (Popper and Hastings, 2009; Slabbekoorn et al., 2010), invertebrates (Moriyasu et al., 2004), and cephalopods (André et al., 2011). Several research papers also describe important experimental results for understanding either physical damage or behavioral effects in cetaceans (e.g., Blackwell et al., 2015; Castellote et al., 2012; Cerchio et al., 2014; Dunlop et al., 2015; Miller et al., 2009; Nieukirk et al., 2012), plankton (e.g., McCauley et al., 2017), sea turtles (e.g., DeRuiter and Larbi Doukara, 2012), fish – auditory damage (McCauley et al., 2003), fish – behavior (e.g., Hassel et al., 2004; Paxton et al., 2017; Slotte et al., 2004; Wardle et al., 2001), and

* Corresponding author at: Praça XV de Novembro, 42 – 9th Floor, Centro, Rio de Janeiro CEP 20.010-010, RJ, Brazil.

E-mail address: cristianovilardo@gmail.com (C. Vilardo).

fish – behavior and catch rates (Engås and Løkkeborg, 1996; Lokkeborg et al., 2012; Streever et al., 2016).

MSS environmental licensing and biodiversity in Brazil

Brazil's first 2D Marine Seismic Survey was conducted in 1957, in Alagoas State continental shelf, whereas the pioneer 3D survey (denser, contiguous lines of survey) took place in 1978, in Cherne Field in Campos Basin (de Mendonça et al., 2004). However, the intensification of Brazilian coast exploration dates back to the opening of the market to foreign companies in 1997, through Federal Law 9.478/97. As a consequence of that opening, the regulatory institutional framework evolved accordingly with the creation of the National Petroleum Agency (ANP) and the environmental licensing office in the Federal Environmental Agency (IBAMA). In Brazil, the environmental licensing is an administrative procedure based upon the environmental impact assessment (EIA) process and applied to potentially harmful activities. The first Marine Seismic Survey licensing application IBAMA reviewed was in early 1999 – probably the first time a MSS went through any environmental oversight in Brazil – a process that is now entering its age of majority at 18 years.

In this period, IBAMA had to develop the expertise to understand and deal with the environmental consequences of Marine Seismic Surveys, as well as the social conflicts eventually associated with the activities. Initially running with the aid of consultants, it was not until 2002 that the first public officers were hired to IBAMA's Oil and Gas Licensing Office (Vilardo, 2007), enabling actual institutional learning and advancement of procedures. Those early days of environmental licensing practice of MSS in Brazil were characterized by a high degree of uncertainty and a very controversial debate between industry and regulators over its potential effects.

IBAMA's strategy to deal with this scenario was two-fold. The first path was to take advantage of the globalized nature of the seismic industry and develop national mitigation guidelines based on international best practices. Those guidelines were first issued in 2005 (IBAMA, 2005) and have been evolving since then using feedback from the field practice. Besides establishing mitigation measures for protection of marine mammals and sea turtles, the guidelines also provided a standardized reporting framework for the monitoring records. This means that data on the occurrence and distribution of marine mammals in Brazilian waters are being recorded in a standardized way since 2005, using trained professionals with a marine biology, oceanography or similar degree. This collection has more than 8000 individual sightings so far, producing information of great value to conservation that would never be available otherwise. Taking into account the potential data quality issues attached to such datasets (observer mistakes, for example), the recorded observations may enable scientific analysis in a variety of ways: as single observations (e.g., Fernandes et al., 2007), using a MSS as sampling unit (e.g., Gurjão et al., 2004) or even a more longitudinal assessment, like Stone and Tasker (2006) did with data from 201 surveys in the United Kingdom.

A great step toward using this register for decision-making and conservation planning was the migration of the IBAMA database to SIMMAM (Univali, 2017), an online opensource platform that integrates sightings and strandings data from the institutions that are part of the Brazilian Aquatic Mammals Stranding Network – REMAB (Barreto et al., 2012) and from academic research groups. The platform is free, and while individual researchers are allowed to establish secrecy of the marine mammals data they input in SIMMAM, all MSS-related data is considered public information and can be accessed by anyone.

Another highlight of the Brazilian MSS mitigation regulation is the existence of seasonal restriction areas for the protection

of sensitive behaviors of marine mammals and sea turtles. In a context of growing concern among the marine conservation community, IBAMA pioneered this kind of approach since 2003, when a Humpback whales' (*Megaptera novaeangliae*) breeding ground was defined as out of reach for seismic surveys from July to November. After this first experience, other restricted areas and periods were developed for protection of Southern Right whales (*Eubalaena australis*), Franciscana dolphins (*Pontoporia blainvillei*), Manatee (*Trichechus manatus*), and sea turtles (main nesting areas of the five species that occur in Brazil). After several years of practical implementation, those closed areas for MSS were finally absorbed into formal regulation in 2011 (IBAMA/ICMBio, 2011a,b). To date, Brazilian guidelines appear to remain one of the few in the world to establish such closed areas – certainly the only in Latin America (Reyes Reyes et al., 2016) – despite the ample recognition of the effectiveness of such measures. Those closed seasons and the mitigation/monitoring guidelines have put Brazil in a highlighted position among international practice (see, for instance, Compton et al., 2008; GHFS, 2015; Reyes Reyes et al., 2016; Weir and Dolman, 2007).

The second path IBAMA trailed to deal with the uncertainties of Marine Seismic Surveys impacts was to foster a national contribution for the international effort aimed to address the existing knowledge gaps. Once the EIA process for offshore activities was usually based on secondary data (due to budgetary and time restrictions), IBAMA's approach involved the commissioning of research initiatives as conditions for the licenses granted. The scope of the research to be commissioned depended on the characteristics of the project being assessed and the area in which it would be developed.

While some of the research initiatives commissioned to date were single undertakings, destined to address a specific question involving seismic surveys impacts on biodiversity, others became regular monitoring programs for areas of higher environmental sensitivity, like the beaches monitoring projects, for example.

Table 1 summarizes the main research initiatives commissioned by IBAMA's request in Marine Seismic Surveys EIA processes to date. One important shortcoming to be highlighted is the low amount of peer-reviewed publications derived from those initiatives. This means that most of the knowledge generated in this process is trapped inside Technical Reports submitted to IBAMA to fulfill administrative obligations – despite the fact that there are several results of potential international relevance. One desired outcome of this short paper is exactly to foster academic interest by providing a useful index of those initiatives. All the information within such reports are considered of public domain and, therefore, can be accessed in IBAMA's EIA documentation center. Unfortunately, some of the older data will not be available in digital formats, but nevertheless it could be retrieved from the printed reports with some effort.

There are, of course, risks and limitations of the knowledge generation embedded in environmental impact assessment processes. Obtaining scientific valid data from monitoring programs depends on several variables, like proper sampling design, adequate survey effort and methodological consistency, as recently argued by Dias et al. (2017). Nevertheless, we believe that the vast majority of the results generated through MSS EIA-related projects can be considered good science and would constitute publishable material if given proper academic treatment.

Other important topic of concern that is worth mentioning is the potential bias that industry funding can impose over such studies, however closely overseen they might be by governmental institutions like IBAMA. Wade et al. (2010) found evidence of bias in papers assessing anthropogenic noise effects in marine mammals, depending on which institution funded the research. One tentative way to minimize this kind of problem would be to prioritize public universities and research centers as implementing institutions,

Table 1
Summary of the main research initiatives commissioned through IBAMA's environmental licensing requirements. All the Environmental Licensing Reports cited are available in the Documentation Center of the IBAMA Oil and Gas Office, located in Rio de Janeiro. Basins: BAR–Barreirinhas, CA–Camamu/Almada, ES–Espírito Santo, PAMA–Pará/Maranhão, PEPB–Pernambuco/Paraíba, POT–Potiguar, S–Santos, and SEAL–Sergipe/Alagoas.

Studied groups	Basin	Type of approach	Type of analysis	References
Shrimps, lobsters, fish	CA	Controlled exposure of caged animals using small airgun array (360 in ³) in shallow waters.	Visual and histopathological damage assessment	GIA/UFPR (2002)
Macrozooplankton	CA	Controlled exposure of zooplankton in nets using small airgun array (360 in ³) in shallow waters.	Mortality rate comparison	GIA/UFPR (2002)
Shrimps	CA	Controlled fishery (trawling)	Catch rate comparison	Andriguetto-Filho et al. (2005), GIA/UFPR (2002)
Zooplankton	CA	Controlled exposure of zooplankton in nets using small airgun array (360 in ³) in shallow waters.	Mortality rate comparison	GIA/UFPR (2004)
Fish	CA	Controlled exposure of caged animals using small airgun array (360 in ³) in shallow waters.	Behavioral (video) and histopathological damage assessment	Boeger et al. (2006), GIA/UFPR (2004)
Fish	CA	Controlled exposure of caged and free animals using full airgun array (3090 in ³) in shallow waters.	Behavioral (video) and histopathological damage assessment	GEIA/UFPR and Everest (2004)
Zooplankton	ES	Controlled plankton tows at 10 m of lateral distance of a full airgun array (5085 in ³)	Mortality rate comparison	Thalassa and Ecology Brasil (2009)
Fish	ES	Controlled exposure of fish living around a fixed platform using full airgun array (5085 in ³)	Video assessment of behavior and abundance	UFES and Petrobras (2012)
Fish (<i>Coryphaena hippurus</i>)	ES	Controlled fishery (longline)	Catch rate comparison	UFRJ et al. (2015)
Cetaceans, birds, fish	POT	Oceanographic cruise in seamount region	Occurrence of species	Engeo (2017)
Sea turtles (<i>L. olivacea</i> and <i>C. caretta</i>)	SEAL	Satellite telemetry of females	Inter-nesting and migratory behavior	Engeo et al. (2017)
Sea turtles (<i>Eretmochelys imbricata</i>)	POT	Satellite telemetry of females	Inter-nesting and migratory behavior	Engeo and Centro TAMAR-ICMBio (2016)
Humpback whales (<i>Megaptera novaeangliae</i>)	ES	Satellite telemetry of migrating whales	Migratory behavior	Instituto Aqualie (2017)
Cetaceans	S	Aerial monitoring	Occurrence, distribution and behavior of species	UESC and Instituto Aqualie (2012)
Marine mammals and Sea turtles	All	Visual monitoring aboard seismic vessels	Occurrence, distribution and behavior of species	Every MSS Environmental Licensing Report in Brazil since 1999
Marine mammals	All	Passive acoustic monitoring aboard seismic vessels	Occurrence, distribution and behavior of species	Every MSS Environmental Licensing Report in Brazil since 2015
Marine mammals and Sea turtles	BAR, PAMA, PEPB	Beaches monitoring	Stranding patterns and causa-mortis	e.g., Amares (2017), Amares and BioMA (2016), IMA (2016), NAV (2010), Soma (2009)

aiming at their independence and public commitment. But unfortunately, practice has shown that this does not completely solve the bias issue – as there are several ways in which the sponsor can influence the outcomes of the research. It can range from direct pressure or even censorship to more subtle means, as the potential for future contracts with the same contractor. Adding to the complexity, university-led projects in Brazil are said to be harder to manage (due to cumbersome administrative bureaucracy) and not always yield the follow-up peer-reviewed publications as expected, due to competing priorities in the Academy (like the need to secure the next grant or contract in a chronically underfunded research environment).

A different approach to minimize possible bias and to avoid environmental licensing constraints of time and scope would be to establish research funds managed by an independent body. This financial mechanism should have a governance design to ensure fair selection of implementation partners and project accountability. There are several examples of such arrangements in Brazil and elsewhere, with specific purposes and formats, and the designing of a tailor-made mechanism to fund MSS-related research should learn from those experiences.

Concluding remarks and reforming risks

Marine Seismic Surveys have been a major source of concern for marine biodiversity conservationists in the past decades. It is critical that the knowledge over its effects on marine life continue to improve – and Brazil's 18-year regulatory experience and well acknowledged mitigation requirements puts the country in a privileged position to contribute to that objective. There have been many relevant research initiatives commissioned through the environmental licensing process so far – but a better communication of the results is still a challenge to be addressed.

Despite all the evolution summarized in this paper, environmental control of Marine Seismic Surveys in Brazil seems endangered from both external (legal) and internal (institutional) menaces. One important external source of risk is the systemic reforming proposals that are currently being discussed in the National Congress, which can severely undermine effectiveness of Brazilian licensing system (see discussion in Bragagnolo et al., 2017; Fonseca et al., 2017), threatening the environmental oversight of activities like MSS. We fear that a new regulation would impose an over-streamlined process, as seen recently in some Brazilian states (Fonseca and Rodrigues, 2017) and in other countries

(Bond et al., 2014), reducing IBAMA's ability to promote effective mitigation and adequate public participation in the environmental licensing decision-making.

The other menace is internal and maybe harder to understand and identify its origin. An institutional reform in IBAMA's Environmental Licensing Directorate (DILIC) has just dismembered the Oil and Gas Office (CGPEG) in Rio de Janeiro, removing the MSS licensing responsibility to the central DILIC office in Brasília, without any reasonable justification. The newly published internal regiment places seismic surveys activities along with ports, marine minerals mining and other marine infrastructure in a Coordination in DILIC's central office in Brasília, while keeping the other activities of the oil and gas chain (drilling and production) in Rio de Janeiro, where the Oil and Gas Office is located since 1998. The experienced staff of the Seismic Surveys team are to be redistributed to other activities in Rio de Janeiro, while IBAMA creates from zero a new team in Brasília office. This obscure management option seems even more unusual in a time of severe budget constraints, when there are no funds available for capacity building and new public tenders for staff increasing are practically out of the question.

This hard-to-explain change has puzzled most of the actors involved in MSS environmental licensing process, gathering important support from outside IBAMA, as shows an open letter signed by 24 researchers and institutions involved in marine biodiversity conservation in Brazil (Magalhães et al., 2017). Despite being already formalized in IBAMA's new internal regiment, the described changes were still not effective at the time of writing (August 2017), but it would certainly be a huge loss to see this 18-year old legacy becoming history for some idiosyncratic reforming. In difficult times such as the present, it is of paramount importance that we preserve the positive experiences and halt the throwbacks in the conservation agenda.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Conflicts of interest

None.

Acknowledgments

The authors would like to express gratitude and acknowledge the great contribution of several Environmental Analysts that have worked in IBAMA's Oil and Gas Office in these 18 years and showed that, despite all the challenges and constraints, it is possible to achieve excellence in the public service in Brazil. We also thank the numerous researchers, analysts and marine mammal observers (MMO) that have contributed to generate science from the environmental licensing process.

References

- Amares, 2017. *Projeto de Monitoramento de Praias – Atividade de Pesquisa Sísmica Marítima 3D, não Exclusiva, na Bacia de Barreirinhas, Projeto Bar Fases Unificadas. Relatório Final. Instituto Amares Pesquisa e Conservação de Ecossistemas Aquáticos. BioMA – Biologia e Conservação de Mamíferos Aquáticos da Amazônia. Processo IBAMA 02022.000605/2013 (DOC24), Portuguese.*
- Amares, BioMA, 2016. *Projeto de Monitoramento de Praias (PMP) – Pesquisa Sísmica Marítima 3D, não-exclusiva, Projeto PAMA, bacia Pará-Maranhão. Relatório Final. Instituto Amares Pesquisa e Conservação de Ecossistemas Aquáticos. BioMA – Biologia e Conservação de Mamíferos Aquáticos da Amazônia. Processo IBAMA 02022.000015/2014 (DOC13), Portuguese.*
- André, M., Solé, M., Lenoir, M., Durfort, M., Quero, C., Mas, A., Lombarte, A., van der Schaar, M., López-Bejar, M., Morell, M., Zaug, S., Houégnigan, L., 2011. Low-frequency sounds induce acoustic trauma in cephalopods. *Front. Ecol. Environ.* 9, 489–493, <http://dx.doi.org/10.1890/100124>.
- Andriguetto-Filho, J.M., Ostrensky, A., Pie, M.R., Silva, U.A., Boeger, W.A., 2005. Evaluating the impact of seismic prospecting on artisanal shrimp fisheries. *Cont. Shelf Res.* 25, 1720–1727, <http://dx.doi.org/10.1016/j.csr.2005.05.003>.
- Barreto, A.S., Sperb, R.M., Barbosa Jr., A.F., da Silva, J.M., 2012. Simmam – Sistema De Apoio Ao Monitoramento De Mamíferos Marinhos: Uma Nova Ferramenta Para a Gestão Ambiental (Marine Mammals Monitoring Support System: a new tool for environmental management). *An. da Rio Oil Gas Expo Conf.*, 1–7 http://simmam.acad.univali.br/site/wp-content/uploads/2014/09/IBP1465_12.pdf, Portuguese.
- Blackwell, S.B., Nations, C.S., McDonald, T.L., Thode, A.M., Mathias, D., Kim, K.H., Greene, C.R., Macrander, A.M., 2015. Effects of airgun sounds on bowhead whale calling rates: evidence for two behavioral thresholds. *PLoS ONE* 10, e0125720, <http://dx.doi.org/10.1371/journal.pone.0125720>.
- Boeger, W.A., Pie, M.R., Ostrensky, A., Cardoso, M.F., 2006. The effect of exposure to seismic prospecting on coral reef fishes. *Note Br. J. Oceanogr.* 54, 235–239, <http://dx.doi.org/10.1590/S1679-87592006000300007>.
- Bond, A., Pope, J., Morrison-Saunders, A., Retief, F., Gunn, J.A.E., 2014. Impact assessment: Eroding benefits through streamlining? *Environ. Impact Assess. Rev.* 45, 46–53, <http://dx.doi.org/10.1016/j.eiar.2013.12.002>.
- Bragagnolo, C., Carvalho Lemos, C., Ladle, R.J., Pellin, A., 2017. Streamlining or sidestepping? Political pressure to revise environmental licensing and EIA in Brazil. *Environ. Impact Assess. Rev.* 65, 86–90, <http://dx.doi.org/10.1016/j.eiar.2017.04.010>.
- Caldwell, J., Dragoset, W., 2000. A brief overview of seismic air-gun arrays. *Lead. Edge* 19, 898–902, <http://dx.doi.org/10.1190/1.1438744>.
- Castellote, M., Clark, C.W., Lammers, M.O., 2012. Acoustic and behavioural changes by fin whales (*Balaenoptera physalus*) in response to shipping and airgun noise. *Biol. Conserv.* 147, 115–122, <http://dx.doi.org/10.1016/j.biocon.2011.12.021>.
- Cerchio, S., Strindberg, S., Collins, T., Bennett, C., Rosenbaum, H., 2014. Seismic surveys negatively affect humpback whale singing activity off Northern Angola. *PLoS ONE* 9, <http://dx.doi.org/10.1371/journal.pone.0086464>.
- Compton, R., Goodwin, L., Handy, R., Abbott, V., 2008. A critical examination of worldwide guidelines for minimising the disturbance to marine mammals during seismic surveys. *Mar. Policy* 32, 255–262, <http://dx.doi.org/10.1016/j.marpol.2007.05.005>.
- Cummings, J., Brandon, N., 2004. Sonic impact: a precautionary assessment of noise pollution from ocean seismic surveys. *Greenpeace* <http://www.greenpeace.org/usa/wp-content/uploads/2013/03/SonicImpactReport.pdf>.
- DeRuiter, S., Larbi Doukara, K., 2012. Loggerhead turtles dive in response to airgun sound exposure. *Endanger. Species Res.* 16, 55–63, <http://dx.doi.org/10.3354/esr00396>.
- Dias, A.M.S., Fonseca, A., Paglia, A.P., 2017. Biodiversity monitoring in the environmental impact assessment of mining projects: a (persistent) waste of time and money? *Perspect. Ecol. Conserv.* 15, 206–208, <http://dx.doi.org/10.1016/j.pecon.2017.06.001>.
- Dragoset, B., 2000. Introduction to air guns and air-gun arrays. *Lead. Edge* 19, 892–897, <http://dx.doi.org/10.1190/1.1438741>.
- Dunlop, R.A., Noad, M.J., McCauley, R.D., Kniest, E., Paton, D., Cato, D.H., 2015. The behavioural response of humpback whales (*Megaptera novaeangliae*) to a 20 cubic inch air gun. *Aquat. Mamm.* 41, 412–433, <http://dx.doi.org/10.1578/AM.41.4.2015.412>.
- Engås, A., Løkkeborg, S., 1996. Effects of seismic shooting on local abundance and catch rates of cod (*Gadus morhua*) and haddock (*Melanogrammus aeglefinus*). *Can. J. Fish. Aquat. Sci.* 53, 2238–2249, <http://dx.doi.org/10.1139/f96-177>.
- Engeo, 2017. *Relatório ambiental – Projeto Vertebrados. Relatório Final. Engeo Soluções Integradas. Processo IBAMA 02022.000148/2014 (DOC18), Portuguese.*
- Engeo, Centro TAMAR-ICMBio, 2016. *Projeto de monitoramento da Tartaruga-de-pente por telemetria satelital. Relatório parcial. In: Processos IBAMA 02022.002094/2013 (DOC19) e 02022.000148/2014 (DOC16), Portuguese.*
- Engeo, Everest, Centro TAMAR-ICMBio, Fundação Pró-TAMAR, 2017. *Projeto de monitoramento de quelônios por telemetria satelital. Relatório Final. In: Processo IBAMA 02022.001383/2013 (DOC11), Portuguese.*
- Fernandes, M.F., Cordeiro, A.S., Carvalho, D.M.R., Santos, W.R., Ramos, R., 2007. An interaction between a juvenile Clymene dolphin (*Stenella clymene*) and seismic survey vessel M/V Ramform Challenger-PGS, Bacia de Santos, Brazil. *Lat. Am. J. Aquat. Mamm.* 6, 189–192, <http://dx.doi.org/10.5597/lajam00124>.
- Fonseca, A., Rodrigues, S.E., 2017. The attractive concept of simplicity in environmental impact assessment: perceptions of outcomes in southeastern Brazil. *Environ. Impact Assess. Rev.* 67, 101–108, <http://dx.doi.org/10.1016/j.eiar.2017.09.001>.
- Fonseca, A., Sánchez, L.E., Ribeiro, J.C.J., 2017. Reforming EIA systems: a critical review of proposals in Brazil. *Environ. Impact Assess. Rev.* 62, 90–97, <http://dx.doi.org/10.1016/j.eiar.2016.10.002>.
- GEIA/UFPR, Everest, 2004. *Efeitos da sísmica com cabo flutuante em peixes tropicais de áreas recifais. Relatório Final. Grupo de Estudos de Impacto Ambiental – Universidade Federal do Paraná. Processo IBAMA 02022.008248/2000, Portuguese.*
- GHFS, 2015. *A review of seismic mitigation measures used along the coast of Northern South America, from North Brazil up to Colombia. Reference Document for the Steering group of the MaMa CoCo SEA. GHFS – Green Heritage Fund Suriname*

- http://www.car-spaw-rac.org/IMG/pdf/seismic_mitigation_measures_review_mamacosea.pdf.
- GIA/UFPR, 2004. Avaliação dos efeitos da sísmica com cabo de fundo sobre peixes recifais e sobre o zooplâncton marinho. Relatório Final. Grupo Integrado de Aquicultura e Estudos Ambientais – Universidade Federal do Paraná. Processo IBAMA 02022.004827/2002, Portuguese.
- GIA/UFPR, 2002. Avaliação dos impactos causados durante a aquisição de dados sísmicos sobre organismos marinhos de interesse comercial. Grupo Integrado de Aquicultura e Estudos Ambientais – Universidade Federal do Paraná. Processo IBAMA 02022.001056/2001, Portuguese.
- Hassel, A., Knutsen, T., Dalen, J., Skaar, K., Løkkeborg, S., Misund, O.A., Østensen, Ø., Fonn, M., Haugland, E.K., 2004. Influence of seismic shooting on the lesser sandeel (*Ammodytes marinus*). ICES J. Mar. Sci. 61, 1165–1173, <http://dx.doi.org/10.1016/j.icesjms.2004.07.008>.
- Gurjão, L.M., Freitas, J.E.F., Araújo, D.S., 2004. Sightings of dolphins during seismic surveys on the coast of Bahia state, Brazil. Lat. Am. J. Aquat. Mamm. 3, 171–175, <http://dx.doi.org/10.5597/lajam00063>.
- Harris, C.M., Thomas, L., Falcone, E.A., Hildebrand, J., Houser, D., Kvadsheim, P.H., Lam, F.-P.A., Miller, P.J.O., Moretti, D.J., Read, A.J., Slabbekoorn, H., Southall, B.L., Tyack, P.L., Wartok, D., Janik, V.M., 2017. Marine mammals and sonar: dose–response studies, the risk–disturbance hypothesis and the role of exposure context. J. Appl. Ecol. 1–9, <http://dx.doi.org/10.1111/1365-2664.12955>.
- Hildebrand, J., 2009. Anthropogenic and natural sources of ambient noise in the ocean. Mar. Ecol. Prog. Ser. 395, 5–20, <http://dx.doi.org/10.3354/meps08353>.
- IBAMA/ICMBio, 2011a. Instrução Normativa Conjunta nº 01-2011 [Joint Normative]. IBAMA – Brazilian Institute of the Environment and Renewable Natural Resources/ICMBio – Chico Mendes Institute of Biodiversity Conservation, Brazil. Portuguese.
- IBAMA/ICMBio, 2011b. Instrução Normativa Conjunta nº 02-2011 [Joint Normative]. IBAMA – Brazilian Institute of the Environment and Renewable Natural Resources/ICMBio – Chico Mendes Institute of Biodiversity Conservation, Brazil. Portuguese.
- IBAMA – Instituto Brasileiro do Meio Ambiente e dos Recursos Naturais Renováveis [Brazilian Institute for the Environment and Renewables Resources], 2005. Biota Monitoring Guidelines for Marine Seismic Surveys. Portuguese. <http://www.ibama.gov.br/phocadownload/licenciamento/petroleo-e-gas/diretrizes/2005-guia-de-monitoramento-da-biota-marinha-em-atividades-aquisicao-de-dados-sismicos.pdf>.
- IMA, 2016. Projeto de monitoramento de praias – Pesquisa sísmica marítima, 3D, nos blocos BAR-M-292, 293, 313, 314, bacia de Barreirinhas. Relatório Final. Instituto Mamíferos Aquáticos. Processo IBAMA 02022.000268/2014 (DOC29), Portuguese.
- Instituto Aqualie, 2017. Estudo sobre a movimentação e comportamento de mergulho da baleia-jubarte (*Megaptera novaeangliae*) em área de interesse de prospecção sísmica na Bacia de Santos. Portuguese.
- IWC SC, 2012. International Whaling Commission – Report of the Scientific Committee. Panama City, Panama.
- IWC SC, 2013. International Whaling Commission – Report of the Scientific Committee. Jeju, Korea.
- IWC SC, 2014. International Whaling Commission – Report of the Scientific Committee. Bled, Slovenia.
- IWC SC, 2015. International Whaling Commission – Report of the Scientific Committee. San Diego, USA.
- IWC SC, 2016. International Whaling Commission – Report of the Scientific Committee. Bled, Slovenia.
- IWC SC, 2017. International Whaling Commission – Report of the Scientific Committee. Bled, Slovenia.
- Jasny, M., Reynolds, J., Horowitz, C., Wetzler, A., 2005. The Rising Toll of Sonar, Shipping and Industrial Ocean Noise on Marine Life. <https://www.nrdc.org/sites/default/files/sound.pdf>.
- Lokkeborg, S., Ona, E., Vold, A., Salthaug, A., Jech, J.M., 2012. Sounds from seismic air guns: gear- and species-specific effects on catch rates and fish distribution. Can. J. Fish. Aquat. Sci. 69, 1278–1291, <http://dx.doi.org/10.1139/f2012-059>.
- Magalhães, W., Azevedo, A., Ristau, N., Borges, J.C., Sousa-Lima, R., Almeida, S., de Castilho, P.V., Andriolo, A., Bertozzi, C., Junior, J.L.B., Mayorga, L.F., Bhering, R.C., Kolesnikovas, C., Moreira, S.C., Cremer, M., Barreto, A., Reis, L., Engel, M., Alencar, A.E., Fragoso, A.B., de Meirelles, A.C., Flach, L., de L. Silva, F.J., Medeiros, P.I., 2017. Open Letter of Support to Marine Seismic Surveys Licensing. <https://asibamario.blogspot.com.br/2017/07/carta-aberta-de-instituicoes-de.html>.
- McCauley, R.D., Day, R.D., Swadlow, K.M., Fitzgibbon, Q.P., Watson, R.A., Semmens, J.M., 2017. Widely used marine seismic survey air gun operations negatively impact zooplankton. Nat. Ecol. Evol. 1, 195, <http://dx.doi.org/10.1038/s41559-017-0195>.
- McCauley, R.D., Fewtrell, J., Popper, A.N., 2003. High intensity anthropogenic sound damages fish ears. J. Acoust. Soc. Am. 113, 631–642, <http://dx.doi.org/10.1121/1.1527962>.
- de Mendonça, P.M.M., Spadini, A.R., Milani, E.J., 2004. Exploração na Petrobras: 50 anos de sucesso. Bol. Geociências da Petrobras 12, 9–58.
- Miller, P.J.O., Johnson, M.P., Madsen, P.T., Biassoni, N., Quero, M., Tyack, P.L., 2009. Using at-sea experiments to study the effects of airguns on the foraging behavior of sperm whales in the Gulf of Mexico. Deep. Res. I Oceanogr. Res. Pap. 56, 1168–1181, <http://dx.doi.org/10.1016/j.dsr.2009.02.008>.
- Moriyasu, M., Allain, R., Benhalima, K., Claytor, R., 2004. Effects of Seismic and Marine Noise on Invertebrates: A Literature Review. <http://www.dfo-mpo.gc.ca/csas-sccs/Publications/ResDocs-DocRech/2004/RES2004.126.e.pdf>.
- NAV, 2010. Projeto de Monitoramento de Praias – Pesquisa Sísmica Marítima 3D, Não-Exclusiva, na Bacia de Pernambuco-Paraíba, Blocos BM-PE-PB-783/837/839. Relatório Final. NAV Oceanografia Ambiental. Processo IBAMA 02022.000994/2009 (DOC11), Portuguese.
- Nelms, S.E., Piniak, W.E.D., Weir, C.R., Godley, B.J., 2016. Seismic surveys and marine turtles: an underestimated global threat? Biol. Conserv. 193, 49–65, <http://dx.doi.org/10.1016/j.biocon.2015.10.020>.
- Nieukirk, S.L., Mellinger, D.K., Moore, S.E., Klinck, K., Dziak, R.P., Goslin, J., 2012. Sounds from airguns and fin whales recorded in the mid-Atlantic Ocean, 1999–2009. J. Acoust. Soc. Am. 131, 1102–1112, <http://dx.doi.org/10.1121/1.3672648>.
- Nowacek, D.P., Clark, C.W., Mann, D., Miller, P.J.O., Rosenbaum, H.C., Golden, J.S., Jasny, M., Kraska, J., Southall, B.L., 2015. Marine seismic surveys and ocean noise: time for coordinated and prudent planning. Front. Ecol. Environ. 13, 378–386, <http://dx.doi.org/10.1890/130286>.
- Nowacek, D.P., Thorne, L.H., Johnston, D.W., Tyack, P.L., 2007. Responses of cetaceans to anthropogenic noise. Mamm. Rev. 37, 81–115, <http://dx.doi.org/10.1111/j.1365-2907.2007.00104.x>.
- Paxton, A.B., Taylor, J.C., Nowacek, D.P., Dale, J., Cole, E., Voss, C.M., Peterson, C.H., 2017. Seismic survey noise disrupted fish use of a temperate reef. Mar. Policy 78, 68–73, <http://dx.doi.org/10.1016/j.marpol.2016.12.017>.
- Popper, A.N., Hastings, M.C., 2009. The effects of anthropogenic sources of sound on fishes. J. Fish Biol. 75, 455–489, <http://dx.doi.org/10.1111/j.1095-8649.2009.02319.x>.
- Popper, A.N., Hawkins, A., 2016. The Effects of Noise on Aquatic Life II, Advances in Experimental Medicine and Biology, vol. 875. Springer, <http://dx.doi.org/10.1007/978-1-4939-2981-8>.
- Popper, A.N., Hawkins, A., 2012. The effect of noise on aquatic life. Adv. Exp. Med. Biol. 730, <http://dx.doi.org/10.1007/978-1-4419-7756-4>.
- Reyes Reyes, M.V., Bessega, M.A.I., Dolman, S.J., 2016. Review of legislation applied to seismic surveys to mitigate effects on marine mammals in Latin America. In: Proceedings of Meetings on Acoustics, p. 32002, <http://dx.doi.org/10.1121/2.0000285>.
- Simmonds, M., Dolman, S., Weilgart, L., 2004. Oceans of Noise 2004, Whale and Dolphin Conservation Society. <https://uk.whales.org/sites/default/files/oceans-of-noise.pdf>.
- Simmonds, M.P., Dolman, S.J., Jasny, M., Parsons, E.C.M., Weilgart, L., Wright, A.J., Leaper, R., 2014. Marine noise pollution – increasing recognition but need for more practical action. J. Ocean Technol. 9, 71–90 <http://animalstudiesrepository.org/acwp-ehlm/9/>.
- Slabbekoorn, H., Bouton, N., van Opzeeland, I., Coers, A., ten Cate, C., Popper, A.N., 2010. A noisy spring: the impact of globally rising underwater sound levels on fish. Trends Ecol. Evol. 25, 419–427, <http://dx.doi.org/10.1016/j.tree.2010.04.005>.
- Slotte, A., Hansen, K., Dalen, J., Ona, E., 2004. Acoustic mapping of pelagic fish distribution and abundance in relation to a seismic shooting area off the Norwegian west coast. Fish. Res. 67, 143–150, <http://dx.doi.org/10.1016/j.fishres.2003.09.046>.
- Soma, 2009. Projeto de Monitoramento de Praias – Pesquisa sísmica 3D no bloco BM-BAR-4, Bacia de Barreirinhas, Maranhão. Relatório Final. Soma Desenvolvimento e Meio Ambiente. Processo IBAMA 02022.000414/2006 (DOC35), Portuguese.
- Southall, B.L., Bowles, A.E., Ellison, W.T., Finneran, J.J., Gentry, R.L., Greene, C.R., Kastak, D., Ketten, D.R., Miller, J.H., Nachtigall, P.E., Richardson, W.J., Thomas, J.A., Tyack, P.L., 2007. Marine mammal noise exposure criteria: initial scientific recommendations. Aquat. Mamm. 33, 411–414, <http://dx.doi.org/10.1578/AM.33.4.2007.411>.
- Stone, C.J., Tasker, M.L., 2006. The effects of seismic airguns on cetaceans in UK waters. J. Cetacean Res. Manage. 8, 255–263.
- Streever, B., Raborn, S.W., Kim, K.H., Hawkins, A.D., Popper, A.N., 2016. Changes in fish catch rates in the presence of air gun sounds in Prudhoe Bay, Alaska. Arctic 69, 346, <http://dx.doi.org/10.14430/arctic4596>.
- Thalassa, Ecology Brasil, 2009. Projeto de avaliação do impacto agudo da atividade sísmica na comunidade zooplancônica. Relatório Final. Thalassa Pesquisa e Consultoria Ambiental. Processo IBAMA 02022.001743/2008, Portuguese.
- UESC, Instituto Aqualie, 2012. Monitoramento aéreo de cetáceos na Bacia de Santos. Relatório Final. Universidade Estadual de Santa Cruz, Ilhéus/BA. Processo IBAMA 02022.000342/2011 (DOC10), Portuguese.
- UFES, Petrobras, 2012. Projeto de avaliação do impacto da pesquisa sísmica marítima no comportamento de peixes. Relatório Final. Universidade Federal do Espírito Santo. Processo IBAMA 02022.003524/2008 (DOC26), Portuguese.
- UFRJ, UFES, Ekman, 2015. Projeto de pesca científica. Relatório Final. Universidade Federal do Rio de Janeiro. Universidade Federal do Espírito Santo. Processo IBAMA 02022.001382/2013 (DOC13), Portuguese.
- Univali, 2017. SIMMAM [website]. Mar. Mamm. Monit. Support Syst. <http://simmam.acad.univali.br/webgis/> (accessed 17.10.17).
- Vilaro, C., 2007. Avaliação Ambiental de Pesquisas Sísmicas Marítimas no Brasil: Evolução e Perspectivas. UFRJ – Federal University of Rio de Janeiro. Masters Dissertation, Portuguese <http://www.ppe.ufrj.br/ppe/production/tesis/mguimaraescvn.pdf>.
- Wade, L., Whitehead, H., Weilgart, L., 2010. Conflict of interest in research on anthropogenic noise and marine mammals: does funding bias conclusions? Mar. Policy 34, 320–327, <http://dx.doi.org/10.1016/j.marpol.2009.08.009>.

- Wardle, C., Carter, T., Urquhart, G., Johnstone, A.D., Ziolkowski, A., Hampson, G., Mackie, D., 2001. Effects of seismic air guns on marine fish. *Cont. Shelf Res.* 21, 1005–1027, [http://dx.doi.org/10.1016/S0278-4343\(00\)00122-9](http://dx.doi.org/10.1016/S0278-4343(00)00122-9).
- Weilgart, L., 2007. The impacts of anthropogenic ocean noise on cetaceans and implications for management. *Can. J. Zool.* 85, 1091–1116, <http://dx.doi.org/10.1139/Z07-101>.
- Weir, C.R., Dolman, S.J., 2007. Comparative review of the regional marine mammal mitigation guidelines implemented during industrial seismic surveys, and guidance towards a worldwide standard. *J. Int. Wildl. Law Policy* 10, 1–27, <http://dx.doi.org/10.1080/13880290701229838>.
- Williams, R., Wright, A.J., Ashe, E., Blight, L.K., Bruintjes, R., Canessa, R., Clark, C.W., Cullis-Suzuki, S., Dakin, D.T., Erbe, C., Hammond, P.S., Merchant, N.D., O'Hara, P.D., Purser, J., Radford, A.N., Simpson, S.D., Thomas, L., Vale, M.A., 2015. Impacts of anthropogenic noise on marine life: publication patterns, new discoveries, and future directions in research and management. *Ocean Coast. Manage.* 115, 17–24, <http://dx.doi.org/10.1016/j.ocecoaman.2015.05.021>.