

Agreement Between REGICOR and SCORE Scales in Identifying High Cardiovascular Risk in the Spanish Population

Vicente Gil-Guillén,^a Domingo Orozco-Beltrán,^b Antonio Maiques-Galán,^b José Aznar-Vicente,^c Jorge Navarro,^b Luis Cea-Calvo,^d Fernando Quirze-Andrés,^b Josep Redón,^e and Jaime Merino-Sánchez^f

^aREDIAPP-CV, Unidad de Investigación, Docencia y Práctica Clínica, Departamento 18 CV, Universidad Miguel Hernández, Elche, Alicante, Spain

^bREDIAPP-CV, Societat Valenciana de Medicina Familiar i Comunitària, Valencia, Spain

^cUnidad de Investigación, Hospital Marina Alta, Denia, Alicante, Spain

^dEspecialista en Medicina Interna, Madrid, Spain

^eServicio de Medicina Interna, Hospital Clínico, Universidad de Valencia, Valencia, Spain

^fUniversidad Miguel Hernández, Elche, Alicante, Spain

Introduction. The aims of this study were to evaluate the consistency between the SCORE (Systematic Coronary Risk Evaluation) and REGICOR (Registre Gironí del Cor) scales in identifying high cardiovascular risk and to describe the characteristics of those individuals for whom scale results were discrepant.

Methods. This cross-sectional study involved 8942 subjects aged 40-65 years who had an indication for a complete lipid profile. The agreement between SCORE (for low-risk countries) and Framingham-REGICOR (with a high risk threshold of 10%) scales in classifying patients as high risk was evaluated using the kappa statistic. Subjects for whom there was a discrepancy between classifications were identified and variables associated with this discrepancy were determined by multivariate analysis involving binary logistic regression.

Results. The REGICOR scale classified 6.7% of subjects (95% confidence interval [CI], 6.2-7.3) as high-risk, while SCORE classified 12.5% (95% CI 11.8-13.2) as high-risk. Discrepant findings were observed in 10.2% of the total population (8% had a high risk on SCORE but not REGICOR, and 2.2% had a high risk on REGICOR but not SCORE; $\kappa=0.420$; $P<.001$). The best agreement was observed between SCORE and REGICOR with a high-risk threshold of 8% ($\kappa=0.463$). Multivariate analysis showed that a high risk on SCORE but not REGICOR was associated with lower age, female sex, a high fasting glucose level, and raised diastolic blood pressure, and a high risk on REGICOR but not SCORE, with male sex,

smoking, and a low high-density lipoprotein (HDL) cholesterol level. These variables accounted for the extent of the discrepancy in 93.2% of cases.

Conclusions. The SCORE and REGICOR (threshold 10%) scales identified different populations as being at a high risk, though the agreement between them was reasonably good. The concurrence of a number of factors (eg, male sex, low HDL-cholesterol, and smoking) in a subject with a low risk on the SCORE scale should be regarded as increasing the cardiovascular risk.

Key words: Cardiovascular risk. Risk factors. SCORE. REGICOR.

Concordancia de las escalas REGICOR y SCORE para la identificación del riesgo cardiovascular alto en la población española

Introducción y objetivos. Los objetivos de este estudio fueron valorar la concordancia entre las escalas SCORE y REGICOR para la identificación de riesgo cardiovascular alto, y describir los perfiles en los que las escalas discrepan.

Métodos. Estudio transversal en 8.942 sujetos de 40-65 años con indicación de perfil lipídico completo. Se valoró la concordancia en la clasificación de riesgo alto entre las escalas SCORE (para países de bajo riesgo) y Framingham-REGICOR (umbral de riesgo alto, 10%) mediante el índice kappa. Se identificó a los sujetos con discrepancia en la clasificación y se realizó un análisis multivariable por regresión logística binaria para identificar las variables relacionadas.

Resultados. REGICOR clasificó como alto riesgo al 6,7% (intervalo de confianza del 95%, 6,2-7,3) y SCORE al 12,5% (11,8%-13,2%). El 10,2% mostró discrepancias (el 8% riesgo SCORE alto y REGICOR no alto, y el 2,2% REGICOR alto y SCORE no alto; $\kappa = 0,420$; $p < 0,001$). La concordancia más elevada fue entre SCORE y REGICOR con umbral de alto riesgo del 8% ($\kappa = 0,463$). En un análisis multivariable, SCORE alto con REGICOR no alto se relacionó con edad inferior, sexo femenino,

Correspondence: Dr. V.F. Gil Guillén.
Departamento de Medicina Clínica. Facultad de Medicina. Universidad Miguel Hernández. Campus Universitario de San Juan.
Ctra. de Valencia N-332, km 87. 03550 San Juan. Alicante. España.
E-mail: info@siccro.com

Received March 6, 2007.
Accepted for publication June 29, 2007.

glucosa basal elevada y presión arterial diastólica elevada, y SCORE no alto con REGICOR alto, con sexo masculino, tabaquismo y colesterol de las lipoproteínas de alta densidad bajo. Estas variables explicaron la variabilidad en las discrepancias en un 93,2%.

Conclusiones. SCORE y REGICOR (umbral, 10%) identificaron poblaciones de riesgo alto diferentes, y la concordancia fue discreta. Se podría considerar que la confluencia de algunas variables (sexo varón, colesterol de las lipoproteínas de alta densidad bajo, tabaquismo) y riesgo SCORE no alto incrementa el riesgo cardiovascular.

Palabras clave: *Riesgo cardiovascular. Factores de riesgo. SCORE. REGICOR.*

ABBREVIATIONS

BP: blood pressure

CEIPC: *Comité Español Interdisciplinario para la Prevención Cardiovascular* (Interdisciplinary Spanish Committee for Cardiovascular Prevention)

DM: diabetes mellitus

HBP: high blood pressure

HDL-C: high density lipoprotein cholesterol

LDL-C: low density lipoprotein cholesterol

PAPPS: *Programa de Actividades Preventivas y Promoción de la Salud de la Sociedad Española de Medicina Familiar y Comunitaria* (Health Promotion and Preventive Activities Program of the Spanish Society for Family and Community Medicine)

INTRODUCTION

Stratifying cardiovascular risk using risk charts is central to decision-making on treatment to prevent cardiovascular disease. The Framingham charts, which are used to calculate coronary morbidity and mortality,¹ and the SCORE risk charts, which are used to calculate cardiovascular mortality,² both at 10 years, are the most widely used in clinical practice. Several analyses have nonetheless concluded that the Framingham-Wilson equation overestimates coronary risk in southern European countries, where there is a lower incidence of acute myocardial infarction.³⁻⁸ Various scientific bodies have therefore proposed carrying out local population cohort studies to obtain risk charts which are applicable to given areas or which allow existing charts to be adapted for use in those areas.^{4,9,10}

In Spain, the Framingham-Wilson score¹ was adapted for use in the Girona region using a validated

methodology⁹ and led to the production of the REGICOR scale.^{11,12}

Cross-sectional studies were also performed to obtain the DORICA (Dislipidemia, Obesity, and Cardiovascular Risk) charts.¹³ On the other hand, following publication of the results of the SCORE project, in which 3 Spanish cohorts were included,² several societies (Third Report of the European Atherosclerosis Societies,¹⁴ Spanish Interdisciplinary Committee for Cardiovascular Prevention [CEIPC],¹⁵ and the Health Promotion and Preventive Activities Program of the Spanish Society for Family and Community Medicine [PAPPS]¹⁶) recommend using the low-risk country SCORE chart to calculate cardiovascular risk.

Discrepancies between the REGICOR and SCORE charts have been the subject of debate.¹⁷ A 10-year retrospective cohort study in a Spanish urban population concluded that the SCORE chart had greater validity than REGICOR for assessing the risk of coronary episodes and cardiovascular mortality, although neither scale accurately reflected the reality of the situation as it was suggested that REGICOR underestimated, and SCORE overestimated the degree of risk.¹⁸ Based on the results of that study, it was proposed that the high risk cut-point on REGICOR should be reduced from 20% to 10%. Using the 20% threshold, REGICOR classifies very few patients as high risk. If the 10% cut-point is used, the number of high risk cases is similar to that produced using the SCORE chart. Agreement with the Framingham-Wilson equation also improves.^{19,20}

Agreement between SCORE (using the CEIPC recommendations) and REGICOR has not been analyzed in populations which are free of cardiovascular disease and there has been no analysis of the sub-groups of patients in which discrepancies are observed between the 2 charts. Likewise, the implications for treatment of discrepancies between the 2 charts have not been studied. The objectives of the present study were to assess the agreement between the 2 charts and to determine the clinical profiles of subjects in whom the 2 charts differed with regard to classifying them as high risk cases.

METHODS

Sample Selection

Subjects were selected from a preventive activities program run by the Ministry of Health of the Valencian Community in 2003 in collaboration with the Valencian Society for Community and Family Medicine (SVMFiC), the Health Care Research Network for Preventive Activities in the Valencia Region (REDIAPP-cardiovascular) and the Community and Family Medicine Unit in the Department of Clinical Medicine at the Miguel Hernández de Elche University (Alicante). As recommended by PAPPS, the program was aimed at the adult population aged 40 years or over.¹⁵ Individuals were

invited to participate by letter and those wishing to be involved were given an appointment at their local health center for an evaluation by medical and nursing staff. The results were entered in a data-base and the final assessment was carried out by the Research, Teaching, and Clinical Practice Unit in Department 18 of the Valencian Community and the Department of Family Medicine at the Miguel Hernández University.

For the present study, subjects who had been clinically examined within the first 6 months of the project were included consecutively if: they had the data necessary to calculate a SCORE (using total cholesterol) and REGICOR classification; if they had no history of cardiovascular disease, and; if they were within the age range in which the 2 charts can be applied. Therefore, for this analysis, individuals were included who: *a*) were aged 40-65 years; *b*) had no established history of cardiovascular disease; and *c*) had a complete lipid profile performed according to CEIPC recommendations. Subjects with cardiovascular disease, those outside the required age range, or without a complete lipid profile or any of the other variables needed to calculate risk were excluded from the analysis.^{2,11}

Variables Analyzed

To calculate coronary risk with REGICOR, the following variables were collected: age, sex, total cholesterol, high density lipoprotein cholesterol (HDL-C), systolic blood pressure (SBP), diastolic blood pressure (DBP), smoking, and history of diabetes mellitus (DM).¹¹ Variables used to calculate risk of cardiovascular death using SCORE (based on total cholesterol figures and using the version for low-risk countries) were age, sex, total cholesterol, SBP, and smoking history.

Blood pressure was measured using mercury sphygmomanometers as recommended in several guidelines.^{21,22} Blood tests were performed in venous blood, after at least 8 h fasting, in reference laboratories in the Valencian community. Lipid profiles were evaluated in accordance with the recommendations of the Adult Treatment Panel III (ATP-III).²³

The low-risk SCORE chart and REGICOR were compared using a 10% high risk cut-point for REGICOR based on the good agreement with the Framingham-Wilson equation using this threshold.^{19,20} A threshold of 5% was used to define high risk on the SCORE chart.² Based on the CEIPC criteria,¹⁵ subjects with diabetes or with blood pressure $\geq 180/110$ mm Hg, total cholesterol ≥ 320 mg/dL, or low density lipoprotein cholesterol (LDL-C) ≥ 240 mg/dL were also considered high risk using the SCORE chart.

Statistical Analysis

Agreement between the 2 charts was assessed using the kappa statistic. Discrepancies were assessed using

McNemar's χ^2 test for paired data, with 1 degree of freedom. The area under the ROC curve was used to assess the diagnostic precision for SCORE using different high risk cut-points on REGICOR. Results were interpreted using Swet's criteria.²⁴

Discrepancies between the 2 scales were analyzed using the Pearson χ^2 test for categorical variables and the Student *t* test for quantitative variables, based on estimates of normality and equality of variances. A multivariate analysis using stepwise binary logistic regression was applied in the population in which discrepancies occurred. The dependent variable consisted of the discrepancies between SCORE and REGICOR and independent variables were those used in the bivariate analysis. Prior history of DM was eliminated from the model as a variable because, according to CEIPC,¹⁵ no individual patients with diabetes were classified as high risk by REGICOR but not by SCORE, so the model would not be able to identify agreement due to the lack of a reference category. Baseline glucose values were included instead. A *P* value less than .05 was considered statistically significant and 95% confidence intervals (CI) were calculated.

RESULTS

Of the 33 440 subjects included, 8942 (26.7%) met inclusion criteria (mean age [SE] was 51.3 [7.3] years; 59.9% were men; 27.7% were smokers; 14.4% had a history of high blood pressure [HBP]; 3.6% were diabetic, and 11.5% had a diagnosis of dyslipidemia). Mean blood pressure was 127.3 (17.1)/78.2 (10.9) mm Hg. Mean SCORE value was 1.54% (95% CI, 1.50-1.57), and the mean REGICOR value was 4.35% (4.28-4.43). The atherogenic index was 4.02 (3.99-4.04). Table 1 shows sample characteristics by gender.

Using a threshold of 20%, 12.5% of the sample was classified as high risk using the SCORE chart compared to 0.5% using REGICOR. When a cut-point of 10% was used, the proportion of high risk subjects on the SCORE chart fell to 6.7%. The proportion of high risk subjects on the REGICOR chart is shown in Figure 1 for different cut-points. Kappa indices between the REGICOR and the low-risk country SCORE chart are also shown. The kappa index increased as the cut-point for high risk decreased on the REGICOR chart. The highest agreement was observed for a threshold of 8%. At this cut-point, the number of high risk cases identified by the 2 charts was similar. Lowering the cut-point to 7% did not increase agreement between the 2 charts.

Table 2 shows the distribution of the population into high risk and non-high risk groups using thresholds of 10% on REGICOR and 5% on SCORE. There was disagreement between the 2 charts for 10.2% (9.6%-10.8%) of cases (high risk on one chart but not on the other; $\kappa=0.42$ [0.39-0.45]). Of those, 8% were classified as high risk by SCORE but not by REGICOR, and 2.2%




Figure 1. Percentage of subjects classified as high risk by SCORE and REGICOR using different cut-points; agreement with SCORE. κ indicates kappa index.

TABLE 1. Sample Characteristics by Gender^a

	General (n=8942)	Men (n=5356)	Women (n= 3586)	P ^b
Age, mean (SD), y	51.3 (7.3)	51.1 (7.4)	51.4 (7.2)	.133
BMI	27.7 (4.7)	28.1 (3.8)	27.4 (4.7)	<.001
Smoking	27.7%	36.8%	21.5%	.001
History of diabetes mellitus	3.6%	4.4%	3.1%	.001
History of high blood pressure	14.4%	14.3%	14.5%	.745
History of dyslipidemia	11.5%	12.2%	11.0%	.08
Atherogenic index	4.02 (1.29)	4.55 (1.38)	3.66 (1.09)	<.001
Glucose, mg/dL	97.3 (24.3)	101.4 (28.1)	94.5 (20.9)	<.001
Total cholesterol, mg/dL	223.3 (39.6)	226.1 (41.3)	221.4 (38.2)	<.001
HDL-C, mg/dL	59.5 (16.9)	52.8 (15.3)	63.9 (16.5)	<.001
LDL-C, mg/dL	139.2 (36.3)	143.5 (36.1)	136.3 (36.1)	<.001
Triglycerides, mg/dL	124.32 (81.4)	151.7 (97.3)	105.9 (62.2)	<.001
Systolic blood pressure, mm Hg	127.3 (17.1)	130.6 (16.6)	125.1 (17.1)	<.001
Diastolic blood pressure, mm Hg	78.2 (10.9)	80.7 (10.7)	76.6 (10.7)	<.001

^aHDL-C indicates high density lipoprotein; LDL-C, low density lipoprotein; BMI, body mass index.

^bDifference between men and women.

The table shows mean (standard deviation) and percentages.

were classified as high risk by REGICOR but not by SCORE (McNemar, $P<.001$).

Table 3 shows the characteristics of the 2 groups in which there were discrepancies between the 2 instruments. Individuals classified as high risk on SCORE but not on REGICOR included a high percentage of cases with diabetes and high blood pressure, and had higher blood pressure than the group with classified as high risk on REGICOR but not on SCORE. In the latter group, there were higher percentages of men and smokers, subjects had lower HDL-C and higher LDL-C, and blood pressure was normal, high-normal, or stage 1 hypertension.

To investigate which variables were significantly associated with the discrepancy between the 2 scales, a multivariate model was constructed for discrepant subjects which included age, gender, smoking, baseline glucose, total cholesterol, HDL-C, and LDL-C, SBP, and DBP as independent variables. Variables independently associated

with discrepancies are shown in Table 4. Discrepancies in which individuals were classified as high risk on SCORE but not on REGICOR were associated with lower age, being female, and having higher levels of DBP and baseline glucose. In the reverse situation (high risk on the REGICOR but not on SCORE), statistically significant variables were being male, smoking, and having low

TABLE 2. Distribution of Subjects According to Classification as High Risk Using SCORE and REGICOR^a

REGICOR Assessment	SCORE Assessment	
	Not High Risk	High Risk
Not high risk	7629 (85.3)	711 (8)
High risk	198 (2.2)	404 (4.5)

^aThe table shows n (%) for the total population analyzed. Kappa=0.42 (0.39-0.45); McNemar, $P<.001$; discrepancies, 10.2% (9.6%-10.8%).

TABLE 3. Characteristics of Individuals With a Discrepancy in Cardiovascular Risk Classification Using SCORE and REGICOR^a

	High Risk Using SCORE but Not REGICOR (n=711)	High Risk Using REGICOR but Not SCORE (n=198)	P ^b
Women	342 (48.1)	25 (12.6)	<.001
Men	369 (51.9)	173 (87.4)	
Age, y	54.2 (53.7-54.7)	55.3 (54.5-56.1)	.028
Age ≥50 years	491 (69.1)	163 (82.3)	<.001
BMI	29.7 (29.3-30.0)	29.2 (28.7-29.8)	.208
BMI ≥30	274 (42.2)	66 (36.1)	.109
Smoking	147 (20.7)	124 (62.6)	<.001
History of diabetes mellitus	191 (26.9)	0	<.001
History of high blood pressure	227 (31.9)	31 (15.7)	<.001
History of dyslipidemia	124 (17.4)	34 (17.2)	.930
Glucose, mg/dL	116.9 (113.7-120.2)	102.3 (98.1-106.5)	<.001
Glucose ≥110 mg/dL	261 (36.7)	39 (19.7)	<.001
Total cholesterol, mg/dL	238.4 (234.3-242.5)	244.7 (240.1-249.3)	.044
Total cholesterol <200 mg/dL	140 (19.7)	18 (9.1)	<.001
Total cholesterol 200-240 mg/dL	274 (38.5)	59 (29.8)	
Total cholesterol ≥240 mg/dL	297 (41.8)	121 (61.1)	
HDL cholesterol, mg/dL	59.7 (58.5-60.9)	40.9 (39.8-41.9)	<.001
HDL-C <40 mg/dL	52 (7.2)	81 (41.1)	<.001
LDL-C, mg/dL	146.8 (143.4-150.2)	158.7 (153.2-164.3)	<.001
LDL-C, <100 mg/dL	73 (11.4)	10 (5.8)	<.001
LDL-C, 100-129 mg/dL	138 (21.6)	19 (11.0)	
LDL-C, 130-159 mg/dL	220 (34.5)	54 (31.4)	
LDL-C, 160-189 mg/dL	119 (18.7)	61 (35.5)	
LDL-C, ≥190 mg/dL	88 (13.8)	28 (16.3)	
Triglycerides, mg/dL	147 (139.3-154.7)	217.3 (199.3-235.3)	<.001
SBP, mm Hg	146.9 (145.4-148.4)	136.4 (134.8-137.9)	<.001
SBP <130 mm Hg	113 (15.9)	31 (15.9)	<.001
SBP 130-139 mm Hg	101 (14.2)	60 (30.3)	
SBP 140-159 mm Hg	257 (36.1)	101 (51.0)	
SBP 160-179 mm Hg	188 (26.4)	6 (3)	
SBP ≥180 mm Hg	52 (7.3)	0	
DBP, mm Hg	91.4 (90.3-92.4)	82.8 (81.7-84.0)	<.001
DBP <80 mm Hg	118 (16.6)	43 (21.7)	<.001
DBP 80-89 mm Hg	144 (20.3)	88 (44.4)	
DBP 90-99 mm Hg	105 (14.8)	67 (33.8)	
DBP 100-109 mm Hg	304 (42.8)	0	
DBP ≥110 mm Hg	40 (5.6)	0	

^aHDL-C indicates high density lipoprotein cholesterol; LDL-C, low density lipoprotein cholesterol; BMI, body mass index; DBP, diastolic blood pressure; SBP, systolic blood pressure.

^bStudent *t* test with quantitative variables and Pearson χ^2 with categorical variables. The table shows mean (95% confidence interval) and n (%).

TABLE 4. Variables Associated With Discrepancies in Classifying Patients (High Risk on SCORE but Not on REGICOR)^a

	B	OR (95% CI)	P
Age	-0.118	0.89 (0.85-0.93)	<.001
Gender (women/men)	1.634	5.12 (2.72-9.64)	<.001
Smoking (no/yes)	2.229	9.29 (5.16-16.71)	<.001
Baseline glucose levels	0.019	1.02 (1.01-1.03)	<.001
Diastolic blood pressure	0.060	1.06 (1.04-1.09)	<.001
HDL-C	0.192	1.21 (1.17-1.25)	<.001

^aHDL-C indicates high density lipoprotein cholesterol; CI, confidence interval; OR, odds ratio.

Dependent variable: 1=(RA SCORE+RB REGICOR); 0=RB SCORE+RA REGICOR). Multivariable analysis (logistic regression). Model significant ($\chi^2=945.781$; $P<.001$). Explained variance = 93.2%.

HDL-C. The multivariate model was statistically significant ($P<.001$) and explained 93.2% of the variability in discrepancies. Smoking and gender were found to have the greatest weight in the model in terms of explaining discrepancies (Table 4).

Figure 2 shows discriminatory power calculated using the ROC curve with a SCORE threshold of 5% and different high risk cut-points for REGICOR. The area under the curve was 0.8 (0.785-0.814). The optimal cut-point for REGICOR was 4% (sensitivity, 86.7%; specificity, 63.9%). At this threshold REGICOR, the kappa index was 0.250 (0.234-0.266); high risk subjects, 42.4% (41.4%-43.4%), and discrepancies with SCORE (using a 5% threshold) occurred in 33.2% of cases (31.2%-34.2%).

DISCUSSION

Overall Interpretation of Results

In this population study in individuals aged 40-65 years with no history of cardiovascular disease, only moderate agreement in identifying cardiovascular risk was observed between the SCORE used according to CEIPC criteria (for low-risk countries and using a 5% cut-point) and REGICOR (using a cut-point of 10%). Factors associated with individuals being classified as high risk using SCORE but not using REGICOR were younger age, being female, and having a higher baseline DBP and glucose values; being classified as high risk by REGICOR but not by SCORE was associated with being male, smoking, and having low HDL-C. Compared to SCORE, REGICOR may underestimate risk in patients with diabetes or high blood pressure, particularly in younger women (young adults). On the other hand, in comparison to REGICOR, SCORE is likely to underestimate risk in non-diabetic males, smokers, or those with low HDL-C. This latter profile, which would be high risk on REGICOR but not on SCORE, could be considered a modifier profile indicating increased risk.

Limitations of the SCORE and REGICOR Charts

Using charts to estimate cardiovascular risk has its limitations. Comparative studies of the SCORE and Framingham charts in the Spanish population revealed discrepancies in the detection of high risk cases^{25,26} and in the recommendations for treatment (SCORE favored intervention in hypertensive women and Framingham in men with raised cholesterol²⁷). Additionally, SCORE only estimates the risk of cardiovascular mortality and not morbidity,² and the original algorithm only included the population aged between 40 and 65 years. Risk assessment in patients with diabetes is another weakness of the SCORE chart, as they were not included in the tables.² Whereas the CEIPC classifies all patients with diabetes as high risk,¹⁵ the authors of the SCORE chart recommend multiplying risk by 4 in women and by 2 in men.²

With regard to REGICOR, using the recommended cut-point of 20% meant that virtually no individuals were identified as being high risk (in the present study it did not reach 1%). Other studies¹⁸⁻²⁰ have indicated that reducing the cut-point to 10% would identify a similar proportion of high risk patients to those identified by SCORE and would lead to greater agreement with the Framingham chart, although the VERIFICA (Validation of the Adapted Framingham Individual Coronary Risk Equation) study showed that REGICOR had greater validity than the original Framingham equation in the population studied.²⁵




Figure 2. ROC curve for REGICOR chart when SCORE threshold is 5%.

Agreement Between SCORE and REGICOR in the Identification of High Risk Cases

The comparison between the 2 risk charts in the Spanish population was performed in selected groups of patients, but not in the adult general population. In a study carried out in individuals attending a health center, SCORE and Framingham would indicate the use of cholesterol-lowering treatment in a greater number of patients than REGICOR using a 10% or 20% cut-point.^{18,28,29} REGICOR indicated a lower risk than SCORE in studies in individuals with HBP or other cardiovascular risk factors,^{30,31} and there were considerable discrepancies between the 2 methods.

In the present study, a population aged 40-65 years with no history of cardiovascular disease was selected to compare the low-risk country SCORE chart with REGICOR using a high risk cut-point of 10%. We chose this threshold as it has been used previously in Spain and because it provided better agreement with the Framingham-Wilson equation.^{19,20} As in other studies,^{18,19,28} using a cut-point of 20% with REGICOR meant that almost no individuals were identified as high risk and agreement with SCORE was very low ($\kappa=0.066$). When the threshold was set at 10%, the percentage of subjects classified as high risk rose to 6.7% and the level of agreement improved ($\kappa=0.420$). Even then, however, SCORE still identified twice as many individuals as high risk compared to REGICOR.




Figure 3. Risk assessment algorithm based on CEIPC recommendations; identification of discrepant profiles. CEIPC indicates Interdisciplinary Spanish Committee for Cardiovascular Prevention; HDL-C, high density lipoprotein cholesterol; PAPPs, Health Promotion and Preventive Activities Program of the Spanish Society for Family and Community Medicine.

In this regard, the discriminatory power of the ROC curve was moderate. The optimal threshold for considering a case to be high risk using REGICOR (4%) does not appear acceptable for use in clinical practice given the substantial discrepancies with SCORE. We identified a cut-point of 8% as the threshold at which the percentage of high risk individuals was similar between the 2 methods and which showed the highest kappa value. This cut-point produced the greatest degree of agreement between the 2 charts and warrant further study in the future when using REGICOR to identify high risk individuals.

Profiles of Subjects Showing Discrepancies Between SCORE and REGICOR

Of the overall sample, 8% of subjects (78.2% of all discrepant cases) were classified as high risk by SCORE but not by REGICOR. Only 2.2% (21.8% of discrepant cases) were classified as high risk by REGICOR but not by SCORE. In other words, for every 5 individuals for whom there were discrepancies between the charts, approximately 4 were classified as high risk by SCORE but not by REGICOR and only 1 was classified as high risk by REGICOR but not by SCORE.

The profiles of the discrepant cases differed. Subjects classified as high risk by SCORE but not by REGICOR were adult women with higher blood pressure, DM, or altered baseline glucose values, and a lipid profile near

the upper limit except for HDL-C, which was normal. Cases classified as high risk by REGICOR but not by SCORE were less frequent, and tended to be adult men, smokers, with no history of DM, normal, high-normal, or stage 1 hypertension, and an altered lipid profile, particularly with low HDL-C.

The multivariate analysis explained 93.2% of the variability in the discrepant cases. Diabetes mellitus was excluded from the multivariate model because none of the patients with diabetes were classified as high risk by REGICOR but not by SCORE; baseline glucose levels were included as a continuous variable. The model showed that a predominantly female population with high DBP and baseline glucose values would be classified as high risk by SCORE but not by REGICOR. On the other hand, male smokers with low HDL-C would be classified as high risk by REGICOR but not by SCORE.

The identification of patients who are classified differently by the 2 methods (men, smokers with low HDL-C classified as high risk by REGICOR but not by SCORE) suggests that this profile may indicate increased risk. In these cases, the true risk may be higher than that estimated using the SCORE chart and treatment could be based on the algorithm shown in Figure 3. As shown in the Figure, if an individual was identified as high risk using the SCORE chart that would indicate a need for action based on the protocols described. Individuals who are not classified as high risk by SCORE but who have

the type of adverse profile described above (males, smokers, low HDL-C) should be considered as being at increased risk of cardiovascular disease and their management should be individualized.

Limitations

The study's main limitation was its cross-sectional design which meant that patients were not followed-up over time. Nevertheless, such a design is appropriate for assessing agreement between the charts and the study was performed in a large sample of the general population aged between 40 and 65 years using a rigorous methodology applied in conditions of usual clinical practice. The sample analyzed could be reasonably representative of the population in the region studied, ie, the Valencian Community, and of the age range studied, although we have no information regarding the characteristics of patients who did not accept the invitation to participate. There may also be variations between autonomous communities. The proposed modifications may help to detect individuals whose true level of cardiovascular risk is higher than that indicated by their SCORE assessment and who may require individualized management. The impact of applying these proposed modifications requires a longitudinal, follow-up study.

CONCLUSIONS

There are discrepancies in risk assessment and the identification of high risk individuals between the SCORE chart for countries with low cardiovascular risk (using a cut-point of 5%) and the REGICOR chart (using a cut-point of 10%). The 2 charts also identify different populations as being at high risk. Defining the type of patient in which discrepancies occur between the 2 charts may help to improve the clinical assessment of cardiovascular risk in patients which the SCORE chart identifies as not being at high risk but whose true cardiovascular risk could be higher than that estimated. The importance of these findings and the impact of their application in clinical practice should be confirmed in future longitudinal studies.

REFERENCES

1. Wilson PW, D'Agostino RB, Levy D, Belenger AM, Silbershatz H, Kannel WB. Prediction of coronary heart disease using risk factor categories. *Circulation*. 1998;97:1837-47.
2. Conroy RM, Pyörälä K, Fitzgerald AP, Sans S, Menotti A, de Backer G, et al. Estimation of ten-year risk of fatal cardiovascular disease in Europe: the SCORE project. *Eur Heart J*. 2003;24:987-1003.
3. Menotti A, Lanti M, Puddu PE, Kromhout D. Coronary heart disease incidence in Northern and Southern European populations: a reanalysis of the seven countries study for an European coronary risk chart. *Heart*. 2000;84:238-44.
4. Menotti A, Puddu PE, Lanti M. Comparison of the Framingham risk function-based coronary chart risk function from an Italian population study. *Eur Heart J*. 2000;21:365-70.
5. Tomás L, Vares C, Pérez I, Puig T, Balaguer I. Factores de riesgo y morbimortalidad coronaria en una cohorte laboral mediterránea seguida durante 28 años. Estudio Manresa. *Rev Esp Cardiol*. 2001;54:1146-54.
6. Kuulasmaa K, Tunstall-Pedoe H, Dobson A, Fortmann S, Sans S, Tolonen H, et al. Estimation of contribution of changes in classic risk factors to trends in coronary-event rates across the WHO MONICA Project populations. *Lancet*. 2000;355:675-87.
7. Pérez G, Pena A, Sala J, Roset PN, Masiá R, Marrugat J, and the REGICOR investigators. Acute myocardial infarction case fatality, incidence and mortality rates in a population registry in Girona, Spain, 1990-1992. *Int J Epidemiol*. 1998;27:599-604.
8. Masiá R, Pena A, Marrugat J, Sala J, Villa JS, Pavesi M, et al. High prevalence of cardiovascular risk factors in Girona, Spain, a province with low myocardial infarction incidence. *J Epidemiol Community Health*. 1998;52:707-15.
9. D'Agostino RB, Grundy S, Sullivan LM, Wilson P. Validation of the Framingham Coronary Heart Disease Prediction Scores: results of a multiple ethnic groups investigation. *JAMA*. 2001;286: 180-7.
10. Hense HW, Schulte H, Lowel H, Assman G, Keil U. Framingham risk function overestimates risk of coronary heart disease in men and women from Germany: results of the MONICA Augsburg and the PROCAM cohorts. *Eur Heart J*. 2003;24:937-45.
11. Marrugat J, Solanas P, D'Agostino R, Sullivan L, Ordovas J, Cordón F, et al. Estimación del riesgo coronario en España mediante la ecuación de Framingham calibrada. *Rev Esp Cardiol*. 2003;56: 253-61.
12. Marrugat J, D'Agostino R, Sullivan L, Elosua R, Wilson P, Ordovas J, et al. An adaptation of the Framingham risk function to southern Europe Mediterranean areas. *J Epidemiol Community Health*. 2003;57:634-8.
13. Aranceta J, Pérez C, Foz M, Mantilla T, Serra L, Moreno B, et al. Grupo colaborativo para el estudio DORICA fase II. Tablas de evaluación del riesgo coronario adaptadas a la población española. Estudio DORICA. *Med Clin (Barc)*. 2004;123:686-91.
14. de Backer G, Ambrosioni E, Borch-Johnsen K, Brotons C, Cifkova R, Dallongeville J, et al. Executive summary. European guidelines on cardiovascular disease prevention in clinical practice. *Eur Heart J*. 2003;24:1601-10.
15. Brotons C, Royo-Bordonada MA, Álvarez-Sala L, Armario P, Artigao RR, Conthe P, et al. Adaptación española de la Guía Europea de Prevención Cardiovascular. Comité Español Interdisciplinario para la Prevención Cardiovascular (CEIPC). *Aten Primaria*. 2004;34: 427-32.
16. Villar F, Maiques A, Brotons C, Torcal J, Banegas JR, Lorenzo A, et al. Recomendaciones preventivas cardiovasculares en atención primaria. Actualización 2005 del Programa de Actividades Preventivas y de Promoción de la Salud (PAPPS). *Aten Primaria*. 2005;36 Supl 2:11-26.
17. Maiques Galán A. Valoración del riesgo cardiovascular. ¿Qué tabla utilizar? *Aten Primaria*. 2003;32:586-9.
18. Buitrago F, Cañón L, Díaz N, Cruces E, Bravo B, Pérez I. Comparación entre la tabla del SCORE y la función Framingham-REGICOR en la estimación del riesgo cardiovascular en una población urbana seguida durante 10 años. *Med Clin (Barc)*. 2006;127: 368-73.
19. Baena JM, Val JL, Salas LH, Sánchez R, Altes E, Deixes B, et al. Comparación de los modelos SCORE y REGICOR para el cálculo de alto riesgo cardiovascular en sujetos sin enfermedad cardiovascular atendidos en un centro de salud de Barcelona. *Rev Esp Salud Pública*. 2005;79:453-64.
20. Ramos R, Solanas P, Cordón F, Rohlfis I, Elosua R, Sala J, et al. Comparación de la función de Framingham original y la calibrada de REGICOR en la predicción del riesgo coronario poblacional. *Med Clin (Barc)*. 2003;121:521-6.

21. 2003 European Society of Hypertension-European Society of Cardiology guidelines for the management of arterial hypertension. *J Hypertens*. 2003;21:1011-53.
22. Chobanian AV, Bakris GL, Black HR, Cushman WC, Green LA, Izzo JL, et al. Seventh report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure. *Hypertension*. 2003;42:1206-52.
23. Executive summary of the Third report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III). *JAMA*. 2001;285:2486-97.
24. Burgueño MJ, García-Bastos JL, González-Buitrago JM. Las curvas ROC en la evaluación de las pruebas diagnósticas. *Med Clin (Barc)*. 1995;104:661-70.
25. Marrugat J, Subirana I, Comín E, Cabezas C, Vila J, Elosua R, et al. Validity of an adaptation of the Framingham cardiovascular risk function: the VERIFICA study. *J Epidemiol Community Health*. 2007;61:40-7.
26. Mostaza JM, Vicente I, Tabeada M, Laguna F, Echaniz A, García-Iglesias F, et al. La aplicación de las tablas del SCORE a varones de edad avanzada triplica el número de sujetos clasificados de alto riesgo en comparación con la función de Framingham. *Med Clin (Barc)*. 2005;124:487-90.
27. Maiques A, Antón F, Taix MF, Albert X, Martí EA, Collado A. Riesgo cardiovascular del SCORE comparado con el de Framingham. Consecuencias del cambio propuesto por las Sociedades Europeas. *Med Clin (Barc)*. 2004;123:681-5.
28. Cristóbal J, Lago F, Fuente J, González-Juanatemy JR, Vázquez-Bellés P, Vila M. Ecuación de Framingham, de Wilson y ecuación de REGICOR. Estudio comparativo. *Rev Esp Cardiol*. 2005;58:910-5.
29. Parrilla F, Segura A, Segú JL. Utilización de la ecuación de Framingham-REGICOR en un centro de atención primaria. Impacto sobre la prevención primaria de las enfermedades cardiovasculares. *Aten Primaria*. 2006;38:490-5.
30. García-Ortíz L, Gómez-Marcos MA, González-Elena LJ, Rodríguez-Sánchez E, García García Á, Parra-Sánchez J, et al. Framingham-Grundy, REGICOR y SCORE en la estimación del riesgo cardiovascular del paciente hipertenso. Concordancias y discrepancias (CICLO-RISK). *Hipertensión (Madrid)*. 2006;23:111-7.
31. García-Mora R, Félix Redondo FJ. Concordancia de dos métodos para el cálculo del riesgo cardiovascular: Framingham calibrado por REGICOR y SCORE. *Hipertensión (Madrid)*. 2005;22:306-10.