

Ten-Year Incidence of Fatal and Non-Fatal Myocardial Infarction in the Elderly Population of Madrid

Blanca Novella,^a Margarita Alonso,^b Francisco Rodríguez-Salvanés,^c Rosario Susi,^c Blanca Reviriego,^b Luisa Escalante,^a Carmen Suárez,^c and Rafael Gabriel^b

^aCentro de Salud Potosí, Área de Salud II, Agencia Laín Entralgo, Consejería de Salud, Madrid, Spain

^bUnidad de Investigación, Red RECAVA, Hospital Universitario La Paz, Madrid, Spain

^cUnidad de Hipertensión, Red RECAVA, Hospital Universitario de la Princesa, Madrid, Spain

Introduction and objectives. To determine the incidence of and mortality due to myocardial infarction and coronary heart disease in the elderly population of Madrid, Spain.

Methods. The study involved a population-based cohort of 1297 individuals aged over 64 years without cardiovascular disease who were recruited in 1995. All cases of fatal and non-fatal myocardial infarction recorded up until December 2004 were investigated and classified using WHO-MONICA (World Health Organization-Multinational MONitoring of trends and determinants in CARdiovascular disease) criteria.

Results. Men had a significantly higher cumulative incidence of ischemic events ($P < .001$) and sudden death ($P < .001$), and a non-significantly higher risk of myocardial infarction (6.30%; 95% confidence interval [CI], 4.33-8.76) than women (4.90%; 95% CI, 3.54-6.70; $P = .181$). While the risk of myocardial infarction increased with age ($P < .05$), gender differences tended to narrow. The incidence was higher in men (889/100 000 person-years) than women (610/100 000 person-years; $P < .001$) and increased with age ($P < .01$). This increase was progressive in women but not in men. The mortality rate was also higher in men (472/100 000 person-years; 95% CI, 248-697) than women (328/100 000 person-years; 95% CI, 188-469; $P < .001$), and was 6 times higher in the ≥ 85 -year-old age group than in those aged 65-74 years ($P < .001$).

Conclusions. The incidence of fatal and non-fatal myocardial infarction was very high in the elderly population of Madrid. Both incidence and mortality rates increased dramatically with age after 64 years. Rates were higher in men than women at all ages, though gender differences decreased with age.

Key words: Myocardial infarction. Incidence. Mortality. Elderly. Madrid.

Incidencia a diez años de infarto de miocardio fatal y no fatal en la población anciana de Madrid

Introducción y objetivos. Estimar las tasas de incidencia y mortalidad por infarto de miocardio (IM) y enfermedad coronaria (EC) en la población anciana de Madrid.

Métodos. En 1995 se estableció una cohorte de base poblacional formada por 1.297 individuos de edad superior a 64 años y sin enfermedades cardiovasculares. Se investigaron todos los casos de infarto de miocardio mortales y no mortales hasta diciembre de 2004 y se clasificaron según los criterios OMS-MONICA.

Resultados. Los varones presentaron incidencias acumulativas significativamente superiores de episodios isquémicos ($p < 0,001$) y de muerte súbita ($p < 0,001$), así como un riesgo de infarto de miocardio superior, aunque no significativamente (6,3%; intervalo de confianza [IC] del 95%, 4,33-8,76), a los de las mujeres (4,9%; IC del 95%, 3,54-6,7) ($p = 0,181$). Aunque el riesgo de infarto de miocardio aumentaba con la edad ($p < 0,05$), las diferencias entre los dos sexos tendían a reducirse. Las tasas de incidencia fueron mayores en los varones (889/100.000 personas-años) que en las mujeres (610/100.000 personas-años) ($p < 0,001$) y con la edad más avanzada ($p < 0,01$). Este aumento era progresivo en las mujeres, pero no en los varones. Las tasas de mortalidad fueron también más altas entre los varones (472; IC del 95%, 248-697) que entre las mujeres (328; IC del 95%, 188-469) ($p < 0,001$), con unos valores en el grupo de edad ≥ 85 años 6 veces superiores a los del grupo de edad de 65-74 años ($p < 0,001$).

Conclusiones. La incidencia de infarto de miocardio mortal y no mortal es muy alta en la población anciana de Madrid. Las tasas de incidencia y de mortalidad aumentan de manera muy notable con la edad a partir de los 64 años. Los varones presentan unas tasas más altas que las mujeres de cualquier edad, pero las diferencias entre los sexos disminuyen con la edad.

Palabras clave: Infarto de miocardio. Incidencia. Mortalidad. Población anciana. Madrid.

This Project has been supported by Fondo de Investigaciones Sanitarias (FIS): PI. 93/0012; 96/1993; and 02/0896; Ministry of Science and Education: PB94-1255-C04 and by Instituto de Salud Carlos III: Red ERICE (Reference: G03/065) and Red RECAVA (Reference:RD06/0014/0015).

Correspondence: Dr. R. Gabriel Sánchez.
Unidad de Investigación. Hospital Universitario La Paz.
P.º de la Castellana, 261. 28046. Madrid. España.
E-mail: rgabriel.hulp@salud.madrid.org

Received July 25, 2007.

Accepted for publication June 9, 2008.

ABBREVIATIONS

CHD: coronary heart disease
ECG: echocardiography
MI: myocardial infarction

INTRODUCTION

Coronary heart disease (CHD) is the main cause of death in most industrialized countries,^{1,2} and myocardial infarction (MI) the most dominant manifestation of the disease. In Spain, 11.2% of deaths in males and 9.6% of deaths in females were related to CHD in 2004.³ Although mortality from CHD has declined steadily both in men and women,^{4,5} it still remains one of the most important mortality causes and it is considered it will be a major public health problem during the decades ahead.⁶

CHD is also a very frequent cause of hospitalization in Spain. The number of hospitalized patients discharged with the diagnosis of CHD has tripled in the last 5 years.⁷ This has made it a priority of health-care policy, not only because of its social repercussions, but also because of its impact on the health-care system.⁸ Incidence studies conducted in the Spanish adult population indicate that CHD is a frequent disease, with rates of 200 and 50 new MI cases per 100 000 in men and women respectively.⁹⁻¹⁵ At present, the higher incidence can be observed in the older age group, being the primary cause of death in those older than 65 years-old (15% of total mortality).³ In Spain more than 58% of discharged patients with the diagnosis of cardiovascular disease are over 65 years and 60% of MI patients admitted to hospitals are over this age.⁷ Although rates are higher after the age of 65,^{3,7} most epidemiological studies focused on the middle-aged adult population, with scarce representation of the elderly.¹⁶ In fact, only few studies include participants older than 65 years old and very few older than 75.¹⁷ The impact of the striking demographic changes in the Spanish elderly population, with 10 million of new elderly during the last 10 years, on the incidence of CHD needs to be evaluated.

The aim of this paper is to estimate the age- and sex-specific incidence and mortality rates of MI and CHD in a cohort of Spanish elderly during 10 years of follow-up (1995-2004).

METHODS

Study Population

The EPICARDIAN project is a multicentre, population-based, epidemiological, cohort study of

cardiovascular diseases in the elderly population of 3 Spanish areas: Lista district (Madrid city); Arévalo county (Avila); and Begonte county (Lugo). Study design and methods in the 3 study areas has been previously published.^{18,19} In this article we report the results available for the Lista district. According to the official census of Madrid City, on March 31, 1991 Lista district comprised 5540 subjects older than 64 years. An age- and sex-matched random sample of residents older than 64 years was selected in 1994 (n=2116). Subjects with evidence of relevant disabilities and handicaps, life-threatening diseases (ie, end-stage cancer), or institutionalized were excluded. In 244 cases (11.5%) no answer was obtained due to registration errors, changes of residence or death prior to the study's commencement. A total of 652 subjects (30.8%) rejected or could not be found after several attempts. In total, it was possible to perform baseline evaluation (year 1995) on 1464 subjects. Of those, 167 subjects (11.4%) were excluded because they had previous history of clinical cardiovascular disease (prevalent cases). The prevalence of possible MI was 4%, possible stroke 3.5%, and peripheral vascular disease 1.5%. The final sample was composed by 1297 subjects free of clinical cardiovascular diseases. The same clinical examinations were repeated in 1998 and 2004. Follow-up information (medical information and vital status) was updated every year. All participants were followed until the occurrence of the first MI, until death or until December 31, 2004 if alive (Figure 1).

Field Study Procedures

The fieldwork was organized into 2 phases. First phase included a home interview. Registered nurses were trained and certified in performing the interview and the examination. Demographic and social data, personal and family history of major cardiovascular diseases, lifestyle information, and regular drug consumption were obtained using questionnaires adapted and validated in Spanish by the MONICA project.²⁰ The interview also included the Rose questionnaire on chest pain and dyspnea.²¹ Those subjects who rejected the home interview received a self-administered questionnaire by mail, and their relatives were contacted in order to obtain information about their general health, particularly risk factors and cardiovascular diseases. We also periodically requested information from the Madrid office of the National Institute of Statistics for those subjects unreachable or dead at the time of the interview.

The second phase pertained to those individuals who responded positively to the questions about cardiac disease or angina in the first phase. We used the methods previously used and validated in the MONICA Project,²⁰ and modified and adapted in Spain by the REGICOR and the IBERICA projects.^{22,23} These pursuit and follow-up procedures were adapted to local characteristics of Lista district. Follow-up information was updated every year through emergency services, public and private hospital and primary care




Figure 1. Flow-chart EPICARDIAN study (1994-2004). CHD indicates coronary heart disease (CHD).

centres within the area, in all members of the cohort regardless of they were localized or rejected the follow-up interviews. Phone calls to the subjects and their relatives were held for those not contacted and rejections. Information on vital status of the cohort members was annually updated with the National Institute of Statistics in Madrid. In 1998 and 2004 interviews and clinical examination were repeated using the same contact procedures and interview methods as in 1994.

Definition of Myocardial Infarction (MI)

The criterion was based on symptoms, enzyme changes, electrocardiographic (ECG) changes (Minnesota code),^{24,25} and post-mortem findings defined in the WHO coding criteria.²⁶ The validity of the diagnosis was ensured by intra-observer studies.²⁷

We considered definite MI in those subjects who presented a definite ECG or typical/atypical symptoms, together with a probable ECG and abnormal enzymes, or those with typical symptoms, abnormal enzymes, and ischemic ECG.

Possible MI or coronary death was considered in living subjects with typical symptoms where enzymes and ECG did not allow classification into the definite category, with no evidence of other diagnosis; or death patients with no clear evidence of other cause of deceased.

No MI was established in living patients when the symptoms and diagnostic tests combinations did not allow classification into the possible MI or when the event could be explained by means of other causes. In fatal events, MI was rejected when death was related to different clinical diagnosis or autopsy.

TABLE 1. Demographic Characteristics and Distribution of Major Cardiovascular Risk Factors of the Cohort at Baseline (Year 1995)

	Men (n=509)	Women (n=788)	Both (n=1297)
Age group			
65-74	273 (21)	430 (33.2)	703 (54.2)
75-84	174 (13.4)	260 (20.0)	434 (33.5)
>85	62 (4.8)	98 (7.6)	160 (12.3)
Educational level			
Illiteracy	3 (0.7)	15 (2.2)	18 (1.6)
Read and write	140 (32.4)	313 (46)	453 (40.7)
Primary studies	78 (18.1)	198 (29.1)	276 (24.8)
Secondary studies	211 (48.8)	155 (22.8)	366 (32.9)
Marital status			
Single	45 (10.4)	141 (20.7)	186 (16.7)
Married	338 (78.4)	213 (31.3)	551 (49.6)
Divorced	10 (2.3)	28 (4.1)	38 (3.4)
Widowed	38 (8.8)	298 (43.8)	336 (30.2)
Cardiovascular risk factors			
Hypertension	273 (21)	464 (35.8)	737 (56.8)
Hypercholesterolemia	86 (6.6)	259 (20)	345 (26.6)
Diabetes	41 (3.2)	46 (3.5)	87 (6.7)
Obesity	345 (26.5)	431 (33.2)	776 (59.7)
Smoking	111 (8.6)	69 (5.3)	180 (13.9)

Number of cases and their corresponding percentages (%).

Diabetes indicates fasting plasma glucose ≥ 126 mg/dL or antidiabetic treatment; hypercholesterolemia, total cholesterol ≥ 250 mg/dL or lipid lowering treatment; hypertension, SBP ≥ 140 mm Hg or DBP ≥ 90 mm Hg or antihypertensive treatment; obesity, body mass index (BMI) ≥ 30 kg/m²; smoking, 1 cigarette per day or 5 per week during the last year

Statistical Methods

Basic statistics of centralization (mean, median and its standard deviation, and distribution values) were used to describe continuous variables. For the calculation of rates and proportions, the distribution of relative frequencies with confidence intervals at 95% by the Poisson approximated method was used.²⁸

To compare proportions the Pearson χ^2 test or the χ^2 for lineal trend for several categories were used. A *P* value less than .05 was considered significant. The Kaplan-Meier²⁹ actuarial method was used in order to estimate the cohort survival and crude incidence rates. Crude incidence rates were standardized by the direct method using the standard European population in 2006 as reference.³⁰

Incidence rate is the number of first fatal and/or non-fatal events divided by midyear population at risk. Mortality rate is the number of fatal events divided by midyear population at risk. Case fatality is the proportion of fatal MI in all MI. Incidence and mortality rates are presented per 100 000 person-years, and case-fatality as a percentage by age and sex groups for the 10-year follow-up.

RESULTS

The general characteristics of the initial cohort (n=1297 persons) are described in Table 1. There were 509 men

(39.2%) and 788 (60.8%) women free of clinical cardiovascular disease. The majority of participants (87.7%) were aged less than 84 years at baseline. The mean age at baseline was 75 (0.39) years (75 in women and 74.9 in men). Approximately 50% of the sample was married, 30% were widowed, 17% were single, and 3% were divorced. There was 1.6% illiteracy. The 56.8% of subjects meet clinical criteria of hypertension according to the V Joint National Committee (JNC-V) definition; 26.6% had hypercholesterolemia by the National Cholesterol Education Program (NCEP) criteria, 6.7% meet the WHO criteria of diabetes, 59.7% were obese (body mass index ≥ 30 kg/m²) and 13.9% were smokers by the WHO criteria (Table 1).

The initial cohort of 1297 subjects accounted for a total 9987 persons/year of follow-up. Mean follow-up for the entire cohort was 7.7 years. During the 10-year follow-up period 434 persons (33.4%) died and 343 persons (26.7%) could not be interviewed personally. Nevertheless, we obtained cardiovascular information through secondary sources (primary-care centres, hospital reports, mail, phone calls, etc) for those subjects not contacted or who rejected the scheduled clinical interview. Death certificates were obtained through the Madrid INE office for all those individuals not localized, who rejected the interviews or died before December 31, 2004 (n=777). In total, 117 coronary events occurred in the entire cohort during the 10-year follow-up period. There were 33 cases of non-fatal definite MI; 38 of fatal definite MI; 19 of

TABLE 2. Cumulative Number of Cases and Risk by Sex and Age of Different CHD Diagnostic Categories in 10 Years of Follow-up (1995-2004). The EPICARDIAN-Lista Study

CHD Diagnostic Categories	Males			Females			Total (95% CI)	P
	All Males (95% CI)			All Females (95% CI)				
	65-74	75-84	>85	65-74	75-84	>85		
Non-fatal definite AMI								
Events	11	2	2	8	5	5	18	
%	4 (2.0-7.1)	1.1 (0.1-4.1)	3.2 (0.4-11.2)	1.9 (0.8-3.6)	1.9 (0.6-4.4)	5.1 (1.7-11.5)	2.3 (1.4-3.6)	.173
Fatal definite AMI								
Events	9	4	4	5	10	6	21	
%	3.3 (1.5-6.2)	2.3 (0.6-5.8)	6.5 (1.8-15.7)	1.2 (0.4-2.7)	3.8 (1.9-6.9)	6.1 (2.3-12.9)	2.7 (1.7-4.0)	<.05
Possible AMI								
Events	8	1	0	5	5	0	10	
%	2.9 (1.3-5.7)	0.6 (0.1-3.1)	0	1.2 (0.4-2.7)	1.9 (0.6-4.4)	0	1.3 (0.6-2.3)	.154
Sudden death								
Events	6	9	4	1	5	2	8	
%	2.2 (0.8-4.7)	5.2 (2.4-9.6)	6.5 (1.8-15.7)	0.2 (0.1-1.3)	1.9 (0.6-4.4)	2 (0.3-7.2)	1.0 (0.4-1.9)	<.01
Fatal and non-fatal definite AMI								
Events	20	6	6	13	15	11	39	
%	7.3 (4.5-11.1)	3.4 (1.3-7.4)	9.7 (3.6-19.8)	3 (1.6-5.1)	5.8 (3.3-9.3)	11.2 (5.7-19.2)	4.9 (3.5-6.7)	<.01
Ischaemic event								
Events	34	16	10	19	25	13	57	
%	12.5 (8.8-16.9)	9.2 (5.3-14.5)	16.1 (8.0-27.6)	4.4 (2.7-6.8)	9.6 (6.3-13.8)	13.3 (7.2-21.6)	7.2 (5.5-9.3)	<.001
Possible and non-fatal definite AMI								
Events	19	3	2	13	10	5	28	
%	7 (4.2-10.7)	1.7 (0.4-4.9)	3.2 (0.4-11.2)	3.0 (1.6-5.1)	3.8 (1.9-6.9)	5.1 (1.7-11.5)	3.6 (2.4-5.0)	.074

P represents differences by age and gender.
AMI indicates acute myocardial infarction.

possible MI; and 27 of sudden death (Table 2). Men showed a higher and significant cumulative risk of ischemic events ($P<.001$) and sudden death ($P<.001$) and a higher, but not significant, risk of definite MI ($P=.181$). Risk of definitive acute myocardial infarction reaches significance when analyzed by age ($P<.01$). The analysis by age also showed that differences by gender decrease as men and women get older, except for the case of sudden death.

Incidence rates (first event) by sex and age of different acute myocardial infarction diagnostic categories are shown in Figure 2. Males older than 85 years showed the highest incidence rates ($P<.001$). Incidence of first definite (fatal and non fatal) MI was 710.9 (95% CI, 555.6-895.9) per 100000 person-years, greater in men (889.3; 95% CI, 609.1-1253.2) than in women (610.4; 95% CI, 434.4-833.5) ($P<.001$). Age-adjusted incidence rate for MI was 883 per 100 000 person-years in men and 696 per 100 000 person-years in women.

Four hundred and thirty four out of 1297 subjects died of any cause during the 10-year follow-up period, which represents a total mortality of 33.5% (95% CI, 30.9-36.1). Mortality was higher in men 42.2% (95% CI, 37.9-46.6) than in women 27.8% (95% CI, 24.7-31.1) ($P<.001$) and higher in older groups than in younger groups in both sexes ($P<.001$). The specific mortality for CHD, MI, or sudden death, was 5% (95% CI, 3.8-6.3), higher in men 7.1% (95% CI, 5.0-9.6) than in women 3.7% (95% CI, 2.5-5.2) ($P<.01$) and increased significantly with age ($P<.001$). Finally, MI as a definite cause of death was confirmed only in 38 subjects, which resulted in a specific MI mortality of 2.9% (95% CI, 2.1-3.9). MI mortality was higher, but not significantly, in men than in women ($P=.482$) and among the older groups ($P<.01$). Case fatality increase with age and was higher




Figure 2. Rates by sex and age of different acute myocardial infarction diagnostic categories. A: men. B: women.

in men than in women of all age groups, except in the 75-84 age-group, but it did not reach statistical significance (Table 3).

CHD mortality rate was 650 cases per 100 000 person-years with higher figures in men (1000 per 100 000 person-years) than in women (453 per 100 000 person-years) ($P < .001$). MI mortality rate was 380 (95% CI, 269-521) cases per 100 000 person-years, greater in men (472; 95% CI, 248-697) than in women (328; 95% CI, 188-469) ($P < .001$). Age-adjusted mortality rate for MI was 470 per 100 000 person-years in men and 386 per 100 000 person-years in women. MI mortality rate in the population aged ≥ 85 years old reached the highest value with 1241 cases per 100 000 person-years, almost 3 times more than in 75-84 age-group (445 per 100 000 person-

years), and 6 times higher than in 65-74 years old (232 per 100 000 person-years) ($P < .001$).

DISCUSSION

In the present study we show that in subjects older than 65 years, the incidence and mortality rates for MI are much higher than in this middle-age adult population (<65 years old).^{31,32} The cohort includes a significant proportion of subjects older than 80 years (22.4%), an aspect of interest, given the scarce data on this subgroup of age.

During the last few years, several epidemiological studies have been initiated in Spain to estimate the MI incidence and mortality. However, they^{9,22,23} were limited

TABLE 3. Case-Fatality for Myocardial Infarction by Sex and Age Among the Study Participants. EPICARDIAN-Lista Study (1995-2004)

Gender	Age, y	Case Fatality, % (Fatal MI/All MI)
Men	65-74	32.1 (9/28)
	75-84	40 (4/10)
	>85	67 (4/6)
Women	65-74	28 (5/18)
	75-84	50 (10/20)
	>85	54 (21/49)
Total	≥65	42 (38/90)

MI indicates myocardial infarction.

Values represent proportion with fatal events within each group.

to middle-age subjects and included few participants from the older age groups. A recent registry, the REGICOR-65 plus, provides information on MI incidence, mortality rates, and case-fatality in the elderly population of Girona, but only MI cases occurring between 1996 and 1997 were registered, which may not represent exactly the real change in risk over time.³³ To our knowledge, this is the first population-based cohort study in the elderly in Spain, with individual follow-up of study participants for a long period of time.

In order to compare our results with other studies we adopted previously used and validated international epidemiological procedures and criteria for the identification and classification of MI.^{34,35} Nevertheless, almost all epidemiological studies share similar limitations when trying to identify cardiac events. One important limitation is the difficulty to identify silent ischemia. The proportion of silent ischemia reaches 30% in some special populations, being particularly high in the elderly and the postmenopausal women.³⁶⁻³⁸ We tried to minimize this potential underestimation using multiple specific strategies such as reviewing clinical reports. Such a limitation should especially be taken into account in this study, with an elderly population in which there is a large proportion of women. Other particular limitation is the possible underestimation of ischemic events, especially during the third period of follow-up (1998-2004), due to the elevated proportion of subjects not contacted (n=343; 26%). Nevertheless we used secondary sources of information in order to detect possible cases of MI in all of these subjects. It should be also emphasized that the diagnosis of MI was not only based on the ECG, but also on the clinical symptoms and the results of laboratory testing (CK-MB elevations). One might always speculate that the application of more recent diagnostic tools, such as troponin levels, might have revealed a higher proportion of MI cases, as has been estimated by other studies.³⁹

Once these potential limitations are assumed, the incidence of definite MI seems to be in accordance with other studies carried out in Spain. These studies show that incidence rates amongst the elderly are much higher than rates in middle aged-adults. This may be related to longer exposure to risk factors and to the increased extension of atherosclerosis. This finding is of crucial importance since elderly people are the fastest growing population segment in western countries and, in consequence, demand for health resources of diagnosis and treatment is bound to increase.⁴⁰ MI incidence rates were significantly higher in men than in women. The incidence increases dramatically from the age group of 65 years to ≥84 years old. This increase was progressive in women but not in men. Other studies in the elderly have observed a similar pattern, with a concentration of MI cases in men in the age group 65-74⁴¹ and over this age (after 75 years old) in women.³³ The question arises of whether males prematurely develop CHD because of a gender-related genetic predisposition or whether their occurrence is delayed in women because of a shorter exposure to relevant cardiovascular risk factors. In our study, the burden of cardiovascular risk factors is bigger in the elderly women than in the elderly men (Table 1), and this could also affect to the gender differences observed in the CHD incidence and mortality, with a relative higher increase in women.

When comparing our results to those of the REGICOR-65 plus study,³³ a registry in a Spanish elderly population, we observed higher absolute values in our study but similar age- and sex- distribution, confirming an exponential increase with age, and higher risk in men than in women. The differences in absolute values could be explained by geographic differences in MI incidence between Mediterranean and Central areas of Spain. These geographic variations have also been demonstrated in the middle-age adult population by the IBERICA study.²³ Besides this, both studies have different designs: REGICOR-65 Plus estimations are based on the registry of MI cases in the elderly of an area of Girona during 1 calendar year. In contrast, the EPICARDIAN study accounted for the total number of cases appearing during a period of 10 years. This approach consider the increase in individual risk as the population gets older. Of interest is also that differences between men and women are narrower as they become older. The same pattern has also been observed in other population-based studies.

Incidence rates in our study are also lower than other European^{36,42} and American⁴³⁻⁴⁵ cohort studies, but age- and sex- distributions are quite similar. This finding can be explained by a real higher MI incidence in north-European and American elderly populations, as has been described for a middle-aged adult population in the International MONICA study. In the Rotterdam study for instance, incidence rates were 1200 cases per

100 000 person-years for males and 700 for females.³⁶ In the US Cardiovascular Health study, incidence was 2070 cases per 100 000 person-years in men and 790 per 100 000 person-years in women. The incidence of MI was also strongly associated with age, increasing from 780 in subjects aged 65-69 to 2560 in subjects aged 85 years and older.⁴³ Other international published studies^{42,44,45} have showed similar rates. Therefore, incidence due to definite coronary events is lower in our study than in other international studies in an elderly population. Since the Spanish elderly, as middle-aged adults, show a high prevalence of cardiovascular risk factors,¹⁹ the intriguing paradox of a low MI incidence and mortality in southern Europe despite the high cardiovascular risk factor prevalence⁴⁶ is likely to hold in older ages.

In Spain, mortality information is usually obtained from official vital statistics published by the National Institute of Statistics. Previous studies^{47,48} have validated death certificate information showing a detection rate of approximately 90%. In the EPICARDIAN study, as in the IBERICA study, when MI could not be confirmed as the cause of death, the case was referred to as "sudden death." Under this epigraph were included those subjects who died within the first 24 hours from the onset of clinical symptoms, deaths without witness or when the subject was well in the previous 24 hours. In our study the proportion of sudden death was similar to the one found by Bayés de Luna et al,⁴⁹ which is amongst the lowest described in the international literature. After taking this into account, we found more similar mortality rates in men and women than those in the REGICOR-65 plus study.³³ Increased mortality with age may be related to lower effectiveness of life-saving treatments, lower hospitalization rates, lower use of diagnostic and therapeutics procedures, longer delay between onset of symptoms and admission to the emergency room and more co-morbidity in the elderly.⁵⁰⁻⁵³ Men showed a significantly higher CHD mortality compared with that of women, but gender differences are less evident when specific mortality rates (MI rates instead of CHD rates) are considered.

In summary, we conclude that incidence of fatal and non-fatal myocardial infarction is very high in the elderly population of Madrid. Incidence and mortality rates increase dramatically with age after 64 years. Men show higher rates than women at any age but, gender differences diminish with age.

ACKNOWLEDGMENTS

To Ana Isabel Ortega for writing and editing assistance; to Eva Martínez-Renedo for statistical assistance; to Javier Muñoz, and Saturio Vega members of the EPICARDIAN Study Group (Lugo and Arévalo), ERICE and RECAVA networks; to Carlos Brotons (member of ERICE and RECAVA networks), to Francisco Fernández-Avilés (coordinator of RECAVA), and to

Pedro Luis Sánchez (secretary of RECAVA) for reviewing and improving this paper with their comments.

REFERENCES

1. Uemura K, Pisa Z. Trends in cardiovascular disease mortality in industrialized countries since 1950. *World Health Statist Quart.* 1988;41:155-78.
2. Tunstall-Pedoe H, Kuulasmaa K, Amouyel P, Arveiler D, Rajakanjans AM, Pajak A. Myocardial infarction and coronary deaths in the World Health Organization MONICA Project. Registration procedures, event rates and case fatality rates in 38 populations from 21 countries in four continents. *Circulation.* 1994;90:583-612.
3. Instituto Nacional de Estadística. Defunciones según la causa de muerte 2004 [cited Jun 30 2006]. Available from: <http://www.ine.es/inebase/cgi/axi>
4. Rosamond WD, Chambless LE, Folsom AR, Cooper LS, Conwill DE, Clegg L, et al. Trends in the incidence of myocardial infarction and in mortality due to coronary heart disease, 1987 to 1994. *N Engl J Med.* 1998;339:861-7.
5. Banegas JR, Rodríguez F, Villar F, Rey J. Perspectiva epidemiológica del riesgo cardiovascular en España. *Iber J Hypertens.* 1996;1 Suppl 2:40-5.
6. Murray CL, López A. Alternative projections of mortality and disability by cause 1990-2020: Global Burden of Disease Study. *Lancet.* 1997;349:1498-504.
7. Morbilidad hospitalaria por cardiopatía isquémica por sexo. España 1977-2002. Madrid: INE; 2004.
8. Plan Integral de Cardiopatía Isquémica 2004-2007. Madrid: Ministerio de Sanidad y Consumo; 2003.
9. Tunstall-Pedoe H, Kuulasmaa K, Mähönen M, Tolonen H, Ruokokoski E, Amouyel P, for the WHO MONICA Project. Contribution of trends in survival and coronary-event rates to changes in coronary heart disease mortality: 10 year results from 37 WHO MONICA Project populations. *Lancet.* 1999;353:1547-57.
10. Marrugat J, Elosua R, Aldasoro E, Tormo MJ, Vanaclocha H, Segura A, et al. Regional variability in population acute myocardial infarction cumulative incidence and mortality rates in Spain 1997 and 1998. *Eur J Epidemiol.* 2004;19:831-9.
11. Marrugat J, Elosua R, Martí H. Epidemiología de la cardiopatía isquémica en España: estimación del número de casos y de las tendencias entre 1997 y 2005. *Rev Esp Cardiol.* 2002;55:337-46.
12. Tormo MJ, García J, Cirera LI, Contreras J, Martínez G, Rodríguez M, et al; por los investigadores y colaboradores del estudio IBERICA-Murcia. Epidemiología del infarto agudo de miocardio en la Región de Murcia: Estudio IBERICA. Serie Informes N.º 34. Murcia: Consejería de Sanidad, Dirección General de Salud Pública; 2003.
13. Tomás L, Varas C, Pérez I, Puig T, Balaguer Vintrolé I. Factores de riesgo y morbimortalidad coronaria en una cohorte laboral mediterránea seguida durante 28 años. Estudio de Manresa. *Rev Esp Cardiol.* 2001;54:1146-54.
14. Gutiérrez JA, Gómez-Jerique J, Gómez de la Cámara A, Rubio MA, García A, Arístegui I. Dieta y riesgo cardiovascular en España (DRECE II). Descripción de la evolución del perfil cardiovascular. *Med Clin (Barc).* 2000;115:726-9.
15. Medrano MJ, Boix R, Cerrato E, Ramírez M. Incidencia y prevalencia de cardiopatía isquémica y enfermedad cerebrovascular en España: revisión sistemática de la literatura. *Rev Esp Salud Pública.* 2006;80:515.

16. Gabriel Sánchez R. Hacia un estudio epidemiológico sobre factores de riesgo cardiovascular en el anciano en España. *Rev Esp Geriatr Gerontol.* 1990;25:383-8.
17. Mc Govern PG, Pankow JS, Sharar E, Doliszny KM, Folsom AR, Blackburn H, et al. Recent trends in acute coronary heart disease-mortality, morbidity, medical care, and risk factors. *N Engl J Med.* 1996;334:884-90.
18. Gabriel R, Novella B, Alonso M, Vega S, López I, Suárez C, et al. El proyecto EPICARDIAN: un estudio de cohortes sobre enfermedades y factores de riesgo cardiovascular en ancianos españoles: consideraciones metodológicas y principales hallazgos demográficos. *Rev Esp Salud Pública.* 2004;78:243-55.
19. Gabriel R, Bermejo F, Vega S, Muñoz J, López I, Suárez C, et al. Survey of cardiovascular diseases (acute myocardial infarction and stroke) and its risk factors in the elderly population of Spain: the Epicardian Study-methods and demographic findings. *Cardiovasc Dis Prev.* 1999;2:290-300.
20. World Health Organization Cardiovascular Diseases Unit: WHO MONICA Project: MONICA Manual. Geneva: World Health Organization; 1990.
21. Rose GA, Blackburn H, Gillyn RF, Prineas RJ. Métodos de encuesta sobre enfermedades cardiovasculares. 2.a ed. Ginebra: Organización Mundial de la salud; 1982.
22. Perez G, Pena A, Sala J, Roset P, Masia R, Marrugat J, and the REGICOR investigators. Acute myocardial infarction case fatality, incidence and mortality rates in a population registry in Gerona, Spain, 1990-1992. *Int J Epidemiol.* 1998;27:599-604.
23. Fiol M, Cabades A, Sala J, Marrugat J, Elosua R, Vega G, et al. Variabilidad en el manejo hospitalario del infarto agudo de miocardio en España. Estudio IBERICA (Investigación, Búsqueda Específica y Registro de Isquemia Coronaria Aguda). *Rev Esp Cardiol.* 2001;54:443-52.
24. Kennedy RD, Caird FI. The application of the Minnesota code to population studies of the electrocardiogram in the elderly. *Gerontol Clin.* 1972;14:5-16.
25. Prineas RJ, Crow RS, Blackburn H. The Minnesota code manual of electrocardiographic findings: standars and procedures for measurement and clasification. Bristol: John Wright; 1962.
26. The Joint International Society and Federation of Cardiology. World Health Organization Task Force on Stardardization of Clinical Nomenclature and Criteria for Diagnosis of Ischemia Heart Disease. *Circulation.* 1979;59:607-8.
27. Novella B. Prevalencia e incidencia de cardiopatía isquémica en ancianos españoles: estudio EPICARDIAN [tesis de Epidemiología]. Madrid: Universidad Autónoma de Madrid. Facultad de Medicina; 2002.
28. Abraira V. Intervalos de confianza de los índices de prevalencia e incidencia. Material docente de la Unidad de Bioestadística. Hospital Universitario Ramón y Cajal [cited Ene 23 2008]. Available from: http://www.hrc.es/investigacion/bioest/Medidas_frecuencia_5.html
29. Kaplan EL, Meier P. Non parametric estimation from incomplete observations. *J Am Statist Ass.* 1958;53:457-81.
30. EuroStat. Población europea 2006 [updated Jul 18 2007]. Available from: <http://ec.europa.eu/eurostat>
31. Centro Nacional de Epidemiología del Instituto de Salud Carlos III. Enfermedades cardiovasculares. Casos y tasas anuales ajustadas por edad y por sexo. España 2002 [cited Jul 23 2007]. Available from: http://www.isciii.es/htdocs/centros/epidemiologia/epi_cardioisquemica.jsp
32. Centro Nacional de Epidemiología del Instituto de Salud Carlos III. Mortalidad por causa, sexo y grupo de edad (2001) [cited Jul 23 2007]. Available from: http://www.isciii.es/htdocs/centros/epidemiologia/anexos/ww9201_ed_cau_tasa.htm
33. Marrugat J, Sala J, Manresa JM, Gil M, Elosua R, Perez G, et al, and the REGICOR. Acute myocardial infarction population incidence and in-hospital management factors associated to 28-day case-fatality in the 65 year and older. *Eur J Epidemiol.* 2004;19:231-7.
34. Tunstall H, Kuulasmaa K, Amouycl P, Arveiler D, Rajakangas A, Pajak A. Myocardial infarction and coronary death in the World Health Organization MONICA Project. Registration procedures, event rates and case-fatality rates in 38 populations from 21 countries in four continents. *Circulation.* 1994;90:583-612.
35. Prineas RJ, Crow RS, Blackburn H. The Code Manual of Electrocardiographic Findings: Standards and Procedures for Measurement and Classification. Bristol: John Wright; 1982.
36. de Torbal A, Boersma E, Kors JA, van Herpen G, Deckers JW, van der Kuip DA, et al. Incidence of recognized and unrecognized myocardial infarction in men and women aged 55 and older: the Rotterdam Study. *Eur Heart J.* 2006;27:729-36.
37. Stehbens WE. Misuse of coronary heart disease. *Heart.* 1999; 82:1-2.
38. Caso C, Ribera JM. Acute myocardial infarction in the elderly. *J Am Geriatr Soc.* 1989;37:194-5.
39. Bardají A. El papel de las troponinas en el diagnóstico y el pronóstico de los síndromes coronarios agudos. *Rev Esp Cardiol.* 2005;5:19-25.
40. Krumholz HM. Cardiopatía isquémica en el anciano. *Rev Esp Cardiol.* 2001;4:819-26.
41. Cabadés A. El registro REGICOR y la epidemiología del infarto de miocardio en España: se hace camino al andar. *Rev Esp Cardiol.* 2007;60:342-5.
42. Volmink JA, Newton JN, Hicks NR, Sleight P, Fowler GH, Neil HA. Coronary event and case fatality rates in an English population: results of the Oxford Myocardial Infarction Incidence Study Group. *Heart.* 1998;80:40-4.
43. Psaty BM, Furberg CD, Kuller LH, Bild DE, Rautaharju PM, Polar JF, et al. Traditional risk factors and subclinical disease measures as predictors of first myocardial infarction in older adults: The Cardiovascular Health Study. *Arch Intern Med.* 1999;159:1339-47.
44. Cupples LA, D'Agostino RB. Some risk factors related to the annual incidence of cardiovascular disease and death using pooled repeated biennial measurements: Framingham Heart Study, 30-year follow-up. Section 34. In: Kannel WB, Wolf PA, Garrison RJ, editors. *The Framingham Study: an epidemiological investigation of cardiovascular disease.* Bethesda: National Institutes of Health; 1987.
45. Nadelmann J, Frishman WH, Ooi WL, Tepper D, Greenberg S, Guzik H, et al. Prevalence, incidence and prognosis of recognized and unrecognized myocardial infarction in persons aged 75 years and older: The Bronx Aging Study. *Am J Cardiol.* 1990;66:533-7.
46. McGovern PG, Pankow JS, Shahar E, Doliszny KM, Folsom AR, Blackburn H, et al. Recent trends in acute CHD. *N Engl J Med.* 1996;334:884-90.
47. Pañella H, Borrell C, Rodríguez C, Roca J. Validación de la causa básica de defunción en Barcelona, 1985. *Med Clin (Barc).* 1989;92:129-34.
48. García J, Cirera L, Tormo MJ, Martínez C, Contreras J, Navarro C. Utilidad del boletín estadístico de defunción para la identificación de muertes extrahospitalarias en un registro poblacional de infartos agudos de miocardio. *Rev Esp Cardiol.* 2001;54:1041-7.
49. Bayes de Luna A, Guindo J. Sudden death in ischaemic heart disease. *Rev Port Cardiol.* 1990;9:473-9.

50. Ayanian JZ, Braunwald E. Thrombolytic therapy for patients with myocardial infarction who are older than 75 years. *Circulation*. 2000;101:2224-6.
51. Bueno H, López-Palop R, Pérez-David E. Combined effect of age and right ventricular involvement on acute inferior myocardial infarction prognosis. *Circulation*. 1998;98:1714-20.
52. Lesnefsky EJ, Lundergan CF, Hodgson J. Increased left ventricular dysfunction in elderly patients despite successful thrombolysis. The GUSTO-I angiographic experience. *Am Coll Cardiol*. 1996;28:331-7.
53. Kannel WB, Vokonas PS. Demographics of the prevalence, incidence, and management of coronary heart disease in the elderly and in women. *Ann Epidemiol*. 1992;2:5-14.