

Original article

Delayed Diagnosis of Hypertension in Diabetic Patients Monitored in Primary Care

Carmen de Burgos-Lunar,^{a,*} Isabel del Cura-González,^{b,c} Miguel A. Salinero-Fort,^d Paloma Gómez-Campelo,^d Leopoldo Pérez de Isla,^e and Rodrigo Jiménez-García^c

^aUnidad de Epidemiología Clínica e Investigación, Hospital Carlos III, Servicio Madrileño de Salud, Madrid, Spain

^bUnidad de Apoyo a la Investigación, Gerencia de Atención Primaria, Servicio Madrileño de Salud, Madrid, Spain

^cDepartamento de Medicina Preventiva y Salud Pública, Universidad Rey Juan Carlos, Madrid, Spain

^dFundación de Investigación Biomédica, Hospital Carlos III, Servicio Madrileño de Salud, Madrid, Spain

^eUnidad de Imagen Cardiovascular, Hospital Clínico San Carlos, Servicio Madrileño de Salud, Madrid, Spain

Article history:

Received 16 August 2012

Accepted 22 January 2013

Available online 16 May 2013

Keywords:

Hypertension

Diabetes mellitus

Delayed diagnosis

Primary care

ABSTRACT

Introduction and objectives: Delayed diagnosis of hypertension may result in inadequate blood pressure control and increased cardiovascular risk. The aim of this study was to estimate the delay in hypertension diagnosis in patients with type 2 diabetes and the likelihood of a diagnosis within a suitable period (first 6 months), and to analyze the patient and physician characteristics associated with delayed diagnosis.

Methods: Retrospective dynamic cohort study, with a 7-year follow-up in primary care, of 8074 adult patients with diabetes who met the diagnostic criteria for hypertension. Two thresholds were considered: 140/90 mmHg and 130/80 mmHg. The time elapsed between meeting these criteria and recording the diagnosis was estimated; the time course of the likelihood of a missed diagnosis and the variables associated with correct diagnosis were assessed by Kaplan-Meier survival analysis and logistic regression analysis, respectively.

Results: The mean diagnostic delay was 8.9 (15.4) months in patients with blood pressure \geq 140/90 mmHg compared to 15.2 (19.6) months for those with $<$ 140/90 mmHg ($P < .001$). The main variables associated with correct diagnosis were baseline blood pressure \geq 140/90 mmHg (odds ratio=2.77; 95% confidence interval, 2.44-3.15), no history of acute myocardial infarction (odds ratio=2.23; 95% confidence interval, 1.67-2.99), obesity (odds ratio=1.70; 95% confidence interval, 1.44-1.99), absence of depression (odds ratio=1.63; 95% confidence interval, 1.27-2.08), female sex (odds ratio=1.29; 95% confidence interval, 1.14-1.46), older age, and taking more intensive antidiabetic therapy. There was an inverse relationship with the age of physicians and a direct relationship with their professional stability.

Conclusions: The mean diagnostic delay in hypertension among diabetic patients was greater than 6 months and varied according to the diagnostic threshold used. Patients with baseline blood pressure \geq 140/90 mmHg were more likely to receive a timely diagnosis.

© 2013 Sociedad Española de Cardiología. Published by Elsevier España, S.L. All rights reserved.

Retraso diagnóstico de la hipertensión arterial en pacientes diabéticos atendidos en atención primaria

RESUMEN

Introducción y objetivos: El retraso diagnóstico de la hipertensión arterial puede favorecer un control deficiente y el incremento del riesgo cardiovascular. El objetivo es estimar el retraso diagnóstico de la hipertensión en los diabéticos tipo 2 y la probabilidad de que se los diagnostique en un plazo adecuado (primeros 6 meses) y analizar las características de pacientes y médicos asociadas al retraso diagnóstico.

Métodos: Cohorte dinámica retrospectiva, con 7 años de seguimiento en atención primaria, de 8.074 adultos diabéticos a los que se incluyó en el momento de cumplir criterios diagnósticos de hipertensión arterial considerando dos umbrales: 140/90 y 130/80 mmHg. Se estimó el tiempo transcurrido desde el cumplimiento de dichos criterios hasta el registro del diagnóstico la evolución temporal de la probabilidad de que no se diagnosticara mediante análisis de supervivencia de Kaplan-Meier y las variables asociadas al diagnóstico adecuado mediante regresión logística.

Resultados: El retraso diagnóstico medio fue $8,9 \pm 15,4$ meses para pacientes que acudieron con presión arterial \geq 140/90 mmHg frente a los $15,2 \pm 19,6$ meses de aquellos con presión $<$ 140/90 mmHg ($p < 0,001$). Las principales variables asociadas al diagnóstico adecuado fueron presión arterial inicial \geq 140/90 mmHg (odds ratio = 2,77; intervalo de confianza del 95%, 2,44-3,15), no tener infarto agudo de miocardio previo (odds ratio = 2,23; intervalo de confianza del 95%, 1,67-2,99), obesidad (odds ratio = 1,70; intervalo de confianza del 95%, 1,44-1,99), no sufrir depresión (odds ratio = 1,63; intervalo de confianza del 95%, 1,27-2,08), ser mujer

Palabras clave:

Hipertensión arterial

Diabetes mellitus

Retraso diagnóstico

Atención primaria

* Corresponding author: Unidad de Epidemiología Clínica, Hospital Carlos III, Sinesio Delgado 10, 28029 Madrid, Spain.

E-mail address: carmenblunar@hotmail.com (C. de Burgos-Lunar).

(odds ratio = 1,29; intervalo de confianza del 95%, 1,14-1,46), tener más edad o tratamiento antidiabético más intensivo. La edad del médico mostró relación inversa y su estabilidad laboral, relación directa.

Conclusiones: El retraso diagnóstico medio de la hipertensión en diabéticos fue > 6 meses y varió según el umbral diagnóstico utilizado. Los pacientes con presión arterial inicial $\geq 140/90$ mmHg presentaron mayor probabilidad de diagnóstico adecuado.

© 2013 Sociedad Española de Cardiología. Publicado por Elsevier España, S.L. Todos los derechos reservados.

Abbreviations

BP: blood pressure
DM2: type 2 diabetes mellitus
EMR: electronic medical record
HT: hypertension
PC: primary care

INTRODUCTION

Hypertension (HT) is a cardiovascular risk factor that affects 35% of the Spanish adult population.¹

The prevalence of HT is between 1.5 and 2.3 times higher in patients with type 2 diabetes mellitus (DM2) than in nondiabetic subjects.^{2,3} When present it hastens the course of microvascular and macrovascular complications of DM2^{4,5} and increases mortality, to the extent that 75% of deaths of cardiovascular origin in diabetic patients are attributable to HT.^{2,6}

Monitoring HT in diabetics reduces mortality and prevents or delays the incidence of vascular complications.^{7,8} However, despite the availability of effective drug therapies, blood pressure (BP) control in these patients is poor, which may be partly due to underdiagnosis and delayed diagnosis. To date no reports have estimated the typical delay in diagnosing HT. Knowledge of these factors may enable implementation of procedures that will improve the management of these patients and reduce their cardiovascular risk.

The aims of this study were to estimate the delay in diagnosing HT in adults with DM2 in primary care (PC) according to the diagnostic threshold considered and the probability of these patients receiving a diagnosis with the first 6 months of the onset of HT, as well as to analyze the association between patient and physician characteristics and the likelihood of a delayed diagnosis.

METHODS

This retrospective, analytical, observational, dynamic cohort study was performed in 21 health centers in northeast Madrid. The study population comprised all patients diagnosed with DM2 in their electronic medical record (EMR) who attended at least 2 annual check-up visits in their PC centers. Patients older than 18 years who met the diagnostic criteria for HT between January 1, 2003 and June 30, 2009 and had at least 2 BP readings recorded in the EMR during the study year were included. Patients with a HT diagnosis at the start of the study and those with a follow-up lasting less than 6 months were excluded.

Patient inclusion and follow-up began on January 1, 2003; inclusion ended on June 30, 2009, and follow-up was completed on December 31, 2009.

Data were obtained from personalized secondary data in the patients' EMR. Diagnoses of HT and DM2 recorded in the EMRs were validated in the same setting where the study was

conducted,⁹ and a positive predictive value for DM of 91.23% and a negative predictive value of 99.98% were obtained. For the HT diagnostic threshold of 140/90 mmHg, the positive and negative predictive values were 82.52% and 97.94%, and for the diagnostic threshold of 130/80 mmHg, 98.68% and 53.92%, respectively.

A patient was considered diabetic when the EMR contained a diagnosis of DM2 (International Primary Care Classification codes K86 or K87).

The patient was considered hypertensive when the measurement of 2 or more systolic blood pressure (SBP) measurements taken on at least 2 consecutive visits was ≥ 130 mmHg or the mean diastolic blood pressure (DBP), ≥ 80 mmHg, according to seventh report of Joint National Committee guidelines.¹⁰

Given the lack of agreement among scientific societies on the diagnostic thresholds for HT in patients with DM2, this study considered a second HT threshold when the mean of 2 or more SBP measurements on at least 2 consecutive visits was ≥ 140 mmHg or the mean DBP was ≥ 90 mmHg, according to the NICE (National Institute for Clinical Excellence) standard.¹¹

HT was considered diagnosed when recorded in the EMR, and as undiagnosed when the diagnostic criteria were met but no HT diagnosis was recorded in the EMR. The use of medication with a hypotensive effect but prescribed for indications other than HT was not considered.

We measured the time elapsed between the visit when the patient met the diagnostic criteria and the date when the diagnosis was recorded.

A diagnosis was defined as "correct" when it was recorded in the EMR during the first 6 months after the patient met the diagnostic criteria and as "incorrect" when it was not recorded until more than 6 months later or not at all. Patient-related variables (sociodemographic, comorbidity, anthropometric, biochemical parameters), use of health resources (consultation and treatments) and PC physician-related variables (sex, age, professional stability, work schedule and professional seniority) were analyzed.

Statistical Analysis

A descriptive analysis was prepared of the study population, overall and stratified by the BP measurement when diagnostic criteria were met. The time elapsed between meeting these criteria and recording of the diagnosis, along with the variation over time of the probability of remaining diagnosis-free, was estimated by Kaplan-Meier survival analysis. Log-rank test was used to study differences related to the degree of initial HT.

A univariate analysis was carried out of the factors associated with a correct diagnosis (ie, diagnosis of HT in the EMR during the first 6 months after the diagnostic criteria were met). The chi-square was used for qualitative variables and Student *t* test for quantitative variables. Variables with a significance < 0.25 were included in the logistic regression analysis.

All estimates were calculated with their 95% confidence intervals (95%CI). A *P* value $< .05$ was considered statistically significant. Data were analyzed using SPSS 19.0 statistical software package (SPSS Inc.; Chicago, Illinois, United States).




Figure 1. Study flowchart. BP, blood pressure; DM2, type 2 diabetes mellitus; EMR, electronic medical record; HT, hypertension.

The study was approved by the Carlos III Hospital Ethics Committee (Madrid).

RESULTS

Figure 1 shows the flow chart of study participation. Of the 27 782 patients with a diagnosis of DM2 recorded in the EMR in PC centers during the study period, 63.8% had HT at the time of the DM2 diagnosis and were excluded from the study population. Of the 10 046 patients who did not have HT when DM2 was diagnosed, 80.4% (n=8074) had BP values $\geq 130/80$ mmHg during follow-up. Of these, 79.2% (n=6395) had SBP of 130 to 139 mmHg or DBP of 80 to 89 mmHg and 20.8% (n=1679) had SBP ≥ 140 mmHg or DBP ≥ 90 mmHg.

For the lower diagnostic threshold (130-139/80-89 mmHg), 41.8% of patients were diagnosed and the diagnosis was recorded on the EMR and 54.8% remained diagnosis-free after a median follow-up of 3.4 [interquartile range, 1.6-5.2] years. For the diagnostic threshold of BP $\geq 140/90$ mmHg, 53.7% were diagnosed and 42.4% remained diagnosis-free after a median follow-up of 3.6 [1.7-5.4] years.

Table 1 summarizes the main characteristics of the study participants and their physicians, in general and stratified by the patients' baseline BP.

A missed diagnosis of HT was lower in patients with BP $\geq 140/90$ mmHg (42.4% compared to 63.1% of those with BP $\geq 130/80$ mmHg and $< 140/90$ mmHg at onset; $P < .001$),

in women (50.5% compared to 58% of men; $P < .001$), and in those aged 65 or over (51% compared to 59.2% of those under 65; $P < .001$).

Hypertensive patients ultimately diagnosed by their physician experienced a mean delay of 11.1 (17.7) months. Of these, 63.2% were diagnosed during the first 6 months; 6.1% between 6 and 12 months; 10.4% between 12 and 24 months; and for 20.2% the diagnosis was delayed more than 24 months. The mean delay in patients who had BP $\geq 140/90$ mmHg was 8.9 (15.4) months compared to 15.2 (19.6) months for those with baseline BP values $\geq 130/80$ and $< 140/90$ mmHg ($P < .001$).

Figure 2 shows the survival curve for the probability of remaining diagnosis-free over time, stratified by BP values at onset of HT. Patients with an initial BP $\geq 140/90$ mmHg were significantly more likely to receive a diagnosis (log rank test, $P < .001$).

The variables associated with correct diagnosis are shown in Table 2. Patients with initial BP $\geq 140/90$ mmHg had a probability of receiving a correct diagnosis that was 2.77 (95%CI, 2.44-3.15) times higher than patients with lower initial BP readings. Other strongly associated variables were not having a record of acute myocardial infarction in the EMR (odds ratio [OR]=2.23; 95%CI, 1.67-2.99), obesity (OR=1.70; 95%CI, 1.44-1.99), not having depression (OR=1.63; 95%CI, 1.27-2.08) and receiving treated antiplatelet agents (OR=1.47; 95%CI, 1.29-1.68). There was a nonsignificant trend toward association with microalbuminuria and macroalbuminuria. In addition, women, older patients, those with a more intensive antidiabetic treatment, and those who attended the health center most frequently were more likely to receive a correct diagnosis. Of the physician-related variables, age and professional

Table 1
Baseline Characteristics of the Sample

	Total (n=8074)	Initial BP 130-139/80-89 mmHg (n=6395)	Initial BP \geq 140/90 mmHg (n=1679)	P
Patient-related variables				
Women, %	41.4	41.4	41.3	.982
Age, years	64.6 (12.2)	64.1 (12.4)	65.6 (12.1)	<.001
<65, %	48	49.2	43.5	<.001
\geq 65, %	52	50.8	56.5	
Duration of diabetes, years	7.4 (5.8)	7.2 (5.5)	7.8 (6.5)	<.001
BMI	29.5 (4.9)	29.4 (4.8)	29.8 (5.2)	.029
<25, %	27	27	27.2	.481
25-30, %	32.7	33	31.4	
\geq 30, %	40.3	40	41.4	
Systolic blood pressure, mmHg	136.2 (12.1)	133.9 (9.9)	148 (12.1)	<.001
Diastolic blood pressure, mmHg	79.3 (8)	79 (6.2)	82 (12)	<.001
HbA _{1c}	7.3 (1.5)	7.3 (1.4)	7.5 (1.5)	<.001
LDL-C	122.6 (33.3)	123.1 (33)	124.8 (34.4)	.088
HDL-C	47.8 (12.5)	47.8 (12.3)	48.1 (13)	.408
Albuminuria	29.9 (73.7)	28.2 (71.8)	36 (80.8)	.027
Normal (<30), %	80.8	81.7	77.3	.063
Microalbuminuria (30-299), %	17.3	16.5	20.5	
Macroalbuminuria (\geq 300), %	1.9	1.8	2.2	
AMI, %	6.5	6.7	5.5	.069
ACVA, %	0.1	0.1	0.1	.252
Atrial fibrillation, %	2.4	2.6	1.9	.167
Heart failure, %	1.3	1.4	1.1	.424
Depression, %	7.8	7.8	7.9	.950
Antihypertensive treatment, %	63	59.8	75	<.001
Antiplatelet treatment, %	65.6	65.1	67.3	.118
Variables related to the use of resources				
Annual visits to the health clinic				
1-12, %	44.9	44.4	46.9	.084
>12, %	55.1	55.6	53.1	
Treatment for diabetes				
No pharmacologic treatment, %	26.3	26.4	25.9	.114
OAD, %	54.1	54.4	52.6	
Insulin (with or without OAD), %	19.6	19.2	21.5	
Physician-related variables				
Women, %	75.5	75.3	76.4	.380
Professional situation				
Permanent, %	64.1	63.6	66	.022
Temporary, %	20.8	21.5	18.3	
Substitute, %	15.1	14.9	15.7	
Work shift				
Morning, %	50.4	49.6	53.4	.018
Split (morning and afternoon), %	10.5	10.5	10.6	
Afternoon, %	39.1	39.9	36	
Age, years	46.2 (7.4)	46.2 (7.4)	46.1 (7.6)	.913
<40, %	22.9	22.6	23.8	.386
40-55, %	64.9	65.3	63.5	
\geq 55, %	12.2	12	12.7	
Professional seniority, years	15.3 (9.4)	15.2 (9.4)	15.7 (9.3)	.064
\leq 5, %	23.1	23.6	21.4	.199
5-15, %	19.4	19.3	19.6	
>15, %	57.5	57.1	59	

ACVA, acute cerebrovascular accident (excluding transient ischemic attacks); AMI, acute myocardial infarction; BMI, body mass index; BP, blood pressure; HbA_{1c}, glycohemoglobin; HDL-C, high density lipoprotein cholesterol; LDL-C, low density lipoprotein cholesterol; OAD, oral antidiabetic. Unless specified otherwise, data are expressed as mean (standard deviation).




Figure 2. Survival curve: likelihood of continuing without a diagnosis of hypertension. HT, hypertension.

stability had a significant inverse and direct relationship with correct diagnosis, respectively.

DISCUSSION

Although the relationship between BP and cardiovascular risk appears continuous up to an SBP of 115 to 110 mmHg and a DBP of 75 to 70 mmHg,^{12,13} decision-making in PC practice requires an operational definition. For the adult general population the diagnosis of HT is established as a mean BP measurement in the clinic $\geq 140/90$ mmHg.^{5,10} However, because of the increased cardiovascular risk in diabetic patients, many scientific societies have set a lower threshold for these patients of 130/80 mmHg,^{5,10,14} although there is no consensus to establish this threshold. Thus, the Spanish Ministry of Health clinical guidelines for DM2¹⁵ set the threshold at $\geq 140/80$ mmHg and NICE maintained the cut-off at $\geq 140/90$ mmHg.¹¹

The reduction in diagnostic threshold to BP values of 130/80 mmHg is controversial and remains a moot point. In the 2009 review, the European guidelines on HT management¹⁶ increased the cut-off point set in 2007 (130/80 mmHg)⁵ to 140/85 mmHg. The results of the ACCORD (Action to Control Cardiovascular Risk in Diabetes) trial¹⁷ and INVEST (International Verapamil-Trandolapril Study)¹⁸ were still unknown at that time; these trials concluded that maintaining SBP below 130 mmHg in patients with HT and DM2 does not lead to a cardiovascular benefit.

The debate caused by lowering diagnostic thresholds may have influenced the absence of a HT diagnosis in diabetic patients with BP $< 140/90$ mmHg. With this cut-off point, the percentage of patients with HT who remained without diagnosis in 2009 decreased from 23.1% to 8.8%.

For the diagnostic threshold of 130/80 mmHg, 23.1% of all diabetic and hypertensive patients in the study area remained without a diagnosis in 2009. This result is slightly lower than the 29% reported in a population sample of 1507 diabetic patients in the United States¹⁹ and a little higher than the 20.2%⁹ and the

Table 2

Variables Associated With Appropriate Diagnosis of Hypertension on Multivariable Logistic Regression Analysis

	aOR (95%CI)	P
Patient-related variables		
<i>Sex</i>		
Male	1	
Female	1.288 (1.138-1.457)	<.001
<i>Age</i>		
	1.006 (1.001-1.011)	.028
<i>BMI</i>		
<25	1	
25-30	1.460 (1.235-1.727)	<.001
≥ 30	1.696 (1.443-1.993)	<.001
<i>Albuminuria</i>		
Normal (<30)	1	
Microalbuminuria (30-299)	1.250 (0.989-1.580)	.061
Macroalbuminuria (≥ 300)	1.785 (0.976-3.262)	.060
<i>Prior AMI</i>		
	0.448 (0.334-0.600)	<.001
<i>No depression</i>		
	1.630 (1.277-2.081)	<.001
<i>Antiplatelet treatment</i>		
	1.469 (1.286-1.677)	<.001
<i>Degree of hypertension at onset</i>		
BP 130-139/80-89 mmHg	1	
BP $\geq 140/90$ mmHg	2.770 (2.436-3.150)	<.001
Variables related to the use of resources		
<i>Visits to the clinic</i>		
	1.006 (1.001-1.011)	.011
<i>Type of antidiabetic treatment</i>		
No pharmacologic treatment	1	
With OAD	1.139 (0.983-1.319)	.083
With insulin (with or without OAD)	1.234 (1.030-1.478)	.023
Physician-related variables		
<i>Physician age</i>		
<40	1	
40-55	0.730 (0.639-0.834)	<.001
≥ 55	0.609 (0.458-0.810)	.001
<i>Professional situation</i>		
Permanent	1	
Temporary	0.849 (0.726-0.992)	.040
Substitute	0.783 (0.654-0.937)	.008

95%CI, 95% confidence interval; AMI, acute myocardial infarction; aOR, adjusted odds ratio; BMI, body mass index; BP, blood pressure; OAD, oral antidiabetic.

20.5%²⁰ in the 2 previous Spanish studies. Irrespective of the differences in methodology between the different studies, our data confirm the low level of HT diagnosis in patients with diabetes.

As in other reports, underdiagnosis of HT occurs mainly in men and younger patients,^{1,3,21,22} regardless of the diagnostic criteria used.

In our study, the mean delay in HT diagnosis by PC physicians after diagnostic BP values are recorded is greater in men, younger patients, and those with lower initial BP. To the best of our knowledge, this is the first study that estimates the delay in HT diagnosis, therefore we cannot compare our results with those of other studies.

Variables that were independently associated with a correct diagnosis include known factors related to cardiovascular risk and poor BP control, such as obesity^{20,23,26} or macroalbuminuria,^{24,25} which indicates that physicians diagnose HT earlier in patients at higher risk. This hypothesis would seem to be confirmed by the earlier diagnosis in patients with receiving more intensive treatment.

However, other associated risk factors are poor control of HT, male sex,^{22,26} younger age,²⁶ and a diagnosis of depression. In DM2 patients, depression is an additional risk factor because it is associated with a higher prevalence of complications and mortality.^{27,28}

Patients with a prior acute myocardial infarction were less likely to be diagnosed with HT than those free of myocardial ischemia history, which may be because the follow-up of HT in these patients is recorded in the acute myocardial infarction episode rather than as an independent diagnosis; ie, there may be underrecording rather than underdiagnosing. This hypothesis is supported by the finding that 98% of these patients had a prescription for hypotensive drugs, regardless of whether or not the HT diagnosis was recorded in the EMR. However, we cannot rule out that the lower probability of acute myocardial infarction patients being diagnosed may translate into a lack of awareness of the importance of high BP in patients with ischemic heart disease.

Regarding physician characteristics, it is known that they accept higher BP values than those recommended in clinical guidelines.^{29,30} This may be because of the "clinical inertia"³¹ that occurs when the physician recognizes the problem but does not make a related decision.

Our results reveal that greater seniority is related to fewer correct diagnoses of HT; this finding is in accordance with those of other studies showing that higher seniority decreases continuous professional development for physicians and compliance with the recommendations of clinical guidelines.^{32,33}

As in other studies, our results did not reveal a statistically significant relationship between physician sex and the accuracy of the diagnosis.^{30,34} However, this finding was associated with stability at work, which may be explained by greater knowledge of the population and increased involvement in patient follow-up.

Strengths and Limitations

Our study may have a classification bias, as the definition of HT does not consider whether patients were treated with drugs with a hypotensive effect that are prescribed for HT or other indications.

The definition of delayed diagnosis used in this study (>6 months from the availability of diagnostic BP values) was based on the fact that, for stable diabetics, clinical guidelines recommend a twice-yearly check-up visits. In addition, many PC physicians postpone recording the diagnosis between 3 and 6 months to recommend lifestyle changes for their patients or to assess the damage in target organs; such assessment may take 3 to 6 months.

Another possible limitation arises from the use of a secondary information source, the EMR, which is designed for care purposes. Nonetheless, the variables used appear sturdy and the quality of care partly depends on correct recording of BP readings.

The influence on delayed diagnosis of an improper referral from the nursing consultation to the medical consultation needs to be assessed. According to the visits scheduled in diabetic care programs, nurses follow-up the BP measurements, which are assessed in 6-monthly medical consultations. Only the physician was included in our analysis.

A strength of this study is that it collates available information for the entire diabetic population monitored in health centers and by all professionals, which reduces the bias related to voluntary participation.

Given that the responsibility for the diagnosis, treatment and monitoring of DM and HT falls mainly on PC professionals, this setting was deemed the most suitable to obtain information on real clinical conditions.

Strategies need to be implemented that improve the care of hypertensive patients in general and in the subgroup of younger

men in particular, to detect and treat HT early with the goal of reducing their cardiovascular risk.

CONCLUSIONS

The delay in PC diagnosis of HT in patients with diabetes varies according to the diagnostic thresholds used and is greater for the 130/80 mmHg diagnostic criteria.

More than half of the patients are diagnosed within the first 6 months.

Older patients, women, and those with initial BP values \geq 140/90 mmHg were more likely to receive a correct diagnosis. In addition, there was an inverse association between correct diagnosis and physician age older than 40 years and working in a less stable professional situation.

CONFLICTS OF INTEREST

None declared.

REFERENCES

- Banegas JR. Epidemiología de la hipertensión arterial en España. Situación actual y perspectivas. *Hipertensión*. 2005;22:353-62.
- Simonson DC. Etiology and prevalence of hypertension in diabetic patients. *Diabetes Care*. 1988;11:821-7.
- Castell MV, Martínez MA, Sanz J, García Puig J; Grupo MAPA-Madrid. Prevalencia, grado de conocimiento y control de la hipertensión arterial en una población española. Estudio MADRIC (Madrid Riesgo Cardiovascular). *Med Clin (Barc)*. 2010;135:671-2.
- Schutta MH. Diabetes and hypertension: epidemiology of the relationship and pathophysiology of factors associated with these comorbid conditions. *J Cardiometab Syndr*. 2007;2:124-30.
- The Task Force for the Management of Arterial Hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). 2007 guidelines for the management of arterial hypertension. *J Hypertens*. 2007;25:1005-87.
- Campbell NR, Leiter LA, Larochelle P, Tobe S, Chockalingam A, Ward R, et al. Hypertension in diabetes: a call to action. *Can J Cardiol*. 2009;25:299-302.
- Hansson L, Zanchetti A, Carruthers SC, Dahlöf B, Elmfeldt D, Julius S, et al. Effects of intensive blood-pressure lowering and low-dose aspirin in patients with hypertension: principal results of the Hypertension Optimal Treatment (HOT) randomised trial. *HOT Study Group. Lancet*. 1998;351:1755-62.
- Adler AI, Stratton IM, Neil HA, Yudkin JS, Matthews DR, Cull CA, et al. Association of systolic blood pressure with macrovascular and microvascular complications of type 2 diabetes (UKPDS 36): prospective observational study. *BMJ*. 2000;321:412-9.
- De Burgos-Lunar C, Salinero-Fort MA, Cárdenas-Valladolid J, Soto-Díaz S, Fuentes-Rodríguez CY, Abanades-Herranz JC, et al. Validation of diabetes mellitus and hypertension diagnosis in computerized medical records in primary health care. *BMC Med Res Methodol*. 2011;11:146.
- Chobanian AV, Bakris GL, Black HR, Cushman WC, Green LA, Izzo JL, et al. The Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure: the JNC 7 report. *JAMA*. 2003;289:2560-71.
- National Institute for Clinical Excellence (NICE). National Collaborating Centre for Chronic Conditions. In: Type 2 diabetes: national clinical guideline for management in primary and secondary care (update). London: Royal College of Physicians; 2008.
- MacMahon S, Peto R, Cutler J, Collins R, Sorlie P, Neaton J, et al. Blood pressure, stroke, and coronary heart disease. Part 1, prolonged differences in blood pressure: prospective observational studies corrected for the regression dilution bias. *Lancet*. 1990;335:765-74.
- Prospective Studies Collaboration. Age-specific relevance of usual blood pressure to vascular mortality: a meta-analysis of individual data for one million adults in 61 prospective studies. *Lancet*. 2002;360:1903-13.
- American Diabetes Association. Treatment of hypertension in adults with diabetes (position statement). *Diabetes Care*. 2003;26 Suppl 1:S80-2.
- Grupo de trabajo de la Guía de Práctica Clínica sobre Diabetes tipo 2. Guía de práctica clínica sobre diabetes tipo 2. Madrid: Plan Nacional para el SNS del MSC. Agencia de Evaluación de Tecnologías Sanitarias de El País Vasco; 2008. Guías de Práctica Clínica en el SNS: OSTEBA n.º 2006/08.
- Mancia G, Laurent S, Agabiti-Rosei E, Ambrosioni E, Burnier M, Caulfield MJ, et al. Reappraisal of European guidelines on hypertension management: a European Society of Hypertension Task Force document. *J Hypertens*. 2009;27:2121-58.
- Cushman WC, Evans GW, Byington RP, Goff Jr DC, Grimm Jr RH, Cutler JA, et al. Effects of intensive blood-pressure control in type 2 diabetes mellitus. *N Engl J Med*. 2010;362:1575-85.

18. Cooper-Dehoff RM, Gong Y, Handberg EM, Bavry AA, Denardo SJ, Bakris GL, et al. Tight blood pressure control and cardiovascular outcomes among hypertensive patients with diabetes and coronary artery disease. *JAMA*. 2010;304:61-8.
19. Geiss LS, Rolka DB, Engelgau MM. Elevated blood pressure among U.S. adults with diabetes, 1988-1994. *Am J Prev Med*. 2002;22:42-8.
20. García Vallejo O, Vicente Lozano J, Vegazo O, Jiménez Jiménez FJ, Llisterri Caro JL, Redón J. Control de la presión arterial de los pacientes diabéticos en el ámbito de atención primaria. Estudio DIAPA. *Med Clin (Barc)*. 2003;120:529-34.
21. Egan BM, Zhao Y, Axon RN. US trends in prevalence, awareness, treatment, and control of hypertension, 1988-2008. *JAMA*. 2010;303:2043-50.
22. Pereira M, Lunet N, Azevedo A, Barros H. Differences in prevalence, awareness, treatment and control of hypertension between developing and developed countries. *J Hypertens*. 2009;27:963-75.
23. Llisterri JL, Barrios V, De la Sierra A, Bertomeu V, Escobar C, González-Segura D. Control de la presión arterial en las mujeres hipertensas de 65 o más años de edad asistidas en atención primaria. Estudio MERICAP. *Rev Esp Cardiol*. 2011;64:654-60.
24. Abellán Alemán J, Prieto Díaz MA, Leal Hernández M, Balanza Galindo SB, De la Sierra Iserte AD, Martell Claros N, et al. Evaluación y control de los hipertensos diabéticos atendidos en centros de Atención Primaria de España. Estudio BRAND II. *Aten Primaria*. 2011;43:297-304.
25. Robles NR, Velasco J, Mena C, Angulo E, Garrote T; en representación de los investigadores del estudio MICREX. Microalbuminuria en pacientes diabéticos y en pacientes con hipertensión arterial: estudio de una cohorte de 979 pacientes. *Med Clin (Barc)*. 2006;25:761-4.
26. Llisterri Caro JL, Rodríguez Roca GC, Alonso Moreno FJ, Barrios Alonso V, Banegas Banegas JR, González-Segura Alsina D, et al. Evolución del control de la presión arterial en España en el período 2002-2006. Estudios PRESCAP. *Hipertension*. 2009;26:257-65.
27. Katon WJ, Rutter C, Simon G, Lin E, Ludman E, Ciechanowski P, et al. The association of comorbid depression with mortality in patients with type 2 diabetes. *Diabetes Care*. 2005;28:2668-72.
28. Egede LE, Nietert PJ, Zheng D. Depression and all-cause and coronary heart disease mortality among adults with and without diabetes. *Diabetes Care*. 2005;28:1339-45.
29. Mehta SS, Wilcox CS, Schulman KA. Treatment of hypertension in patients with comorbidities: results from the study of hypertensive prescribing practices (SHyPP). *Am J Hypertens*. 1999;12:333-40.
30. Midlöv P, Ekesbo R, Johansson L, Gerward S, Persson K, Nerbrand C, et al. Barriers to adherence to hypertension guidelines among GPs in southern Sweden: a survey. *Scand J Prim Health Care*. 2008;26:154-9.
31. Phillips LS, Branch WT, Cook CB, Doyle JP, El-Kebbi IM, Gallina DL, et al. Clinical inertia. *Ann Intern Med*. 2001;135:825-34.
32. Choudry NK, Fletcher RH, Soumerai SB. Systematic review: the relationship between clinical experience and quality of health care. *Ann Intern Med*. 2005;142:260-73.
33. Moride Y, Du Fort GG, Monette J, Ducruet T, Boivin JF, Champoux N, et al. Suboptimal duration of antidepressant treatments in the older ambulatory population of Quebec: association with selected physician characteristics. *J Am Geriatr Soc*. 2002;50:1365-71.
34. Janssen PG, Gorter KJ, Stolk RP, Rutten GE. Do characteristics of practices and general practitioners influence the yield of diabetes screening in primary care? The ADDITION Netherlands study. *Scand J Prim Health Care*. 2008;26:160-5.