

Original article

Health-related Quality of Life of Patients With Chronic Systolic Heart Failure in Spain: Results of the VIDA-IC Study



Josep Comín-Colet,^{a,b,c,*} Manuel Anguita,^d Francesc Formiga,^e Luis Almenar,^f María G. Crespo-Leiro,^g Luis Manzano,^h Javier Muñoz,ⁱ José Chaves,^j Trinidad de Frutos,^j and Cristina Enjuanes^{a,b,c} on behalf of the VIDA-IC (Quality of Life and Heart Failure in Spain: Current Situation) multicenter study researchers[◇]

^a Programa de Insuficiencia Cardíaca, Servicio de Cardiología, Hospital del Mar, Barcelona, Spain

^b Grupo de Investigación Biomédica en Enfermedades del Corazón, Programa de Investigación en Procesos Inflamatorios y Cardiovasculares, Instituto Hospital del Mar de Investigaciones Médicas (IMIM), Barcelona, Spain

^c Departamento de Medicina, Universidad Autónoma de Barcelona, Barcelona, Spain

^d Programa de Insuficiencia Cardíaca y Trasplante, Servicio de Cardiología, Hospital Universitario Reina Sofía, Córdoba, Spain

^e Servicio de Medicina Interna, Hospital Universitario de Bellvitge, L'Hospitalet de Llobregat, Barcelona, Spain

^f Unidad de Insuficiencia Cardíaca y Trasplante, Servicio de Cardiología, Hospital Universitario La Fe, Valencia, Spain

^g Unidad de Insuficiencia Cardíaca Avanzada y Trasplante Cardíaco, Servicio de Cardiología, Instituto de Investigación Biomédica de A Coruña (INIBIC), Complejo Hospitalario Universitario de A Coruña (CHUAC), SERGAS, Universidade da Coruña (UDC), A Coruña, Spain

^h Unidad de Insuficiencia Cardíaca y Riesgo Vascular en el Anciano, Servicio de Medicina Interna, Hospital Universitario Ramón y Cajal, Madrid, Spain

ⁱ Instituto Universitario de Ciencias de la Salud, Instituto de Investigación Biomédica de A Coruña (INIBIC), Universidade da Coruña, A Coruña, Spain

^j Departamento Médico de Pfizer S.L.U., Madrid, Spain

Article history:

Received 17 March 2015

Accepted 16 July 2015

Available online 23 December 2015

Keywords:

Heart failure

Health-related quality of life

Specific and generic quality of life

questionnaires

Real life or routine clinical practice

ABSTRACT

Introduction and objectives: Although heart failure negatively affects the health-related quality of life of Spanish patients there is little information on the clinical factors associated with this issue.

Methods: Cross-sectional multicenter study of health-related quality of life. A specific questionnaire (Kansas City Cardiomyopathy Questionnaire) and a generic questionnaire (EuroQoL-5D) were administered to 1037 consecutive outpatients with systolic heart failure.

Results: Most patients with poor quality of life had a worse prognosis and increased severity of heart failure. Mobility was more limited and rates of pain/discomfort and anxiety/depression were higher in the study patients than in the general population and patients with other chronic conditions. The scores on both questionnaires were very highly correlated (Pearson $r = 0.815$; $P < .001$). Multivariable linear regression showed that being older (standardized $\beta = -0.2$; $P = .03$), female (standardized $\beta = -10.3$; $P < .001$), having worse functional class (standardized $\beta = -20.4$; $P < .001$), a higher Charlson comorbidity index (standardized $\beta = -1.2$; $P = .005$), and recent hospitalization for heart failure (standardized $\beta = 6.28$; $P = .006$) were independent predictors of worse health-related quality of life.

Conclusions: Patients with heart failure have worse quality of life than the general Spanish population and patients with other chronic diseases. Female sex, being older, comorbidity, advanced symptoms, and recent hospitalization are determinant factors in health-related quality of life in these patients.

© 2015 Sociedad Española de Cardiología. Published by Elsevier España, S.L.U. All rights reserved.

* Corresponding author: Programa de Insuficiencia Cardíaca, Servicio de Cardiología, Hospital del Mar (Parc de Salut Mar), Pg. Marítim 25-29, 08003 Barcelona, Spain.
E-mail address: josepcomin@gmail.com (J. Comín-Colet).

◇ The names of the VIDA-IC study researchers are shown in the [supplementary material](#).

Calidad de vida relacionada con la salud de los pacientes con insuficiencia cardiaca crónica sistólica en España: resultados del estudio VIDA-IC

RESUMEN

Palabras clave:

Insuficiencia cardiaca
Calidad de vida relacionada con la salud
Cuestionarios de calidad de vida específicos y genéricos
Vida real o práctica clínica habitual

Introducción y objetivos: La calidad de vida relacionada con la salud de los pacientes con insuficiencia cardiaca está afectada. Hay poca información sobre los factores clínicos asociados a esta mala calidad de vida de la población española con insuficiencia cardiaca.

Métodos: Estudio multicéntrico transversal de calidad de vida relacionada con la salud aplicando un cuestionario específico (*Kansas City Cardiomyopathy Questionnaire*) y otro genérico (*EuroQoL-5D*) a 1.037 pacientes ambulatorios consecutivos con insuficiencia cardiaca sistólica.

Resultados: Los pacientes con peor calidad de vida presentaron en su mayoría datos asociados a peor pronóstico y mayor gravedad de la enfermedad. Los pacientes del estudio presentaron mayor incidencia de limitaciones en movilidad, dolor/malestar y ansiedad/depresión cuando se realizó una comparación externa con población general y con pacientes con otras afecciones crónicas. La correlación entre las puntuaciones totales de ambos cuestionarios fue muy alta (r de Pearson = 0,815; $p < 0,001$). Con regresión lineal multivariable, se observó que mayor edad (β estandarizada = -0,2; $p = 0,03$), sexo femenino (β estandarizada = -10,3; $p < 0,001$), peor clase funcional (β estandarizada = -20,4; $p < 0,001$), mayor comorbilidad según índice de Charlson (β estandarizada = -1,2; $p = 0,005$) y el ingreso reciente por insuficiencia cardiaca (β estandarizada = 6,28; $p = 0,006$) son factores independientes predictores de peor calidad de vida relacionada con la salud.

Conclusiones: Los pacientes con insuficiencia cardiaca tienen muy afectada su calidad de vida respecto a la población general española y a otras enfermedades crónicas. Sexo femenino, edad avanzada, comorbilidad, síntomas avanzados y hospitalización reciente son factores determinantes en la calidad de vida relacionada con la salud de estos pacientes.

© 2015 Sociedad Española de Cardiología. Publicado por Elsevier España, S.L.U. Todos los derechos reservados.

Abbreviations

CHF: chronic heart failure
EQ-5D: EuroQoL-5D overall quality of life questionnaire
HF: heart failure
HRQoL: health-related quality of life
KCCQ: Kansas City Cardiomyopathy Questionnaire
VAS: visual analogue scale

INTRODUCTION

Patients with chronic heart failure (CHF) have far worse health-related quality of life (HRQoL) than the general population and patients with other chronic diseases.¹ Improving HRQoL is one of the main objectives of the comprehensive management of CHF patients.^{2–4}

In CHF patients, HRQoL is a multidimensional measure with good correlation with disease severity,⁵ provides independent prognostic information, and can assist in assessing the cost-effectiveness of implementing new therapeutic options.^{6,7}

The deterioration of HRQoL in CHF patients is reflected in the dimensions that capture information on the functional limitations that have a particular impact on the mobility or daily activities domains.⁸

Several authors have addressed the extent to which the HRQoL of CHF patients differs from that of the general population or patients with other chronic diseases, which dimensions or domains of HRQoL are the most affected, and which clinical and demographic factors influence HRQoL. However, there is little information on HRQoL in heart failure (HF) patients in Spain, since publications to date in this field come from substudies of clinical trials or studies conducted in other geographical and cultural settings, and thus it remains unknown if the results are fully transferable to the Spanish setting.^{9,10}

Thus, the aims of the prespecified analysis of the VIDA-IC study, whose first results were published in 2014,¹¹ were: a) to determine

the clinical-demographic factors associated with HRQoL in patients with CHF and left ventricular systolic dysfunction followed up in cardiology or internal medicine clinics; b) to assess which dimensions were most affected in these patients, and c) to explore if there was a gradient of total scores and by specific domains in HRQoL instruments between the study patients and the general population or in patients with other chronic conditions in Spain.

METHODS

Study Design

The VIDA-IC observational descriptive study was conducted throughout Spain from October 2011 to January 2012 by 115 specialists (cardiologists and internal medicine specialists), who included consecutive CHF patients seen in an outpatient clinic.¹¹ The objectives of the study were to assess the level of correlation between specific and generic HRQoL measures in CHF patients, study the factors determining the level of HRQoL, and contextualize the quality of life of HF patients measured with generic scales with the quality of life measured with the same scales in the general population or patients with other chronic diseases in Spain. The latter objective was fulfilled by comparing the general information available in the literature and in public national health surveys on quality of life in the general Spanish population and the population with chronic diseases in Spain. The study protocol was approved by the Ethics and Clinical Research Committee of the *Instituto Hospital del Mar de Investigaciones Médicas* (IMIM; Barcelona, Spain). All patients gave written informed consent before inclusion in the study.

Study Population and Inclusion and Exclusion Criteria

The study consecutively included patients attending a specialized outpatient clinic (cardiology or internal medicine) who fulfilled the following inclusion criteria: clinically stable, older

Table 1
Demographic and Clinical Characteristics of all Patients Included in the Study According to Health-related Quality of Life

Variables	Total (n = 1037)	Patients with better HRQoL* (n = 696)	Patients with worse HRQoL (n = 327)	P
Age, y	70.6 ± 11.1	69.2 ± 11.2	73.6 ± 10.2	< .0001
Women	309 (30.1)	175 (25.3)	129 (39.9)	< .001
BMI	27.7 ± 3.9	27.6 ± 3.6	27.9 ± 4.5	.343
Systolic blood pressure, mmHg	127.2 ± 18.7	127.3 ± 17.7	127 ± 20.7	.807
Heart rate, bpm	73.9 ± 15.7	73.4 ± 15.7	75.2 ± 15.6	.09
NYHA I-II/III-IV	550 (54.9)/452 (45.1)	481 (71.8)/189 (28.2)	59 (18.5)/260 (81.5)	< .001
LVEF, %	33.7 ± 6.8	34.4 ± 6.4	32.2 ± 7.5	< .0001
Charlson Index	4.4 ± 2.8	3.9 ± 2.5	5.2 ± 3.1	< .0001
Ischemic etiology	527 (50.8)	345 (49.6)	175 (53.5)	.239
Comorbidities				
Hypertension	821 (79.2)	539 (77.4)	271 (82.9)	.046
Diabetes mellitus	456 (44.0)	288 (41.4)	160 (48.9)	.023
Significant kidney failure	244 (23.5)	126 (18.1)	115 (35.2)	< .001
Atrial fibrillation	447 (45.5)	279 (42.1)	161 (52.8)	.002
Anemia	202 (21.3)	110 (17.1)	90 (30.6)	< .001
Treatment				
ACE inhibitors or ARB	929 (89.6)	633 (91.0)	283 (86.5)	.032
Beta blockers	794 (76.6)	544 (78.2)	238 (72.8)	.059
Aldosterone antagonists	689 (66.4)	451 (64.8)	228 (69.7)	.12
Ivabradine	91 (8.8)	64 (9.2)	27 (8.3)	.623
Digoxin	225 (21.7)	137 (19.7)	85 (26.0)	.022
Diuretics	925 (89.2)	605 (86.9)	306 (93.6)	.001
Statins	786 (75.8)	533 (76.6)	240 (73.4)	.269
Antiplatelet agents	622 (60.0)	419 (60.2)	193 (59.0)	.72
Anticoagulants	414 (39.9)	253 (36.4)	156 (47.7)	.001
Laboratory values				
Hemoglobin, g/dL	12.9 ± 1.7	13.0 ± 1.6	12.5 ± 1.7	< .0001
EGFR, mL/min/1.73 m ²	61.2 ± 27.6	64.9 ± 27.7	53.7 ± 26.2	< .0001
Creatinine clearance < 60	260 (45.2)	138 (36.4)	117 (63.2)	< .001
NT-proBNP, pg/mL	1854.1 ± 1829.8	1560.2 ± 1361.6	2491.6 ± 2489.2	.005
BNP, pg/mL	515.0 ± 1871.8	616.2 ± 2342.8	341.0 ± 280.1	.253

ACE, angiotensin-converting enzyme; ARB, angiotensin receptor blockers; BMI, body mass index; BNP, brain natriuretic peptide; EGFR, estimated glomerular filtration rate; HRQoL, health-related quality of life; LVEF, left ventricular ejection fraction; NT-proBNP, N-terminal pro-brain natriuretic peptide; NYHA, New York Heart Association. Data are expressed as No. (%) or mean ± standard deviation.

* Better health-related quality of life was defined as a Kansas City Cardiomyopathy Questionnaire overall summary score of ≥ 50 points.

than 18 years, and a diagnosis of CHF with systolic dysfunction (left ventricular ejection fraction ≤ 40%) within the last 12 months. Exclusion criteria were: waiting for heart transplantation or correction of valvular lesions, inability to understand or complete the HRQoL questionnaires, noncardiac disease with a life expectancy of less than 1 year, noncardiovascular hospitalization in the month prior to inclusion, or hospitalization at the time of inclusion. Patient inclusion was stratified according to recent (less than 1 month) admission for HF and nonrecent (more than 6 months) admission for HF at a ratio of 1:1 for each of the recruiters. Baseline data were obtained from eligible patients or medical records after they had given informed consent, provided the patients were stable and had no signs of acute decompensation.

Evaluation of Results in Patient-centered Health: Quality of Life

All patients in the study were asked to complete the self-administered Kansas City Cardiomyopathy Questionnaire (KCCQ)¹² and the EuroQoL-5D overall quality of life questionnaire (EQ-5D).¹³ The KCCQ is specific to patients with HF. It comprises 23 items in 7 domains.

The score on each domain can, in theory, range from 0 to 100, with 100 corresponding to the best state. In addition, 3 summary scores are calculated: the symptoms summary score is derived by summing the scores on the frequency and severity of symptoms (excluding stability); the clinical summary score is derived by summing the scores on the physical limitations and symptoms domains; and the overall summary score is derived by summing the clinical summary score and the quality of life and social limitation scores. The EQ-5D is a generic instrument comprising a visual analogue scale (VAS) of self-rated general health and 5 dimensions (mobility, self-care, daily activities, pain/discomfort, and anxiety/depression). Scores on the VAS can range from 0 (worse state) to 100 (best state). Scores on the 5 dimensions can be expressed as an overall summary index (EQ-5D index) or as the percentage of patients who indicate some kind of problem on each of the dimensions. Both scales have been validated for the Spanish general population.¹³

The HRQoL of the study patients, the Spanish general population, and patients with other chronic diseases was compared using summary data of the VAS and the 5 dimensions of the EQ-5D. These data were obtained from the most recent Spanish National Health Survey of the general population¹⁴ and

publications using the EQ-5D to assess HRQoL in Spanish patients with various chronic diseases.^{15–18}

Statistical Analysis

Continuous variables are expressed as mean \pm standard deviation and discrete variables as absolute and relative values. Groups with good and poor HRQoL were compared using the chi square test and Student *t* test (or Mann-Whitney U test as needed) for discrete and continuous variables, respectively. The level of correlation between the overall KCCQ and EQ-5D scores was assessed using correlation coefficients and Spearman's ρ and Pearson's *r*. Clinical and demographic factors associated with HRQoL were assessed using univariable logistic regression models and univariable linear regression models in which the dependent variables were the overall summary scores of the KCCQ, the EQ-5D index, and the VAS, and the independent variables were specific demographic and clinical factors included in this study. The independent variables were used to construct several exploratory multivariable linear regression models using the backward stepwise method to determine which factors remained independently associated with patient-centered health outcomes. A *P* value of $< .05$ was used as a cutoff for statistical significance. All analyses were performed using the SPSS statistical

software package version 18 and the Stata statistical software package version 11.

RESULTS

A total of 1037 patients with HF and left ventricular systolic dysfunction were included in the study. Of these, 63.2% were recruited by cardiologists and the remaining 36.8% were recruited by internal medicine specialists. A total of 1037 KCCQ, 1020 VAS, and 1009 EQ-5D completed HRQoL questionnaires were available for analysis. Table 1 shows the characteristics of the study patients. Mean age was 72 (interquartile range: 64–78) years and there was a predominance of men. About half of the patients had ischemic HF and were in New York Heart Association (NYHA) functional class III/IV. In general, patients with worse HRQoL scores on the KCCQ showed data associated with poor prognosis and increased severity of CHF.

Compared with the general reference population (Figure 1),^{14–20} the study patients reported more limitations on all the dimensions of the EQ-5D. In some dimensions, such as mobility, pain/discomfort, and anxiety/depression, HF patients had more limitations than patients with chronic diseases, such as diabetes mellitus, cancer, or Alzheimer's disease. Patients with HF and NYHA III/IV comprised




Figure 1. Comparison of the impact on health-related quality of life in study patients with heart failure compared with the Spanish general population and people with other chronic diseases in Spain. A: percentage of people and patients reporting any limitation in each dimension of the EuroQoL-5D. B: comparative analysis of scores (mean \pm standard deviation) on the EuroQoL-5D visual analogue scale. COPD, chronic obstructive pulmonary disease; DM, diabetes mellitus; EuroQoL-5D, EuroQoL-5D overall quality of life questionnaire; HF, heart failure; NYHA, New York Heart Association functional class.




Figure 1. (Continued).

almost half the study population and reported similar or higher levels of limitations in the dimensions studied than patients with a history of stroke or those with chronic kidney failure on dialysis. Similar results were obtained when the mean scores of the VAS were analyzed. According to the VAS score, the perceived overall state of health of the study patients with CHF was worse than that of the general population, patients with chronic obstructive pulmonary disease or cancer, and similar to that of patients with diabetes or

pulmonary hypertension. Patients with CHF in NYHA functional class III/IV had a lower mean VAS score, indicating a worse perceived overall health state even when compared with patients with a history of stroke or Alzheimer's disease or patients on dialysis.

Table 2 shows the average scores of each subdomain of the KCCQ, the summary scores, the mean scores of the EQ-5D index and the VAS, and the percentage of patients with some degree of limitation on each of the EQ-5D. As expected, patients with worse

Table 2

Distribution of the Summary Scores, Dimensions, and Various Domains of Quality-of-life Questionnaires Specific to Heart Failure (Kansas City Cardiomyopathy Questionnaire) and Generic Quality-of-life Questionnaires Used in the Total Study Population According to Health-related Quality of Life

	Total (n = 1037)	Patients with better HRQoL (n = 696)	Patients with worse HRQoL (n = 327)	P
<i>KCCQ, domains</i>				
Physical limitation	61.1 ± 28.1	75.7 ± 18.2	29.9 ± 18.1	< .0001
Symptom stability	59.5 ± 23.2	63.0 ± 21.2	51.9 ± 25.4	< .0001
Frequency of symptoms	66.3 ± 26.1	79.9 ± 15.3	37.1 ± 19.2	< .0001
Symptom burden	67.1 ± 26.1	80.7 ± 16.0	37.5 ± 17.6	< .0001
Self-efficacy	69.1 ± 22.5	72.9 ± 20.2	60.6 ± 24.6	< .0001
Quality of life	54.4 ± 24.1	66.6 ± 16.6	28.1 ± 14.4	< .0001
Social limitation	61.6 ± 29.4	77.7 ± 17.9	27.3 ± 17	< .0001
<i>KCCQ, summary measures</i>				
Overall summary score	60.9 ± 24.5	75.1 ± 13.5	30.6 ± 12.3	< .0001
Clinical summary score	63.9 ± 25.2	78.0 ± 14.4	33.6 ± 14.4	< .0001
Symptom summary score	66.7 ± 25.4	80.3 ± 15.0	37.3 ± 17.0	< .0001
<i>EQ-5D, patients who reported problems</i>				
Mobility	586 (58.1)	273 (40.7)	304 (93.5)	< .001
Self-care	382 (38.0)	132 (19.7)	246 (76.4)	< .001
Daily activities	619 (61.4)	307 (45.8)	305 (93.8)	< .001
Pain/discomfort	510 (50.6)	256 (38.1)	248 (76.8)	< .001
Anxiety/depression	493 (48.9)	237 (35.3)	249 (76.9)	< .001
<i>EQ-5D, summary measures</i>				
Overall EQ-5D Index	0.6 ± 0.3	0.8 ± 0.2	0.4 ± 0.2	< .0001
Visual analogue scale	60.8 ± 20	68.7 ± 15.8	43.5 ± 16.8	< .0001

EQ-5D, EuroQoL-5D overall quality of life questionnaire; HRQoL, health-related quality of life; KCCQ, Kansas City Cardiomyopathy Questionnaire. Values are expressed as no. (%) or mean ± standard deviation.

* Better health-related quality of life was defined as a Kansas City Cardiomyopathy Questionnaire overall summary score of ≥ 50 points.




Figure 2. Demographic and clinical factors associated with the health-related quality of life perceived by patients assessed using the Kansas City Cardiomyopathy Questionnaire (A), the EuroQoL-5 dimensions (B), and the EuroQoL-5D visual analogue scale (C). Quality of life related to poor health: overall Kansas City Cardiomyopathy Questionnaire summary score < 50; EuroQoL-5D index < 0.5; visual analogue scale < 50. Analyses were performed using univariable binary logistic regression models. 95%CI, 95% confidence interval; ACE, angiotensin converting enzyme; ARB, angiotensin receptor blockers; BMI: body mass index; HF, heart failure; HRQoL: health-related quality of life; NT-proBNP: N-terminal pro-brain natriuretic peptide. *Reference category.




Figure 2. (Continued)




Figure 2. (Continued).

Table 3
Correlation Matrix (R-values and Confidence Intervals) of the Different Items, Dimensions, Domains, and Summary Scores of the Questionnaires Used to Assess Health-related Quality of Life

	1	2	3	4	5	6	7	8	9	10	11	12
1. EuroQoL-5 dimensions score	1											
2. Visual analogue scale	0.682 (0.647-0.714)	1										
3. Physical limitations	0.785 (0.759-0.807)	0.634 (0.596-0.669)	1									
4. Symptom stability	0.207 (0.147-0.266)	0.229 (0.170-0.286)	0.176 (0.116-0.235)	1								
5. Frequency of symptoms	0.717 (0.685-0.745)	0.601 (0.560-0.639)	0.756 (0.729-0.781)	0.203 (0.144-0.261)	1							
6. Symptom burden	0.721 (0.689-0.749)	0.633 (0.595-0.669)	0.756 (0.729-0.781)	0.240 (0.182-0.297)	0.903 (0.891-0.914)	1						
7. Overall symptom score	0.736 (0.706-0.763)	0.633 (0.594-0.668)	0.775 (0.749-0.798)	0.227 (0.168-0.284)	0.976 (0.972-0.978)	0.976 (0.972-0.978)	1					
8. Self-efficacy	0.349 (0.293-0.402)	0.301 (0.244-0.356)	0.322 (0.266-0.375)	0.0909* (0.030-0.151)	0.340 (0.284-0.392)	0.345 (0.290-0.398)	0.351 (0.296-0.403)	1				
9. Quality of life	0.701 (0.668-0.731)	0.629 (0.591-0.665)	0.695 (0.662-0.726)	0.210 (0.151-0.268)	0.779 (0.754-0.802)	0.788 (0.763-0.810)	0.802 (0.780-0.823)	0.318 (0.262-0.372)	1			
10. Social limitations	0.751 (0.723-0.777)	0.645 (0.607-0.679)	0.822 (0.801-0.841)	0.208 (0.149-0.266)	0.790 (0.766-0.812)	0.795 (0.772-0.817)	0.812 (0.790-0.832)	0.300 (0.243-0.354)	0.815 (0.793-0.834)	1		
11. Overall summary score	0.815 (0.792-0.834)	0.698 (0.665-0.729)	0.905 (0.893-0.916)	0.224 (0.165-0.281)	0.897 (0.884-0.908)	0.903 (0.891-0.914)	0.921 (0.912-0.930)	0.352 (0.297-0.404)	0.898 (0.885-0.909)	0.944 (0.937-0.951)	1	
12. Clinical summary score	0.807 (0.784-0.828)	0.672 (0.637-0.705)	0.948 (0.941-0.954)	0.211 (0.152-0.269)	0.913 (0.903-0.923)	0.913 (0.903-0.923)	0.936 (0.928-0.943)	0.358 (0.303-0.410)	0.792 (0.768-0.814)	0.866 (0.850-0.881)	0.968 (0.964-0.972)	1

* Correlation coefficient: $P = .004$; all other correlation coefficients: $P < .0001$.

Table 4

Univariable and Multivariable Linear Regression Models used to Assess Demographic and Clinical Factors Associated with the Health-related Quality of Life as Measured by the Kansas City Cardiomyopathy Questionnaire Summary Score, the EuroQoL-5 Dimensions Overall Index, and the Visual Analogue Scale

	Univariable								
	KCCQ OSS			EQ-5D Index			VAS		
	β^*	R ²	P	β^*	R ²	P	β^*	R ²	P
Age, 1 y	-0.275	0.076	<.001	-0.287	0.082	<.001	-0.223	0.050	<.001
Sex, male/female	-0.157	0.025	<.001	-0.169	0.029	<.001	-0.108	0.012	.001
BMI, 1	-0.056	0.003	.077	-0.082	0.007	.010	-0.016	<0.001	.610
Systolic blood pressure, 1 mmHg	-0.010	<0.001	.743	0.001	<0.001	.965	0.041	0.002	.194
Heart rate, 1 bpm	-0.103	0.011	.001	-0.093	0.009	.003	-0.059	0.004	.059
NYHA functional class I-II/III-IV	-0.562	0.316	<.001	-0.465	0.216	<.001	-0.453	0.206	<.001
LVEF, 1%	0.156	0.024	<.001	0.129	0.017	<.001	0.165	0.027	<.001
Charlson index, 1 point	-0.285	0.081	<.001	-0.318	0.101	<.001	-0.240	0.058	<.001
Ischemic etiology, no/yes	-0.068	0.005	.030	-0.086	0.007	.006	-0.050	0.003	.114
EGFR, 1 mL/min/ 1.73 m ²	0.193	0.037	<.001	0.187	0.035	<.001	0.173	0.030	<.001
Hypertension, no/yes	-0.097	0.010	.002	-0.126	0.016	<.001	-0.040	0.002	.202
Atrial fibrillation, no/yes	-0.152	0.023	<.001	-0.172	0.030	<.001	-0.145	0.021	<.001
DM, no/yes	-0.149	0.022	<.001	-0.164	0.027	<.001	-0.106	0.011	.001
Hemoglobin, 1 g/dL	0.227	0.051	<.001	0.245	0.060	<.001	0.214	0.046	<.001
Optimal treatment, no/yes	0.028	0.001	.366	0.042	0.002	.188	0.038	0.001	.230
Inclusion service, CAR/IM	-0.186	0.035	<.001	-0.197	0.039	<.001	-0.185	0.034	<.001
Recent admission, yes/no	0.259	0.067	<.001	0.201	0.041	<.001	0.195	0.038	<.001
Time since diagnosis <1 y, no/yes	-0.070	0.005	.034	-0.067	0.004	.046	-0.072	0.005	.029
	Multivariable (stepwise backward method)								
	KCCQ OSS			EQ-5D index			VAS		
	β^*		P	β^*		P	β^*		P
Age, 1 y	-0.230		.030	-0.004		.002	-0.178		.072
Sex, male/female	-10.258		<.001	-0.105		<.001	-3.683		.095
BMI, 1									
Systolic blood pressure, 1 mmHg									
Heart rate, 1 bpm									
NYHA functional class I-II/III-IV	-20.373		<.001	-0.180		<.001	-12.586		<.001
LVEF, 1%	0.254		.135				0.263		.086
Charlson index, 1 point	-1.258		.005	-0.008		.136	-1.029		.009
Ischemic etiology, no/yes				-0.053		.055			
EGFR, 1 mL/min/ 1.73 m ²									
Hypertension, no/yes				-0.060		.085			
Atrial fibrillation, no/yes									
DM, no/yes				-0.041		.140			
Hemoglobin, 1 g/dL						1.023	0.087		
Optimal treatment, no/yes									
Inclusion service, CAR/IM	-4.595		.049	-0.035		.185	-4.761		.022
Recent admission, yes/no	6.286		.006	0.046		.075			
Time since diagnosis <1 y, no/yes									
R ² adjusted for each model		0.3690			0.3151			0.2534	

β , standardized beta coefficient; BMI, body mass index; CAR/IM, cardiology/internal medicine; DM, diabetes mellitus; EGFR, estimated glomerular filtration rate; EQ-5D, EuroQoL-5D overall quality of life questionnaire; KCCQ OSS, Kansas City Cardiomyopathy Questionnaire overall summary score; LVEF, left ventricular ejection fraction; NYHA, New York Heart Association.

* The first dichotomous variable is the reference category.

Table 5 Unadjusted Analysis of the Mean Scores of the Overall Kansas City Cardiomyopathy Questionnaire Summary Score, Visual Analogue Scale, and the Overall EuroQoL-5D Index Quality of Life Questionnaire in the Patient Subgroups With an Independent Association With The Kansas City Cardiomyopathy Questionnaire Summary Score in the Multivariable Linear Regression Analysis

	Sex		NYHA functional class		Age		Charlson comorbidity index		Service		Recent hospitalization	
	Men (n = 719)	Women (n = 309)	I-II (n = 550)	III-IV (n = 452)	< 75 y (n = 601)	≥ 75 y (n = 418)	≤ 4 (n = 324)	> 4 (n = 207)	Cardiology (n = 638)	Internal medicine (n = 367)	No (n = 647)	Yes (n = 386)
KCCQ OSS	63.4 ± 24.0	55.0 ± 24.6	73.4 ± 18.8	45.6 ± 22.0	64.8 ± 24.2	55.2 ± 23.9	62.5 ± 23.8	54.5 ± 23.9	66.1 ± 22.2	56.6 ± 23.9	65.3 ± 23.6	53.7 ± 24.5
Visual analogue scale	62.2 ± 19.4	57.5 ± 20.8	68.9 ± 16.6	50.7 ± 19.1	63.5 ± 19.7	56.9 ± 19.9	61.6 ± 20.3	56.5 ± 20.1	63.7 ± 18.6	56.0 ± 23.9	63.1 ± 19.9	57.1 ± 19.7
EQ-5D Index	0.7 ± 0.3	0.6 ± 0.3	0.8 ± 0.2	0.5 ± 0.2	0.7 ± 0.3	0.6 ± 0.2	0.7 ± 0.3	0.6 ± 0.3	0.7 ± 0.2	0.6 ± 0.3	0.7 ± 0.2	0.6 ± 0.3

EQ-5D, EuroQoL-5D overall quality of life questionnaire; KCCQ, Kansas City Cardiomyopathy Questionnaire; NYHA, New York Heart Association functional class. All differences between groups: $P < .05$. Mean ± standard deviation according to sex (male vs female), NYHA functional class (I-II vs IV-III), age (< 75 years vs ≥ 75 years), Charlson comorbidity index (≤ 4 [median] vs > 4), time since last admission (< 30 days vs ≥ 30 days), and clinical service (cardiology vs internal medicine).

HRQoL scored worse on all these items. Items such as self-efficacy or symptom stability, which are not included in the overall summary, were significantly worse in patients with worse HRQoL. Similarly, patients with an overall summary score on the KCCQ less than 50 points had more limitations on the 5 dimensions of EQ-5D and lower mean scores on this questionnaire and the VAS.

A correlation matrix was used to study the associations between the HRQoL questionnaire variables used in this study (Table 3). The table shows a horizontal and vertical list of the same variables with their correlation coefficients (R) expressed as a number ranging from 0 to 1 and their confidence intervals. Table 3 shows the correlations between the scores on the KCCQ domains and the summary scores of the KCCQ, the EQ-5D index, and the VAS. The correlations between the overall scores of the EQ-5D and KCCQ were very high (Pearson's $r = 0.815$; Spearman's $\rho = 0.811$; $P < .001$ for both coefficients). Significantly high correlations (> 0.6) were found between the KCCQ domains and between these domains and the KCCQ and EQ-5D summary scores in all cases where convergent correlation would be expected. Regarding the overall KCCQ scores, the physical limitation domain and total symptoms summary score had very high convergence (coefficients > 0.8). These correlations were relatively lower in relation to the VAS and EQ-5D index. The correlations that, although significant, were more divergent regarding the other dimensions and the KCCQ or the EQ-5D summary scores were those obtained from symptom stability and the self-efficacy domain of the KCCQ (ranges between 0.1 and 0.2 in most cases).

Figure 2 and Table 4 show the clinical factors associated with worse HRQoL. The multiple linear regression analysis (Table 4) showed that advanced age, female sex, worse functional class, and greater comorbidity were independent predictors of worse quality of life. The treatment of patients in cardiology services was independently associated with better quality of life, which is probably associated with a better clinical profile. These clinical features had an effect on the statistical significance of the differences in the raw summary scores of the instruments (Table 5) and the percentage of problems identified in each dimension of the EQ-5D (Figure 3).

DISCUSSION

This multicenter Spanish study showed that patients with HF and systolic dysfunction experience marked changes in HRQoL. Specifically, in patients with HF and advanced functional class, the level of HRQoL is similar to or even worse than that seen in patients with chronic obstructive pulmonary disease, pulmonary hypertension, Alzheimer's disease, or a history of stroke or in patients on dialysis.¹⁴⁻²⁰ The average overall KCCQ summary score, particularly that of the subgroup of patients in NYHA functional class III-IV, was lower than that described in patients in international clinical trials on CHF.^{9,21,22} This finding highlights not only the worse HRQoL of patients with HF in the real world, but also the differences between populations included in trials and those treated in clinical practice.²³

This study found high correlations between specific and generic HRQoL measures. Higher correlations were found between the overall scores of quality of life measured with the KCCQ and the dimensions or items that measured physical limitation imposed by the disease. These results suggest that physical limitations and symptoms related to HF (physical component) determine the decreased HRQoL of these patients.^{8,24-26} Notably, although high correlations were found between the items related to physical limitations or symptoms on the KCCQ and the generic overall measures of the EQ-5D, they were slightly lower than those associated with the overall scores on the KCCQ for HF. This result




Figure 3. Unadjusted analysis of each dimension of the EuroQol-5D in patient subgroups with an independent association with health-related quality of life in the multivariable analysis. Percentage of patients who had some type of limitation in each of the 5 dimensions of EuroQol-5 dimensions according to sex (A); New York Heart Association functional class (B); age (C); Charlson comorbidity index (D); recent hospitalization (E), and clinical service (F).




Figure 3. (Continued)

suggests that the HRQoL of patients with HF could also be affected by other factors beyond the physical limitations imposed by the disease and other aspects unaddressed by specific instruments for HF. These aspects include impaired ability to maintain self-care, pain, anxiety or mood symptoms, all of which are equally relevant in the perceived state of health of patients with HF.^{6,8,9,24} This aspect highlights the relevance of assessing HRQoL in these patients using specific and generic instruments and underlines the multidimensional character of HRQoL.⁸

A prominent aspect of our study was to assess the clinical determinants of HRQoL in these patients. Thus, we draw attention to the originality of this study, given the current lack of Spanish multicenter studies that have assessed the factors that determine HRQoL in such a large number of patients with systolic HF. The

factors associated with worse disease progression, such as advanced age, comorbidity, recent hospitalization, or poor functional class, were independently associated with poor HRQoL. Many of these factors are not only associated with poor HRQoL,^{5,6,24-26} but are also associated with an increased risk of death or hospitalization.²³ In this sense, previous studies have shown that HRQoL is an independent predictor of these clinical events.^{27,28}

The association between sex and HRQoL found in this study may be related to the loss of the social role of women due to the limitations imposed by HF or to the possibility that the instruments that are designed to measure HRQoL better capture this information in women. Although the analyzes were adjusted for variables of severity of HF, patients attending cardiology




Figure 3. (Continued).

services had better HRQoL scores, which were probably due to their clinical profile being better than that of patients attending internal medicine services. It is also likely that these differences may have been due to factors not collected prospectively and which better define patients regarding social aspects or frailty.

No independent association was found between HRQoL and several variables commonly used to stratify patient risk factors (such as left ventricular ejection fraction, renal function, or hemoglobin function). This result highlights the importance of incorporating HRQoL as an additional measure when assessing patients with HF, because other clinical variables used to stratify

risk do not provide the information obtained from instruments that measure the patients' perception of health, nor do they provide information on limitations, which differs from those obtained using physiological or biological measures.⁸

Finally, the impact of HF on HRQoL should be assessed in specific geographic areas. The European Commission has drawn attention to the differences between European countries in perceived health status and the importance of conducting studies on these specific aspects in each geographical area. This study is of relevance, because it provides novel data on HRQoL in Spanish patients with HF and adds new data on the determinants of HRQoL

that complement published material on patients from other cultural or geographical environments.²⁹

Limitations

This study has the limitations inherent to all cross-sectional studies because it does not provide information on longitudinal changes of the study variable or on its association with the clinical determinants under investigation. The study population represented a subgroup of patients with HF and systolic dysfunction who are typically assessed in Spanish outpatient clinics. It is therefore not possible to determine if the results can be extrapolated to other populations of patients with HF, such as those with preserved left ventricular ejection fraction or who do not attend follow-up at outpatient clinics. This study addressed clinical variables and thus does not provide specific information on the impact of psychosocial variables or lifestyle and dietary habits on HRQoL.

CONCLUSIONS

This Spanish multicenter study found that patients with CHF have worse HRQoL than the general population and other patients with chronic diseases. High correlations were found between specific and generic measures of HRQoL. Several clinical factors, such as advanced age, female sex, advanced functional class, recent hospitalization, and greater comorbidity were associated with HRQoL independently of other prognostic factors. Patient treatment in Spanish cardiology services is independently associated with better quality of life, which is probably due to their better clinical profile.

ACKNOWLEDGEMENTS

We would like to thank all the researchers and patients who collaborated in the VIDA-IC study.

The study was acknowledged and supported by the Heart Failure and Transplantation Section of the Spanish Society of Cardiology and the Heart Failure Section of the Spanish Society of Internal Medicine.

Fieldwork was conducted by SANED. Statistical analyses were performed with the collaboration of ODDS S.L.

FUNDING

Pfizer S.L.U. promoted and funded the VIDA-IC study.

CONFLICTS OF INTEREST

J. Comín-Colet, M. Anguita, F. Formiga, L. Almenar, M.G. Crespo-Leiro, and L. Manzano received honoraria as members of the VIDA-IC assessment committee. J. Muñiz received honoraria for his collaboration in the independent statistical analysis. J. Chaves and T. de Frutos are employees of the Medical Department of Pfizer S.L.U. and collaborated in the VIDA-IC study.

SUPPLEMENTARY MATERIAL



Supplementary material associated with this article can be found in the online version available at [doi:10.1016/j.rec.2015.07.030](https://doi.org/10.1016/j.rec.2015.07.030).

REFERENCES

- Juenger J, Schellberg D, Kraemer S, Haunstetter A, Zugck C, Herzog W, et al. Health related quality of life in patients with congestive heart failure: comparison with other chronic diseases and relation to functional variables. *Heart*. 2002;87:235–41.
- Anker SD, Agewall S, Borggrefe M, Calvert M, Caro JJ, Cowie MR, et al. The importance of patient-reported outcomes: a call for their comprehensive integration in cardiovascular clinical trials. *Eur Heart J*. 2014;35:2001–9.
- Formiga F, Chivite D, Ortega C, Casas S, Ramón JM, Pujol R. End-of-life preferences in elderly patients admitted for heart failure. *QJM*. 2004;97:803–8.
- Rumsfeld JS, Alexander KP, Goff Jr DC, Graham MM, Ho PM, Masoudi FA, et al. Cardiovascular health: the importance of measuring patient-reported health status: a scientific statement from the American Heart Association. *Circulation*. 2013;127:2233–49.
- Myers J, Zaheer N, Quaglietti S, Madhavan R, Froelicher V, Heidenreich P. Association of functional and health status measures in heart failure. *J Card Fail*. 2006;12:439–45.
- Sullivan MD, Levy WC, Russo JE, Crane B, Spertus JA. Summary health status measures in advanced heart failure: relationship to clinical variables and outcome. *J Card Fail*. 2007;13:560–8.
- Heidenreich PA, Spertus JA, Jones PG, Weintraub WS, Rumsfeld JS, Rathore SS, et al. Health status identifies heart failure outpatients at risk for hospitalization or death. *J Am Coll Cardiol*. 2006;47:752–6.
- Comín-Colet J, Garin O, Lupón J, Manito N, Crespo-Leiro MG, Gómez-Bueno M, et al. Validación de la versión española del *Kansas City Cardiomyopathy Questionnaire*. *Rev Esp Cardiol*. 2011;64:51–8.
- Comin-Colet J, Lainscak M, Dickstein K, Filippatos GS, Johnson P, Lüscher TF, et al. The effect of intravenous ferric carboxymaltose on health-related quality of life in patients with chronic heart failure and iron deficiency: a subanalysis of the FAIR-HF study. *Eur Heart J*. 2013;34:30–8.
- Kind P, Dolan P, Gudex C, Williams A. Variations in population health status: results from a United Kingdom national questionnaire survey. *BMJ*. 1998;316:736–41.
- Anguita M, Comin-Colet J, Formiga F, Almenar L, Crespo-Leiro M, Manzano L. Tratamiento de la insuficiencia cardiaca con función sistólica deprimida: situación actual en España. Resultados del estudio VIDA-IC. *Rev Esp Cardiol*. 2014;67:769–70.
- Green CP, Porter CB, Bresnahan DR, Spertus JA. Development and evaluation of the Kansas City Cardiomyopathy Questionnaire: a new health status measure for heart failure. *J Am Coll Cardiol*. 2000;35:1245–55.
- Badia X, Roset M, Herdman M, Kind P. A comparison of United Kingdom and Spanish general population time trade-off values for EQ-5D health states. *Med Decis Making*. 2001;21:7–16.
- Encuesta Nacional de Salud 2011-2012 [cited 2015 Jul 16]. Madrid: Instituto Nacional de Estadística; 2013. Available at: <http://www.ine.es/jaxi/menu.do?type=pcaxis&path=/t15/p419&file=inebase&L=0>
- Mata Cases M, Roset Gamisans M, Badia Llach X, Antozanzas Villar F, Ragel Alcázar J. Impacto de la diabetes mellitus tipo 2 en la calidad de vida de los pacientes tratados en las consultas de atención primaria en España. *Aten Primaria*. 2003;31:493–9.
- Baquero M, Peset V, Burguera JA, Salazar-Cifre A, Boscá-Blasco ME, Del Olmo-Rodríguez A, et al. Calidad de vida en la enfermedad de Alzheimer. *Rev Neurol*. 2009;49:337–42.
- Miravittles M, Soriano JB, García-Río F, Muñoz L, Duran-Tauleria E, Sanchez G, et al. Prevalence of COPD in Spain: impact of undiagnosed COPD on quality of life and daily life activities. *Thorax*. 2009;64:863–8.
- Cobo Sánchez JL, Pelayo Alonso R, Ibarquien Rodríguez E, Aja Crespo A, Sáenz de Buruaga Perea A, Incera Setién ME, et al. Factores sociológicos y calidad de vida relacionada con la salud en pacientes en hemodiálisis. *Rev Soc Esp Enferm Nefrol*. 2011;14:98–104.
- Alonso J, Ferrer M, Gandek B, Ware Jr JE, Aaronson NK, Mosconi P, et al. Health-related quality of life associated with chronic conditions in eight countries: results from the International Quality of Life Assessment (IQOLA) Project. *Qual Life Res*. 2004;13:283–98.
- Badia Llach X, director. Estudios sobre la calidad de vida de pacientes afectados por determinadas patologías [cited 2015 Jul 16]. Madrid: Ministerio de Sanidad y Política Social; 2009. Available at: http://www.mssi.gob.es/organizacion/sns/planCalidadSNS/docs/Estudios_calidad_vida_pacientes.pdf
- McMurray JJ, Packer M, Desai AS, Gong J, Lefkowitz M, Rizkala AR, et al. Baseline characteristics and treatment of patients in Prospective comparison of ARNI with ACEI to determine impact on global mortality and morbidity in heart failure trial (PARADIGM-HF). *Eur J Heart Fail*. 2014;16:817–25.
- Ponikowski P, Van Veldhuisen DJ, Comin-Colet J, Ertl G, Komajda M, Mareev V, et al. Beneficial effects of long-term intravenous iron therapy with ferric carboxymaltose in patients with symptomatic heart failure and iron deficiency. *Eur Heart J*. 2015;36:657–68.
- Frigola-Capell E, Comin-Colet J, Davins-Miralles J, Gich-Saladich IJ, Wensing M, Verdú-Rotellar JM. Supervivencia de pacientes ambulatorios con insuficiencia cardiaca crónica del área mediterránea. Un estudio de base poblacional. *Rev Esp Cardiol*. 2013;66:539–44.
- Gutzwiller FS, Pfeil AM, Comin-Colet J, Ponikowski P, Filippatos G, Mori C, et al. Determinants of quality of life of patients with heart failure and iron deficiency treated with ferric carboxymaltose: FAIR-HF sub-analysis. *Int J Cardiol*. 2013;168:3878–83.

25. Enjuanes C, Klip IT, Bruguera J, Cladellas M, Ponikowski P, Banasiak W, et al. Iron deficiency and health-related quality of life in chronic heart failure: results from a multicenter European study. *Int J Cardiol.* 2014;174:268–75.
26. Comín-Colet J, Enjuanes C, González G, Torrens A, Cladellas M, Meroño O, et al. Iron deficiency is a key determinant of health-related quality of life in patients with chronic heart failure regardless of anaemia status. *Eur J Heart Fail.* 2013;15:1164–72.
27. Rodríguez-Artalejo F, Guallar-Castillón P, Rodríguez Pascual C, Montoto Otero C, Ortega Montes A, Nieto García A, et al. Health-related quality of life as a predictor of hospital readmission and death among patients with heart failure. *Arch Intern Med.* 2005;165:1274–9.
28. Zuluaga MC, Guallar-Castillón P, López-García E, Banegas JR, Conde-Herrera M, Olcoz-Chiva M, et al. Generic and disease-specific quality of life as a predictor of long-term mortality in heart failure. *Eur J Heart Fail.* 2010;12:1372–8.
29. Masseria C, Allin S, Sorenson C, Papanicolas I, Elias Mossialos. What are the methodological issues related to measuring health and drawing comparisons across countries? [cited 2015 Jul 16]. Available at: [http://ec.europa.eu/social/](http://ec.europa.eu/social/BlobServlet?docId=3951&langId=en)