

Original article

Childhood Excess Weight in Spain From 2006 to 2012. Determinants and Parental Misperception



María D. Ramiro-González,^{a,*} Belén Sanz-Barbero,^{b,c} and Miguel Ángel Royo-Bordonada^b

^a Servicio de Medicina Preventiva y Gestión de Calidad, Hospital General Universitario Gregorio Marañón, Madrid, Spain

^b Escuela Nacional de Sanidad, Instituto de Salud Carlos III (ISCIII), Madrid, Spain

^c CIBER de Epidemiología y Salud Pública, CIBERESP, Madrid, Spain

Article history:

Received 9 August 2016

Accepted 14 November 2016

Available online 18 March 2017

Keywords:

Childhood obesity
Childhood overweight
Parental misperception
Nutrition
Diet
Hours of sleep
Sedentary activity
Spain

ABSTRACT

Introduction and objectives: Due to the high prevalence of childhood obesity in Spain, we analyzed changes in its prevalence from 2006 to 2007 and from 2011 to 2012, as well as diet, sleep, and sedentary habits in 5- to 14-year-olds and parental misperceptions about their children's excess weight.

Methods: The sample was from the Spanish National Health Surveys for 2006 to 2007 (n = 5590) and for 2011 to 2012 (n = 3601). Data were collected by trained personnel through telephone interviews with parents/guardians. Weight and height were self-reported and the International Obesity Task Force cutpoints were used to define overweight and obesity.

Results: The prevalence of childhood excess weight was 30.1% from 2006 to 2007 and 29.7% from 2011 to 2012, while that of childhood obesity was 9.6% and 9%, respectively. Parental misperception of childhood excess weight increased from 60.8% to 71.4% ($P < .001$). Daily consumption of vegetables increased by 7.8%, while that of soft drinks and snacks decreased. This decrease was greatest in children from families with a low socioeconomic status, who also decreased their consumption of sweets and fast food. Adherence to sleep recommendations decreased by 5%, but adherence to recommended sedentary time did not change.

Conclusions: High childhood overweight and obesity rates remained stable in Spain from 2006 to 2007 and from 2011 to 2012, but there was an increase in parental misperception of childhood excess weight. Despite reduced consumption of soft drinks and snacks, there was low adherence to dietary recommendations, hours of sleep, and sedentary habits.

© 2016 Sociedad Española de Cardiología. Published by Elsevier España, S.L.U. All rights reserved.

Exceso de peso infantil en España 2006-2012. Determinantes y error de percepción parental

RESUMEN

Introducción y objetivos: Dada la gran prevalencia de obesidad infantil en España, se analiza su evolución entre 2006-2007 y 2011-2012, junto con dieta, hábitos de sueño y sedentarismo en la población de 5-14 años y la percepción parental del exceso de peso infantil.

Métodos: La muestra procede de las Encuestas Nacionales de Salud de 2006-2007 (n = 5.590) y 2011-2012 (n = 3.601). Las encuestas se realizaron mediante entrevista telefónica a los padres/tutores por personal entrenado. La información sobre peso y talla es la declarada por los encuestados, y se aplicaron los puntos de corte de sobrepeso/obesidad de la *International Obesity Task Force*.

Resultados: La prevalencia de exceso de peso fue del 30,1% en 2006-2007 y del 29,7% en 2011-2012, y la de obesidad, del 9,6 y el 9% respectivamente. La percepción errónea del exceso de peso pasó del 60,8 al 71,4% ($p < 0,001$). La ingesta diaria de verdura aumentó un 7,8% y la de refrescos y aperitivos disminuyó. Esta caída fue mayor en menores de clase social baja, que también redujeron el consumo de dulces y comida rápida. Mientras el cumplimiento de las recomendaciones sobre horas de sueño disminuyó un 5%, el de las relativas al tiempo máximo de actividad sedentaria no varió.

Conclusiones: Las altas cifras de obesidad y sobrepeso infantil permanecieron estables en España entre 2006-2007 y 2011-2012. El error perceptivo del exceso de peso aumentó entre los padres. Aunque disminuyó el consumo de refrescos y aperitivos, se observó un bajo cumplimiento de las recomendaciones dietéticas y las relativas a horas de sueño y ocio sedentario.

© 2016 Sociedad Española de Cardiología. Publicado por Elsevier España, S.L.U. Todos los derechos reservados.

Palabras clave:

Obesidad infantil
Sobrepeso infantil
Percepción parental errónea
Nutrición
Dieta
Horas de sueño
Actividad sedentaria
España

SEE RELATED CONTENT:

<http://dx.doi.org/10.1016/j.rec.2017.02.047>, Rev Esp Cardiol. 2017;70:629–630.

* Corresponding author: Servicio de Medicina Preventiva y Gestión de Calidad, Hospital General Universitario Gregorio Marañón, Dr. Esquerdo 46, 28007 Madrid, Spain. E-mail address: mariaramiro@gmail.com (M.D. Ramiro-González).

<http://dx.doi.org/10.1016/j.rec.2017.02.026>

1885-5857/© 2016 Sociedad Española de Cardiología. Published by Elsevier España, S.L.U. All rights reserved.

Abbreviation

ENSE: *Encuesta Nacional de Salud de España* (Spanish National Health Survey)

INTRODUCTION

Childhood overweight and obesity increased worldwide at the end of the 20th century,¹ with the prevalence of childhood obesity in Spain jumping from 4.9% in 1984 to 13.9% in 2000.² This trend was paralleled by an increase in the proportion of parents unable to recognize excess weight status in their children,³ reflecting the social normalization of obesity.⁴

From about 2000, childhood obesity prevalence began to stabilize in many countries.^{5,6} In Spain, the rate of increase slowed between 2001 and 2006 except among adolescents from a less privileged socioeconomic background.⁷ These developments were accompanied by increasing parental misperception of childhood weight status.³ Recent data for Spain are inconclusive. Compared with data from 2000,² the results of a 2011 study indicated an overall stabilization except for a continuing increase in obesity among 6- to 9-year-olds girls.⁸ In contrast, a 2012 study detected a reduced prevalence among children aged between 8 and 13 years.⁹ To our knowledge, no study since 2006 has analyzed changes in parental misperception of childhood excess weight.

Factors associated with obesity include consumption of sugar-sweetened soft drinks, a high-energy diet lacking in fruit and vegetables,^{10,11} insufficient hours of sleep,¹² and an excessively sedentary lifestyle.^{13,14} In Spain, a 1998 survey found that more than 40% of children and adolescents consumed below the recommended amount of fruit and vegetables,¹⁵ while other studies found that sugar intake among children accounted for 21% of total caloric intake (double the recommended amount)¹⁶ and that the consumption of sugar-sweetened soft drinks increased between 2001 and 2004.¹⁷ The proportion of 5- to 15-year-olds who slept less than 8 h/d rose from 2.9% in 2001 to 4% in 2003,¹⁸ and the proportion watching television for more than 2 h/d increased by 4% between 1997 and 2003.^{19,20}

In the present study, we analyzed data from the Spanish National Health Surveys (*Encuesta Nacional de Salud de España* [ENSE]) for changes in childhood obesity in Spain between 2006 to 2007²¹ and 2011 to 2012²² and their relationship with diet, hours of sleep, sedentary activity, and parental misperception of childhood excess weight.

METHODS

Data

We analyzed completed ENSE questionnaires from the most recent surveys of children and adolescents, carried out in 2011 to 2012²³ and 2006 to 2007.²⁴ These surveys have been conducted at varying intervals since 1987 by the Spanish National Institute of Statistics, in partnership with the Ministry of Health, Social Services, and Equality. Noninstitutionalized minors were included in the surveys by stratified multistage sampling. The first-stage sampling units were census tracts, second-stage units were family dwellings, and the third-stage units were individuals. Information was collected between July 2006 and June 2007 and between July 2011 and June 2012. The surveys were conducted by telephone interview; parents or guardians were informed about the survey and the voluntary and anonymous nature of their participation and, after giving verbal consent, answered questions about

children and young adolescents in their care. All questions and response options used to measure study variables were identical in the 2 surveys.

The 2011-to-2012 ENSE survey included 5495 0- to 14-year-olds, whereas the 2006-to-2007 survey included 9122 0- to 15-year-olds. The present analysis excluded 0- to 4-year-olds (n = 1894 in the 2011-to-2012 survey and n = 2869 in the 2006-to-2007 survey) and 15-year-olds (n = 663 in the 2006-to-2007 survey). The final sample thus included children and adolescents between the ages of 5 and 14 years: 3601 from the 2011-to-2012 survey and 5590 from the 2006-to-2007 survey. Due to incomplete weight and height data, body mass index was analyzed in subsamples of 2938 individuals from the 2011-to-2012 survey and 4341 from the 2006-to-2007 survey.

Variables

Overweight and Obesity

Overweight and obesity status was assigned according to body mass index calculated from self-reported weight and height measurements provided during the interview. Overweight and obesity together constituted excess weight and were defined according to International Obesity Task Force cutpoints.²⁵

Misperception of Excess Weight and Obesity

Parental misperception of childhood excess weight was evaluated with the following question: "In relation to his/her height, which of the following options best describes your child's weight: 1) substantially above normal, 2) slightly above normal, 3) normal, 4) below normal?" Misperception was defined as answers 3 or 4 in relation to a child with excess weight.

Daily Soft Drink and Food Consumption

Daily consumption of soft drinks and a range of foods (fruit, vegetables, sweets, fast foods, and snacks) was obtained with the question "How often does your child consume the following foods? 1) Every day, 2) Three or more times a week, 3) Once or twice a week, 4) Less than once a week, 5) Never or almost never."

Sleep Duration

Hours of sleep were obtained with the following question: "Can you tell me approximately how many hours your child normally sleeps per day, including naps." Reported sleep duration was evaluated by comparison with National Sleep Foundation and Centers for Disease Control and Prevention recommendations: a minimum 10 h/d for 5- to 9-year-olds and 9 h/d for 10- to 14-year-olds.^{26,27}

Sedentary Leisure Activity

Sedentary leisure activities include television viewing, playing video games, and computer use in free time. Time dedicated to these activities was obtained from the following questions: "Does your son/daughter watch television every day or almost every day?", "Approximately how much time per day does your son/daughter spend watching television?", "Does your son/daughter play video games, play on the computer, or access the internet every day or almost every day?" and "Approximately how much time per day does your son/daughter spend playing

Table 1
Sociodemographic Characteristics

	2006 to 2007, no. (%)	2011 to 2012, no. (%)
Age		
5 to 9 years	2631 (47.1)	1829 (50.8)
10 to 14 years	2959 (52.9)	1772 (49.2)
Sex		
Boys	2890 (51.7)	1851 (51.4)
Girls	2700 (48.3)	1750 (48.6)
Socioeconomic status		
High	1312 (23.8)	760 (22.5)
Middle	3508 (63.7)	2158 (64.0)
Low	683 (12.4)	454 (13.5)
Total	5590 (100)	3601 (100)

Results were obtained after application of the Spanish National Health Survey weighting coefficients.

video games, playing on the computer, or accessing the internet?" Reported sedentary activity was evaluated in relation to American Academy of Pediatrics recommendations of a maximum 2 h/d.²⁸

Sociodemographic variables

Collected sociodemographic variables were sex (boy or girl) and age (5 to 9 years or 10 to 14 years) and family socioeconomic status (high, middle, or low) as defined by the Spanish National Institute of Statistics according to criteria proposed by the Spanish Society of Epidemiology in 1995 and 2012. Family socioeconomic status was assigned according to the occupation and educational level of the

main family breadwinner.²⁹ Socioeconomic categories corresponded to Spanish Society of Epidemiology classifications as follows: high status to classes I and II; middle status to classes III, IVa, and IVb; and low status to class V.

Analysis

We conducted a descriptive analysis of each ENSE survey. Overall frequency distributions were calculated for the following qualitative variables: excess weight, obesity, parental misperception, daily food intake, and adherence to recommendations for hours of sleep and sedentary activity; frequency distributions were further stratified by age, sex, and family socioeconomic status. Overall prevalence was shown when stratified prevalence showed no differences. Quantitative variables (hours of sleep and sedentary activity) were expressed as means and 95% confidence intervals (95%CI). Because sleep recommendations differ for different age groups, sleep duration was stratified by age.

Qualitative variables in the 2 survey periods were compared by chi-square test. Sleep duration was compared by the Student *t* test, and hours dedicated to sedentary activities were compared by the nonparametric Mann-Whitney *U* test.

To estimate relative increases or decreases in dependent variables between the 2006-to-2007 and 2011-to-2012 surveys, prevalence ratios were obtained by Poisson regression with robust variance and adjustments for age, sex, and family socioeconomic status. Prevalence ratios for excess weight and obesity are not shown because there were no statistically significant between-survey differences.

The weighting coefficients included in the ENSE surveys were applied in all the analyses conducted. Data were analyzed with SPSS version 13 (SPSS, Inc.; Chicago, Illinois, United States) and

Table 2
Prevalence of Childhood Excess Weight and Obesity Stratified by Age, Sex, and Socioeconomic Status

	2006 to 2007		2011 to 2012		<i>P</i> [*]
	% (95%CI)	no.	% (95%CI)	no.	
Excess weight					
Age					
5 to 9 years	36.8 (34.6-39.0)	695	37.0 (34.5-39.5)	526	.913
10 to 14 years	25.0 (23.3-26.7)	614	22.9 (20.7-25.0)	347	.125
Sex					
Boys	31.8 (29.9-33.7)	714	30.3 (28.0-32.6)	465	.323
Girls	28.4 (26.5-30.3)	595	29.1 (26.7-31.5)	408	.654
Socioeconomic status					
High	24.4 (21.8-27.0)	257	21.1 (18.0-24.3)	136	.118
Middle	31.6 (29.8-33.3)	867	31.1 (29.0-33.2)	563	.736
Low	34.3 (30.1-38.5)	168	36.0 (30.7-41.2)	116	.611
Obesity					
Age					
5 to 9 years	15.4 (13.7-17.0)	290	14.7 (12.9-16.6)	209	.595
10 to 14 years	5.1 (4.2-5.9)	124	3.7 (2.7-4.6)	56	.050
Sex					
Boys	9.9 (8.7-11.2)	223	8.8 (7.4-10.2)	136	.268
Girls	9.1 (7.9-10.4)	192	9.2 (7.7-10.7)	129	.967
Socioeconomic status					
High	6.2 (4.7-7.7)	65	4.2 (2.6-5.7)	27	.080
Middle	10.3 (9.1-11.4)	282	10.5 (9.0-11.9)	189	.853
Low	12.5 (9.6-15.5)	62	8.1 (5.1-11.1)	26	.199

95%CI, 95% confidence interval; ENSE, *Encuesta Nacional de Salud de España* (Spanish National Health Survey).

Results were obtained after application of the ENSE weighting coefficients.

* Chi-square test comparison of prevalence between ENSE survey years 2006 to 2007 and 2011 to 2012.




Figure. Prevalence of childhood excess weight and obesity and the percentage of parents unable to recognize these characteristics in their children in 2006 to 2007 and 2011 to 2012.

STATA version 12.0 (StataCorp.; College Station, Texas, United States).

RESULTS

Sociodemographic characteristics of the samples are shown in Table 1. The prevalence of excess weight was 30.1% in 2006 to 2007 and 29.7% in 2011 to 2012 ($P = .696$). Obesity prevalence

values for the same survey years were 9.6% and 9%, respectively ($P = .485$). No significant changes were detected in the prevalence of excess weight or obesity adjusted for age, sex, or family socioeconomic status (Table 2). In both survey years, excess weight and obesity were more prevalent among 5- to 9-year olds ($P = .001$ for both survey years). The relationship between family socioeconomic status and overall excess weight strengthened between the 2006-to-2007 and 2011-to-2012 surveys, but the same trend was not observed for obesity.

The proportion of parents failing to recognize childhood excess weight increased from 60.8% (95%CI, 58.1%-63.4%) in 2006 to 2007 to 71.4% (95%CI, 68.4%-74.4%) in 2011 to 2012 ($P < .001$) (Figure 1). This increase was observed in all subgroups except for children from a high or low socioeconomic background (Table 3). Misperception of obesity increased by 10% between 2006 to 2007 and 2011 to 2012 ($P = .011$). Stratification revealed increases in parental misperception in relation to 10- to 14-year-olds, boys, and a middle socioeconomic status, whereas misperception decreased among parents with high socioeconomic status. Parental misperception of both overall excess weight and obesity was highest in relation to 5- to 9-year olds.

Daily consumption of fruit, vegetables, sweets, soft drinks, fast food, and snacks in 2006 to 2007 and 2011 to 2012 is shown in Table 4. These figures show an overall increase of 7.8% in vegetable consumption and decreases of 6.4% and 3.7% in the consumption of soft drinks and snacks, respectively. Stratification by socioeconomic status revealed that vegetable consumption increased most in children from a high socioeconomic status background and that consumption of soft drinks decreased most among children from families with a low socioeconomic status; children in this group

Table 3
Prevalence of Parental Misperception of Childhood Excess Weight and Obesity Stratified by Age, Sex, and Family Socioeconomic Status

	2006 to 2007		2011 to 2012		P^*	2011 to 2012:2006 to 2007 Prevalence Ratio (95%CI)
	% (95%CI)	no.	% (95%CI)	no.		
Parental misperception of childhood excess weight						
<i>Age</i>						
5 to 9 years	71.6 (68.2-74.9)	495	79.7 (76.2-83.1)	419	.001	1.64 (1.24-2.16)
10 to 14 years	48.6 (44.6-52.5)	298	58.7 (53.5-63.9)	203	.002	1.53 (1.16-2.01)
<i>Sex</i>						
Boys	59.8 (56.2-63.4)	426	71.1 (67.0-75.2)	331	< .001	1.60 (1.23-2.08)
Girls	61.9 (57.9-65.8)	367	71.6 (67.2-76.0)	292	.002	1.55 (1.16-2.08)
<i>Socioeconomic status</i>						
High	64.9 (59.0-70.8)	166	69.1 (61.2-77.0)	94	.433	1.17 (0.74-1.84)
Middle	60.8 (57.5-64.0)	524	73.5 (69.9-77.2)	414	< .001	1.67 (1.32-2.12)
Low	53.1 (45.5-60.7)	89	64.7 (55.9-73.6)	75	.052	1.74 (1.05-2.88)
Parental misperception of childhood obesity						
<i>Age</i>						
5 to 9 years	67.9 (62.7-73.5)	197	70.3 (64.3-76.7)	147	.624	1.26 (0.84-1.88)
10 to 14 years	20.3 (12.8-27.1)	25	37.5 (24.8-51.1)	21	.017	2.23 (1.06-4.68)
<i>Sex</i>						
Boys	45.7 (39.2-52.4)	101	60.7 (52.5-69.1)	82	.006	1.72 (1.07-2.77)
Girls	62.8 (56.0-69.8)	120	66.7 (58.4-74.9)	86	.552	1.11 (0.64-1.92)
<i>Socioeconomic status</i>						
High	69.2 (57.7-80.6)	45	51.9 (31.0-71.4)	14	.152	0.36 (0.14-0.93)
Middle	50.9 (45.1-56.8)	143	68.8 (62.3-75.6)	130	< .001	1.85 (1.20-2.83)
Low	45.9 (33.0-58.6)	28	46.2 (27.0-68.1)	12	.998	1.07 (0.38-2.03)

95%CI, 95% confidence interval; ENSE, Encuesta Nacional de Salud de España (Spanish National Health Survey).

Results were obtained after application of the ENSE weighting coefficients. Prevalence ratios are adjusted for age, sex, and family socioeconomic status.

* Chi-square test comparison of prevalence between ENSE survey years 2006 to 2007 and 2011 to 2012.

Table 4

Prevalence of the Daily Consumption of Fruit, Vegetables, Sweets, Soft Drinks, Fast Food, and Snacks in the Total Study Population and After Adjustment for Socioeconomic Status

Daily consumption	2006 to 2007		2011 to 2012		P [*]	2011 to 2012:2006 to 2007 prevalence ratio (95%CI)
	% (95%CI)	no.	% (95%CI)	no.		
<i>Fruit</i>						
Total	62.3 (61.0-63.6)	3357	60.3 (58.7-61.9)	2169	.064	0.90 (0.82-0.98)
High	70.7 (68.2-73.2)	894	66.6 (63.3-70.0)	506	.055	0.83 (0.68-1.01)
Middle	60.2 (58.5-61.8)	2049	59.0 (56.9-61.1)	1271	.368	0.94 (0.84-1.05)
Low	57.4 (53.6-61.1)	381	52.7 (48.1-57.4)	239	.127	0.83 (0.65-1.05)
<i>Vegetables</i>						
Total	29.0 (27.8-30.2)	1563	36.8 (35.2-38.3)	1322	< .001	1.44 (1.31-1.58)
High	33.4 (30.8-36.0)	422	42.8 (39.3-46.4)	325	< .001	1.50 (1.25-1.81)
Middle	28.0 (26.5-29.5)	952	35.4 (33.4-37.5)	763	< .001	1.41 (1.25-1.58)
Low	26.0 (22.7-29.4)	172	33.7 (29.3-38.1)	153	.005	1.43 (1.10-1.86)
<i>Candy</i>						
Total	46.5 (45.2-47.8)	2502	45.9 (44.3-47.6)	1651	.591	0.98 (0.90-1.07)
High	48.0 (45.2-50.8)	606	46.3 (42.7-49.8)	351	.455	0.94 (0.79-1.13)
Middle	44.5 (42.8-46.1)	1511	46.2 (44.1-48.3)	995	.202	1.07 (0.96-1.19)
Low	53.9 (50.1-57.7)	356	45.3 (40.7-49.9)	206	.005	0.70 (0.55-0.89)
<i>Soft drinks</i>						
Total	17.4 (16.4-18.4)	934	11.1 (10.0-12.1)	397	< .001	0.56 (0.49-0.63)
High	8.9 (7.4-10.5)	113	6.5 (4.7-8.3)	49	.044	0.73 (0.51-1.03)
Middle	17.7 (16.4-19.0)	601	10.8 (9.5-12.1)	233	< .001	0.57 (0.49-0.67)
Low	31.7 (28.2-35.3)	210	16.7 (13.3-20.2)	76	< .001	0.43 (0.32-0.58)
<i>Fast food</i>						
Total	5.4 (4.8-6.0)	290	4.7 (4.0-5.4)	169	.147	0.88 (0.73-1.08)
High	4.4 (3.3-5.5)	56	4.4 (3.0-5.8)	33	.920	1.00 (0.64-1.55)
Middle	5.6 (4.8-6.3)	189	5.4 (4.4-6.3)	116	.781	0.98 (0.77-1.24)
Low	6.6 (4.7-8.4)	43	2.6 (1.1-4.1)	12	.004	0.37 (0.20-0.72)
<i>Snacks</i>						
Total	6.2 (5.6-6.9)	334	2.4 (1.9-3.0)	88	< .001	0.36 (0.29-0.47)
High	3.2 (2.2-4.2)	40	0.4 (0.0-0.9)	3	< .001	0.13 (0.04-0.41)
Middle	6.2 (5.4-7.0)	211	2.9 (2.2-3.6)	63	< .001	0.46 (0.35-0.62)
Low	12.3 (9.8-14.8)	82	3.3 (1.7-5.0)	15	< .001	0.25 (0.14-0.44)

95%CI, 95% confidence interval; ENSE, *Encuesta Nacional de Salud de España* (Spanish National Health Survey).

Results were obtained after application of the ENSE weighting coefficients. Prevalence ratios are adjusted for age, sex, and family socioeconomic status.

* Chi-square test comparison of prevalence between ENSE survey years 2006 to 2007 and 2011 to 2012.

Table 5

Prevalence of Adherence to Recommendations for Childhood Sleep Duration and Sedentary Activities

Adherence to recommendations	2006 to 2007		2011 to 2012		P [*]	2011 to 2012:2006 to 2007 prevalence ratio (95%CI)
	% (95%CI)	no.	95%CI	no.		
<i>Sleep recommendations</i>						
5 to 9 years (10 h/d)	61.4 (59.5-63.3)	1606	55.7 (53.4-58.0)	1018	< .001	0.91 (0.85-0.97)
10 to 14 years (9 h/d)	65.5 (63.8-67.2)	1929	61.1 (58.9-63.4)	1084	.026	0.92 (0.87-0.97)
<i>Recommendations for TV and internet use (≤ 2 h/d)</i>	48.1 (46.7-49.4)	2497	46.8 (45.2-48.5)	1671	.260	0.95 (0.87-1.04)

95%CI, 95% confidence interval; ENSE, *Encuesta Nacional de Salud de España* (Spanish National Health Survey); TV, Television.

Results were obtained after application of the ENSE weighting coefficients. Prevalence ratios are adjusted for age, sex, and family socioeconomic status.

* Chi-square test comparison of prevalence between ENSE survey years 2006 to 2007 and 2011 to 2012.

also decreased their consumption of fast food and sweets. In 2011 to 2012, sweets consumption did not differ between socioeconomic groups.

In 2006 to 2007, 5- to 9-year-olds slept 9.67 h/d (95%CI, 9.63-9.71) and in 2011 to 2012 children in the same age group slept 9.52 h/d (95%CI, 9.47-9.57), a decrease of 9 min/d ($P < .001$).

Children aged 10 to 14 years slept 8.94 h/d in 2006 to 2007 (95%CI, 8.89-8.98) and 8.84 h/d in 2011 to 2012 (95%CI, 8.78-8.88), a 6-minute decrease ($P = .003$). Adherence to sleep recommendations thus decreased by 5.9% among 5- to 9-year-olds and by 4.5% among 10- to 14-year-olds (Table 5). After adjustment for sex and family socioeconomic status, the sleep duration values for 2011 to

2012 represent a reduction of approximately 10% from those for 2006 to 2007.

In 2006 to 2007, Spanish 5- to 14-year-olds dedicated an average 2.4 h/d (95%CI, 2.47-2.57) to sedentary activities. In 2011 to 2012, the figure was 2.5 h/d (95%CI, 2.38-2.48), representing an increase of 5 min/d ($P = .004$). Adherence to recommendations for maximum sedentary time did not vary, remaining close to 50% (Table 5). In both survey years, a positive association was observed between adherence to recommendations and higher family socioeconomic status (data not shown in the tables).

DISCUSSION

The prevalence of excess weight and obesity among Spanish 5- to 14-year-olds was similar in 2006 to 2007 and 2011 to 2012, both overall and after adjustment for age, sex, and family socioeconomic status. However, the intersurvey period showed a marked increase in the proportion of parents unable to recognize excess weight in their children. There was an overall increase in the consumption of vegetables and a decrease in the consumption of soft drinks and snacks. Children from a low socioeconomic background consumed fewer sweets and less fast food. There was an overall reduction in sleep duration and an increase in the time dedicated to sedentary activities.

These data reveal a stabilization of the prevalence of childhood excess weight and obesity in Spain. The reported prevalence values are similar to those among children aged between 8 and 9 years in the 1998-to-2000 eKid study² and the 2011 ALADINO study.⁸ The results presented here are the first to show a stabilization of excess weight prevalence among 10- to 14-year-olds. Despite this stabilization, Spanish rates of childhood excess weight and obesity are among the highest in the world.³⁰ The results of the present analysis coincide with those of other studies showing an accentuation of the relationship between childhood excess weight and family socioeconomic status in 2011 to 2012³¹ and a higher prevalence of excess weight and obesity in 5- to 9-year-olds.³²

Up until 2006, parental misperception of childhood excess weight in Spain tracked the increases in childhood obesity.³ Our study shows that parental misperception in Spain has continued to grow despite the stabilization of childhood weight, and is now one of the highest in the world.⁴ Although parental misperception of excess weight may be skewed by erroneous estimates of children's weight, there was a similar increase in parental misperception of obesity, which has a lower risk of weight misclassification. Possible causes of parental misperception include comparison with other children who are more obese,³³ fear of social stigmatization, and avoidance of guilt feelings.³ Other possible influences are the general increase in adult obesity and the associated social normalization of elevated body weight. As in other studies, misperception was highest among parents of younger children, for whom excess weight is widely accepted as a transitory characteristic that will disappear with growth.³³ The increasing misperception of childhood weight is worrying because parents who do not detect weight problems in their children are unlikely to take steps to modify obesogenic habits.^{34,35} The exception to this trend was the decreased misperception of childhood obesity among parents with a high socioeconomic status, possibly reflecting a higher capacity to resist social pressures among people with greater wealth or a higher educational level.

There was an overall positive trend to 2011 to 2012 in the consumption of vegetables, soft drinks, and snacks; however, consumption of sweets, fast foods, and fruit did not change. Vegetable consumption increased most in children from a high socioeconomic background, whereas the largest reductions in soft

drink, sweets, and fast food consumption were observed in children from a low socioeconomic background. In 2011 to 2012, sweets consumption did not differ according to family socioeconomic status. Consumption of fruit, vegetables, and soft drinks continues to show a relationship with socioeconomic status.³⁶ Vegetable consumption remains below WHO recommendations,³⁷ and the reduced soft drink consumption, while marked, is insufficient.³⁸ Consumption of unhealthy foods remained highest among children from a low socioeconomic background. Fruit consumption, among the highest in Europe,³⁹ did not change.

In 2011 to 2012, 5- to 9-year-olds slept 9 min/d less than the same age group in 2006 to 2007, whereas 10- to 14-year-olds slept 6 min/d less. These decreases translate into a drop in adherence to sleep recommendations of around 50%, in line with international studies.⁴⁰ Adherence to recommendations for maximum sedentary time did not change, remaining slightly below 50%. These findings are concerning because habits established in childhood and adolescence set the pattern for adulthood. There is, moreover, an established inverse relationship among children between sleep duration and the time devoted to sedentary activities.⁴¹ Sedentary behavior among children is widespread throughout the world,⁴² and is a major health determinant, independently of the level of physical activity.¹³

The stabilization of childhood weight and the positive trend in the consumption of some foods come in the wake of the 2005 launch of the Spanish strategy on nutrition, physical activity, and obesity prevention (NAOS),⁴³ and to some degree may reflect success of this strategy. However, a number of factors bring the effectiveness of the NAOS strategy into question; these include the discontinuation of some program activities,^{44,45} nonadherence to self-regulation agreements by the food and drink vending machine sector,⁴⁶ the doubtful effectiveness of the PAOS self-regulation code governing food advertising directed at children,⁴⁷ and the negative trends reported here in parental misperception, sleep duration, and sedentary activity. A possible alternative explanation for the observed changes is the effects of the economic crisis in Spain since 2008. Between 2006 and 2012, the price of food and nonalcoholic drinks increased by 12%,⁴⁸ and 41% of Spaniards claimed to have adjusted their dietary habits in order to save money.⁴⁹ These adjustments may have reduced the consumption of superfluous foods such as soft drinks and snacks and their replacement with other, healthier foods such as vegetables. This hypothesis is supported by the larger reduction in the consumption of these unhealthy foods among children from a low socioeconomic background. An earlier study attributed the major drop in body weight among Cubans during the 1991 to 1995 economic crisis to food shortages and an increase in manual labor.⁵⁰

Limitations

Self-reported survey data generally produce underestimates of body mass index because respondents tend to overestimate height and underestimate weight.⁵¹ However, studies in representative Spanish populations indicate that self-reported weight status is a valid parameter for classifying people as nonobese or obese.^{52,53} Our goal was not to produce precise estimates of obesity, but rather to analyze its temporal development from survey data obtained with the same methodology. Assuming that body mass index would be underestimated to the same degree in both survey years, the temporal analysis should be valid.

Sleep duration was not analyzed independently for weekdays and weekends because the questionnaires do not make this distinction.

There is a well-known association between childhood obesity and physical inactivity; however, this parameter was

not considered in the analysis due to the lack of quantitative data in the ENSE surveys related to the time dedicated to physical activity and its intensity.

CONCLUSIONS

Despite the stabilization of childhood obesity in Spain between 2006 to 2007 and 2011 to 2012, its prevalence remains excessively high, especially among children from a less privileged background. During the intersurvey period, parental misperception of childhood excess weight grew alarmingly. The ENSE surveys reveal that Spanish children do not adhere to national and international recommendations on diet, sleep duration, or sedentary behavior.

Policy directed at improving childhood and adolescent health should include obesity awareness programs. These programs should be targeted at parents, who play a fundamental role in establishing healthy habits, and at families with a low socioeconomic status, among whom there is greater potential for improving health through changes to dietary, sleep, and sedentary habits.^{54,55}

FUNDING

This project was financed by the Carlos III Institute of Health through the *Fondo de Investigación en Salud* (Project ENPY 1015/13).

CONFLICTS OF INTEREST

None declared.

WHAT IS KNOWN ABOUT THE TOPIC?

- The most recently published studies indicate stabilization of childhood overweight and obesity prevalence in Spain between 2000 and 2010 at around 30% (International Obesity Task Force cutpoints). Up until 2006, parental misperception of childhood weight tracked the increasing prevalence of childhood obesity. The increasing prevalence of childhood obesity in Spain during this period was accompanied by worsening dietary, sleep, and sedentary habits.

WHAT DOES THIS STUDY ADD?

- Childhood obesity prevalence in Spain has stabilized (9.6% in 2006 to 2007 and 9% in 2011 to 2012). Over the same period, the frequency of parental misperception of childhood excess weight increased by 11%, breaking with the previous tendency to track the prevalence of childhood obesity.
- Spanish children slept between 6 min/d and 9 min/d less in 2011 to 2012 than in 2006 to 2007 and spent 5 min/d more in front of an electronic device screen.
- Childhood consumption of soft drinks, sweets, and snacks decreased in the intersurvey period, with a more pronounced decline among children from a low socioeconomic background. Vegetable consumption increased.

REFERENCES

1. Wang Y, Lobstein T. Worldwide trends in childhood overweight and obesity. *Pediatr Obes*. 2006;1:11–25.
2. Serra L, Ribas L, Aranceta J, et al. Obesidad infantil y juvenil en España. Resultados del Estudio enKid (1998–2000). *Med Clin (Barc)*. 2003;121:725–732.
3. Salcedo V, Gutiérrez-Fisac JL, Guallar-Castillón P, et al. Trends in overweight and misperceived overweight in Spain from 1987 to 2007. *Int J Obes*. 2010;34:1759–1765.
4. Parry LL, Netuveli G, Parry J, et al. A systematic review of parental perception of overweight status in children. *J Ambul Care Manage*. 2008;31:253–268.
5. Canoy D, Buchan I. Challenges in obesity epidemiology. *Obes Rev*. 2007;8(Suppl 1):S1–S11.
6. Rokholm B, Baker JL, Sørensen TIA. The levelling off of the obesity epidemic since the year 1999—a review of evidence and perspectives. *Obes Rev*. 2010;11:835–846.
7. Miquelz E, Lostao L, Ortega P, et al. E. Trends in the prevalence of childhood overweight and obesity according to socioeconomic status: Spain, 1987–2007. *Eur J Clin Nutr*. 2014;68:209–214.
8. Pérez-Farínós N, López-Sobaler AM, Dal Re MA, et al. The ALADINO study: a national study of prevalence of overweight and obesity in Spanish children in 2011. *Biomed Res Int*. 2013;2013:163687.
9. Sánchez-Cruz JJ, Jiménez-Moleón JJ, Fernández-Quesada F, et al. Prevalencia de obesidad infantil y juvenil en España en 2012. *Rev Esp Cardiol*. 2013;66:371–376.
10. Te Morenga L, Mallard S, Mann J. Dietary sugars and body weight: systematic review and meta-analyses of randomised controlled trials and cohort studies. *BMJ*. 2012;346:e7492.
11. Monasta L, Batty GD, Cattaneo A, et al. Early-life determinants of overweight and obesity: a review of systematic reviews. *Obes Rev*. 2010;11:695–708.
12. Magee L, Hale L. Longitudinal associations between sleep duration and subsequent weight gain: A systematic review. *Sleep Med Rev*. 2012;16:231–241.
13. De Rezende LF, Rodrigues M, Rey-López JP, et al. Sedentary behavior and health outcomes: an overview of systematic reviews. *PLoS One*. 2014;9:e105620.
14. Amigo I, Busto R, Fernández C. La obesidad infantil como resultado de un estilo de vida obesogénico. *Endocrinol Nutr*. 2007;54:530–534.
15. Aranceta J, Pérez-Rodrigo C, Ribas L, et al. Sociodemographic and lifestyle determinants of food patterns in Spanish children and adolescents: the enKid study. *Eur J Clin Nutr*. 2003;57(Suppl 1):S40–S414.
16. Royo-Bordonada MA, Gorgojo L, Martín-Moreno JM, et al. Spanish children's diet: compliance with nutrient and food intake guidelines. *Eur J Clin Nutr*. 2003;57:930–939.
17. Últimos datos de consumo alimentario. Madrid: Ministerio de Agricultura, Alimentación y Medio Ambiente [accessed 6 Jun 2014]. <http://www.magrama.gob.es/es/alimentacion/temas/consumo-y-comercializacion-y-distribucion-alimentaria/panel-de-consumo-alimentario/ultimos-datos/default.aspx>.
18. Instituto Nacional Estadística. Encuesta Nacional de Salud. Tablas nacionales. Año 2003 [accessed 6 Jun 2014]. <http://www.ine.es/jaxi/tabla.do?path=/t15/p419/p02/a2003/I0/&file=02093.px&type=pcaxis>.
19. Ministerio de Sanidad, Servicios Sociales e Igualdad. Encuesta Nacional de Salud España 1997 [accessed 6 Jun 2014]. <https://www.msssi.gob.es/estadEstudios/estadisticas/encuestaNacional/encuesta1997.htm>.
20. Instituto Nacional Estadística. Encuesta Nacional de Salud. Tablas nacionales. Año 2003 [accessed 6 Jun 2014]. <http://www.ine.es/jaxi/tabla.do?path=/t15/p419/p02/a2003/I0/&file=02108.px&type=pcaxis>.
21. Encuesta Nacional de Salud España 2006. Ministerio de Sanidad, Servicios Sociales e Igualdad [accessed 2 Mar 2014]. <https://www.msssi.gob.es/estadEstudios/estadisticas/encuestaNacional/encuesta2006.htm>.
22. Encuesta Nacional de Salud de España 2011/12. Ministerio de Sanidad, Servicios Sociales e Igualdad [accessed 2 Mar 2014]. <https://www.msssi.gob.es/estadEstudios/estadisticas/encuestaNacional/encuesta2011.htm>.
23. Encuesta Nacional de Salud 2011/2012 Cuestionario de Menores. Ministerio de Sanidad, Servicios Sociales e Igualdad [accessed 2 Mar 2014]. http://www.msssi.gob.es/estadEstudios/estadisticas/encuestaNacional/encuestaNac2011/Cuestionario_Menores.pdf.
24. Encuesta Nacional de Salud 2006 Cuestionario de Menores. Ministerio de Sanidad, Servicios Sociales e Igualdad [accessed 2 Mar 2014]. http://www.msssi.gob.es/estadEstudios/estadisticas/encuestaNacional/encuestaNac2006/ENS_06_Menores_definitivo.pdf.
25. Cole TJ, Bellizzi MC, Flegal KM, et al. Establishing a standard definition for child overweight and obesity worldwide: international survey. *BMJ*. 2000;320:1240–1243.
26. How much sleep do we really need? National Sleep Foundation [accessed 4 Jun 2014]. <http://sleepfoundation.org/how-sleep-works/how-much-sleep-do-we-really-need>.
27. Insufficient sleep is a public health epidemic. Centers for Disease Control and Prevention [accessed 4 Jun 2014]. <http://www.cdc.gov/features/dss/sleep/>.
28. American Academy of Pediatrics. Committee on Public Education. American Academy of Pediatrics: Children, adolescents, and television. *Pediatrics*. 2001;107:423–426.
29. Metodología Encuesta Nacional de Salud de España 2011/2012. Ministerio de Sanidad, Servicios Sociales e Igualdad [accessed 4 Jun 2014]. http://www.msssi.gob.es/estadEstudios/estadisticas/encuestaNacional/encuestaNac2011/MetodologiaENSE2011_12.pdf.
30. Ng M, Fleming T, Robinson M, et al. Global, regional, and national prevalence of overweight and obesity in children and adults during 1980–2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet*. 2014;384:766–781.

31. Font-Ribera L, García-Continente X, Davó-Blanes MC, et al. El estudio de las desigualdades sociales en la salud infantil y adolescente en España. *Gac Sanit*. 2014;28:316–325.
32. Posso M, Brugulat-Guiteras P, Puig T, et al. Prevalencia y condicionantes de la obesidad en la población infantojuvenil de Cataluña, 2006–2012. *Med Clin (Barc)*. 2014;143:475–483.
33. Jones AR, Parkinson KN, Drewett RF, et al. Parental perceptions of weight status in children: the Gateshead Millennium Study. *Int J Obes (Lond)*. 2011;35:953–962.
34. Tiberio SS, Kerr DC, Capaldi DM, et al. Parental monitoring of children's media consumption: the long-term influences on body mass index in children. *JAMA Pediatr*. 2014;168:414–421.
35. Santiago-Torres M, Cui Y, Adams AK, et al. Familial and individual predictors of obesity and insulin resistance in urban Hispanic children. *Pediatr Obes*. 2016;11:54–60.
36. Miqueleiz E, Lostoa L, Ortega P, et al. Patrón socioeconómico en la alimentación no saludable en niños y adolescentes en España. *Aten Primaria*. 2014;46:433–439.
37. Diet, nutrition and the prevention of chronic diseases. WHO Technical Series 916 [accessed 10 Sep 2014]. http://whqlibdoc.who.int/trs/WHO_TRS_916.pdf.
38. Valoración de la Dieta Española de acuerdo al Panel de Consumo Alimentario. Fundación Española de Nutrición y Ministerio de Medio Ambiente y Medio Rural y Marino [accessed 9 Jul 2014]. http://www.mapama.gob.es/es/alimentacion/temas/consumo-y-comercializacion-y-distribucion-alimentaria/valoracion_panel_tcm7-7983.pdf.
39. Naska A, Fouskakis D, Oikonomou E, et al. Dietary patterns and their socio-demographic determinants in 10 European countries: data from the DAFNE databank. *Eur J Clin Nutr*. 2006;60:181–190.
40. Galland BC, Taylor BJ, Elder DE, et al. Normal sleep patterns in infants and children: A systematic review of observational studies. *Sleep Med Rev*. 2012;16:213–222.
41. Marinelli M, Sunyer J, Alvarez-Pedrerol M, et al. Hours of television viewing and sleep duration in children: a multicenter birth cohort study. *JAMA Pediatr*. 2014;168:458–464.
42. Eurodata TV Worldwide. Kids' TV Trends Global insight into the animation marketplace. 2012 [accessed 1 Jun 2014]. http://www.miptv.com/RM/RM_MIP-WORLD/2012/documents/white-papers/mipcom-2012-eurodata-white-paper-international-kids-tv-trends.pdf.
43. Estrategia NAOS. Ministerio de Sanidad, Servicios Sociales e Igualdad [accessed 3 May 2014]. <http://www.aesan.msssi.gob.es/AECOSAN/docs/documentos/nutricion/estrategianaos.pdf>.
44. Programa PERSEO. Agencia Española de Consumo, Seguridad Alimentaria y Nutrición [accessed 2 May 2014]. http://www.aecosan.msssi.gob.es/AECOSAN/docs/documentos/nutricion/educanaos/profesores_escuela_activa.pdf.
45. Programa PERSEO. Agencia Española de Consumo, Seguridad Alimentaria y Nutrición [accessed 2 May 2014]. <http://www.mecd.gob.es/educacion-mecd/areas-educacion/comunidades-autonomas/melilla/programas-educativos/programa-perseo.html>.
46. Royo-Bordonada MA, Martínez-Huedo MA. Evaluation of compliance with the self-regulation agreement of the food and drink vending machine sector in primary schools in Madrid, Spain, in 2008. *Gac Sanit*. 2014;28:65–68.
47. Romero-Fernández MM, Royo-bordonada MA, Rodríguez-Artalejo F. Evaluation of food and beverage television advertising during children's viewing time in Spain using the UK nutrient profile model. *Public Health Nutr*. 2013;16:1314–1320.
48. Productos y Servicios. El IPC en un clic. ¿Cuánto ha variado el IPC desde...? Cálculo variaciones del Índice de Precios de Consumo. Instituto Nacional Estadística [accessed 2 Jul 2014]. <http://www.ine.es/varipc/>.
49. Barómetro de diciembre. Avance de resultados. Estudio 2923. Diciembre 2011. Centro de Investigaciones Sociológicas (CIS) [accessed 2 Jul 2014]. http://datos.cis.es/pdf/Es2923mar_A.pdf.
50. Franco M, Bilal U, Orduñez P, et al. Population-wide weight loss and regain in relation to diabetes burden and cardiovascular mortality in Cuba 1980–2010: repeated cross sectional surveys and ecological comparison of secular trends. *BMJ*. 2013;346:f1515.
51. Nawaz H, Chan W, Abdulrahman M, et al. Self-reported weight and height: implications for obesity research. *Am J Prev Med*. 2001;20:294–298.
52. Basterra-Gortari FJ, Bes-Rastrollo M, Forga L, et al. Validity of self-reported body mass index in the National Health Survey. *An Sist Sanit Navar*. 2007;30:373–381.
53. Galan I, Gandarillas A, Febrel C, et al. Validation of self-reported weight and height in an adolescent population. *Gac Sanit*. 2001;15:490–497.
54. Laurson KR, Lee JA, Gentile DA, et al. Concurrent associations between physical activity, screen time, and sleep duration with childhood obesity. *ISRN Obes*. 2014;2014:204540.
55. Uso de las nuevas tecnologías por la infancia y adolescencia. Informe 2010. Observatorio de la Infancia en Andalucía (OIA) [accessed 20 May 2014]. http://www.observatoriodelainfancia.es/oia/esp/documentos_ficha.aspx?id=3022.