



apunts

MEDICINA DE L'ESPORT

www.apunts.org



SPECIAL ARTICLE

General guidelines in the rehabilitation process for return to training after a sports injury



Toni Caparrós^{a,b,*}, Montserrat Pujol^{c,d}, Carlos Salas^{e,f}

^a *Institut Nacional d'Educació Física de Catalunya (INEFC), Barcelona, Spain*

^b *SPARG Research Group, Universitat de Vic, Vic, Spain*

^c *Consell Català de l'Esport, Generalitat de Catalunya, Esplugues de Llobregat, Barcelona, Spain*

^d *Departament de Fisioteràpia, Facultat de Medicina y Ciències de la Salut, Universitat Internacional de Catalunya (UIC), Barcelona, Spain*

^e *Servicios Médicos Sanitas-Real Madrid CF, Madrid, Spain*

^f *Unidad de Medicina del Deporte, Hospital Universitario Sanitas La Moraleja, Madrid, Spain*

Received 27 December 2016; accepted 25 February 2017

Available online 15 May 2017

KEYWORDS

Rehabilitation;
Sports injury;
Planning;
Return to play;
Interdisciplinary

Abstract Rehabilitation for return to training is an interdisciplinary, specific and individualised process that is geared towards the sportsman's optimal availability for competition after a sports injury. This process begins after medical discharge and continues up to the return to play, involving the different professionals in the field of healthcare and led by the sports physician. Planning consists of three phases, defined by muscular action (isometric, concentric and eccentric); kinetic chain (closed or open); range of movement (internal, medial, external and total); and exercise orientation (general, directed and specific). Return to play must be agreed upon on the basis of objective information on the recovery stage of the injury, the state of fitness, and the sportsman's own perception.

© 2017 FC Barcelona. Published by Elsevier España, S.L.U. All rights reserved.

PALABRAS CLAVE

Readaptación;
Lesión deportiva;
Planificación;
Return to play;
Interdisciplinar

Pautas generales en el proceso de readaptación al entrenamiento después de una lesión deportiva

Resumen La readaptación a la práctica deportiva es el proceso interdisciplinar, específico e individualizado, que tiene como objetivo la óptima disponibilidad del deportista para la competición después de una lesión deportiva. Este proceso se lleva a cabo a partir del alta médica hasta el alta deportiva, e intervienen los diferentes profesionales del ámbito de la salud,

* Corresponding author.

E-mail address: info@tonicaparrós.eu (T. Caparrós).

liderados por el médico deportivo. Su planificación consta de 3 fases definidas por la acción muscular (isométrica, concéntrica y excéntrica), la cadena cinética (cerrada o abierta), la amplitud de trabajo (interna, media, externa y total) y la orientación de los ejercicios (general, dirigida y específica). La vuelta a la competición (*return to play*) debe ser consensuada a partir de información objetiva sobre la evolución de la lesión, el estado de forma, así como la propia percepción del deportista.

© 2017 FC Barcelona. Publicado por Elsevier España, S.L.U. Todos los derechos reservados.

Introduction

In its different disciplines, sport today is defined not only on the basis of performance level or fitness but also on availability and health.¹ In this context, the assessment of training load as well as injury prevention become an element which is an integral part of the training process.²⁻⁶ In spite of this, and considering this is one of the objectives to which sports organisations assign most resources,⁷ injuries continue to occur because certain levels of risk are intrinsic to the practice of sport.⁷⁻¹⁰

Once the sportsman is injured and must interrupt his professional routine,¹¹ regardless of severity, the recovery period begins. The first question asked by the sportsman¹² is, "What's the matter with me?", and the question that follows is ".and how long does that mean not playing?". The answer to the first question is purely medical, aimed at providing an exact diagnosis.^{11,13} The second question introduces a biopsychosocial working model to confront the complex¹⁴ process of injured sportsman until their return to play. This process is specific^{15,16} to each sportsman, injury and sport, and can be influenced by many different factors^{8,9} not only clinical but also involving coaches, clubs and, of course, the sportsman himself.¹⁷ When it comes to professional sportsmen, many other interests come into play, even economic ones. This is therefore a current topic of particular interest.^{8,9,18}

Background

Growing interest in the recovery of injuries has led to numerous studies being presented on the subject. One of the most important fields has been research directed towards the prevention of injuries.^{1,16,19-23} For more than three decades these studies have been used for rehabilitation^{24,25} since the profile of the sportsman is described with data that is used to set goals if the player gets injured. At present, and thanks to technological advance, it is possible to describe the condition of the sportsman better and in more detail.^{26,27} Other researchers have looked for elements that provide prognostic value and help clinicians in the decision-making process, whether these are derived from clinical examinations²⁸⁻³⁰ or from objective assessments based on findings in diagnostic tests.^{31,32} Finally, another line of research attempts to guide clinicians in the preparation or adaptation of programmes for the recovery of injured sportsmen.³³⁻³⁸ Within

this group we can include the present article, which presents an interdisciplinary, cross-sectional and current view of how to face up to the challenge presented whenever a sportsman is injured.

Definitions

Recently the scientific community has attempted to reach a consensus on the return to the play after injury.¹⁸ Rehabilitation for return to play has been defined as the interdisciplinary, specific and individualised process, the objective of which is the sportsman's optimal availability for competition after a sports injury.

There are several concepts that we need to pinpoint in order to standardise the language of health professionals in sport.¹⁸ The concept of *return to participation*, as it is termed internationally, refers to the point at which the sportsman begins to take part in activities specific to his sport, but to a lesser degree than the target set. When the target level is reached, participation in the activity is full and he/she can compete with no restriction. This is classified as *return to sport*. This term is the equivalent of "*return to play* (RTP),^{11,13,39} most commonly used in team sports.

It is necessary to distinguish between medical discharge and the return to full involvement in sport. Both will be decided by the physician.^{18,33} Medical discharge marks the healing of the sportsman's injury, as well as full functionality. In contrast, the sportsman will receive the all-clear for sport when he is fully able to compete.¹⁸

To complete the analysis, we must describe the professional scope of this rehabilitation process. Initially, a medical diagnosis will be fundamental, as well as treatment guidelines that must be determined by a physician specialised in sports medicine. Any injury should be treated by a specialist in rehabilitation of the functionality of the affected area. This specific training and these skills are encompassed within the profile of the physiotherapist. It will then be the specialist in sport and physical activity who will lead this phase of the process, with the objective of improving performance.⁴⁰ There is also the figure of the sports therapist who, depending on his training and skills, will be able to undertake the whole process or a significant proportion of it. This structure of different professionals is further extended in professional sport.^{14,17}

The profile of the sports therapist should be interdisciplinary, with specific training in sports injuries, kinesiology

and anatomy. To this specific training must be added a profile of friendliness, flexibility and empathy. The psychological management of an injury is not always easy, so the care required by the injured sportsman requires a high degree of dedication.¹³

In addition, over and above any other aspect, none of these professionals can ignore the fact that the true protagonist of the process of rehabilitation is the sportsman himself. During the injury period, the sportsman undergoes an internal process of *disidentification*. For an indefinite period, he ceases to be himself, without being able to do a job that is as vital as it is professional, since his body, seen as a working tool as well as of expression, is not capable of this.²⁹ His well-being,¹³ recovery^{28,31} and progress⁴⁰ to an optimal state of fitness^{14,30,33,35,36} are the only aspects which determine the process.

Description of the working model

The process of rehabilitation is cross-sectional and is based on teamwork by the different specialists in the field of sports medicine. Strictly speaking, rehabilitation to training begins at the moment the patient is discharged. From this point on, up to the point at which he can compete with full potential, rehabilitation is given. This will be an individualised and specific process.⁴¹ Its planning and programming should be oriented specifically to the sport practised and its competition requirements^{14,15,28}; to the sportsman, taking into consideration age, medical history, position of play, role and sports career; and, of course, to the injury: type, cause, location and recurrence.^{1,9,13}

The management of the phases^{30,32-34} and the information on them will be the responsibility of the doctor. Other professionals taking on this role could be counterproductive for the sportsman, his confidence and stability, as well as for achieving the goals. Taking into account these phases, the therapist and other professionals will design certain content for each of the phases the sportsman is at.

There are variables inherent in the injury that must be monitored and assessed at all times by the professionals who work with the sportsman. On this point, perception of pain is important.^{31,33,34} Professional care and its functionality, as well as the qualitative management of the treatment given and its intensity, should be defined as asymptomatic.^{33,40} Providing a workload that is restricted merely by pain, or progressing in intensity until reaching thresholds that are not fully tolerable, means taking that injury to limits not prepared for and running the premature and unnecessary risks of both relapse and overload of other areas, due to compensation.

The sportsman must be educated in pain management, as well as in the expression of perceiving it,^{32,34} investing all the necessary time until obtaining optimal communication with him. At the same time, personal trust between sportsman and therapist is a fundamental requirement that the professionals involved must know how to achieve in the day-to-day.

This will require that professionals set specific goals in the short term to prevent channelling motivation and concentration from going beyond that set of achievable achievements.³² Effort must be made to make

objectives assessable,¹⁶ and therefore have tests that mark the achievement of these.^{16,42-45} It is important that these objectives be defined for each phase and that the sportsman is aware of them, since, by definition, he is competitive and will be more motivated throughout the entire process.

Conditional orientation of the rehabilitation process

Contemporary sources already offer proposals on periodisation of the rehabilitation process, depending on the particular sport practised and its competition programming.¹⁴ This process must adopt the same training patterns as in the case of competition. That is, if team planning involves the project of a competitive programme, rehabilitation and its phases should also be planned as one single programme. This would be the case of individual sports and amateur team sports where only one competition is held. In the case of professional team sports with several different competitions, where planning can take up to seven phases, rehabilitation must understand and approach these in its final (directed and specific) stages.

Depending on the terms and specific conditions of both the injury and the sportsman, it may be considered necessary to undertake a certain degree of conditional work. This may be of a general, preventive or compensatory nature, without any intervention in the affected area prior to medical discharge.^{46,47} Subsequent to medical discharge, this could lead to more specific levels of intensity or directions (if deemed appropriate). Physical conditioning should be worked on in parallel with the injury, with the aim of being integrated within the rehabilitation phase considered.^{14,46,47} Its objectives are not only to reduce the duration of the period until return to sport but also prevention, maintenance (and improvement) of conditional modifications, both general and specific.

Planning the rehabilitation process

The proposal must be open; adapted to the specificity of each injury and type of sport; individualised for that particular sportsman; and restricted to the context of professionals who will undertake the process. Planning of the rehabilitation process will be divided into 3 different phases^{14,46,47,35,36,48} (Fig. 1) and the duration of each one will be determined by premises previously set out – as well as the achievement of specific clinical and conditional objectives.¹⁶ These phases define a progression for load and intensity^{49,50} as regards the exercises actually done. These will be defined by kinetic chain^{51,52}; type of muscular action; number of joints involved⁵³⁻⁵⁵; range of movement; and the technical and tactical approach to the actual practice of sport.⁵⁶

The first phase aims to achieve active early mobilisation, which provides a mechanical and circulatory benefit⁵⁷ without ignoring the fact that the whole process should be asymptomatic. Generally, this is characterised by proposing exercises in closed kinetic chain (CKC) and by muscular, isometric or dynamic action in internal or medial ranges. Their orientation will be of a general nature, that is to say, oriented towards the basic conditional and cognitive

	Phase 1	Phase 2	Phase 3
Kinetic chain	CKC	CKC/OKC	OKC
Muscular action	Isométric	Concentric	Eccentric
Extent of work	Internal - medial	External - total	Total
Orientation of content	General	Directed	Specific

Figure 1 General guidelines in the rehabilitation process for return to training after a sports injury.

capacities required in the specific type of sport practised.

The second phase is aimed at increasing functionality, intensity and load. There will be a progression of content in CKC to open kinetic chain (OKC) and extent of work, which enables concentric muscular work patterns, with an orientation directed at the particular work patterns (gestures) and needs of the specific sport. Specific sport exercises can be designed but without any opposition (defence) whatsoever.

Finally, the third phase will aim to reach competition-level intensity and loading, albeit proposed under training conditions. The content will bear in mind the intensity and maximum loading of the particular type of sport, as well as all its tactical complexity. This progression is achieved with the gradual entry of the sportsman into different phases of the training sessions with the team, focussed on qualitative (intensity) and quantitative (volume) approach.⁶ For most sports these exercises will be in OKC and the development of strength will be based on exercises which are eccentric in nature.

Objective assessment for return to play

Determining suitable progression in rehabilitation after sports injuries, up to achieving return to sport requires objective information,^{1,2} especially considering the fact that absence of pain is not synonymous with complete healing of the injured tissues.²⁶ Monitoring is needed which evaluates tissue response to loading from a multifactorial view.^{40,58} This requires the design of functional assessment that will enable us to control and monitor both with respect to conditional abilities and state of fitness,^{59,60} as well as adaptation to the training process.⁶¹ This process must be handled by means of reliable and applicable,¹⁶ non-invasive,⁶² submaximal,^{59,60} specific,⁴⁴ individualised^{63,64} tests.

These should be oriented to the effects of training on the sportsman, both in terms of internal load (based on parameters such as variability in heart rate, perception of fatigue or effort, among others), such as external load (time of exposure, distances, impacts, accelerations, decelerations).^{61,65,66} In this design, two types of parameters should not be ignored, such as subjective assessment of the sportsman himself^{26,29} (relative to internal load), due to its usefulness in the management of envisaged periods, and the use of isometric tests (external load). This will facilitate monitoring throughout the rehabilitation process.

The validity of the assessment will be determined by data collected prior to the injury. This will enable progress to be compared and assessed.

Future prospects

Sports training and the integration of technology^{2,5,6} offer the possibility of obtaining a multitude of different data.^{66,67} Accelerometry, tracking and GPS provide a non-invasive way of achieving specific and affordable values from speeds and acceleration rates during competition or training. This data is mainly related to manifestations of specific force in the form of acceleration (concentric) and deceleration (eccentric) rates; jumps; impacts; as well as achieved velocities; displacement; and changes in direction or distances.^{66,67} Beyond descriptive studies, research in this field focusses on the development of profiles based on the relationship of workloads with other types of parameters, through simple and ecological questionnaires (Borg, POMS, RPE) or Heart Rate Variability,^{68,69} which enables us to relate external and internal loading⁶⁵ and therefore, the effect of training on the development of the sportsman, as well as the management of optimal loads.²⁷

Conclusions

Once a sportsman has been injured, the recovery period begins, in order to obtain the sportsman's optimum availability for competition. Taking into account the factors that may influence the process, a framework will be designed using a biopsychosocial model, where the roles of each professional are clearly defined and where communication and trust between them and the player are maximum. The management of deadlines; content programming in each phase of work; and the creation of objectives are all essential elements for recovery of the injury, which requires specific, reliable and reproducible tests that will facilitate decision-making in an objective way.

Conflict of interests

Authors declare that they don't have any conflict of interests.

References

- Fuller CW, Junge A, Dvorak J. Risk management: FIFA's approach for protecting the health of football players. *Br J Sports Med.* 2011;46:11–7. <http://dx.doi.org/10.1136/bjsports-2011-090634>.
- Hugues H, Franks IM. Notational analysis of sport: systems for better coaching and performance in sport. 2nd edition London: Taylor & Francis; 2004. p. 320. ISBN: 0415290058.
- Carling C, Court M. Match & motion analysis of soccer. In: Williams M, editor. Science and soccer, developing elite performers. London: Routledge; 2012. p. 173–98.
- Gabbett TJ. Reductions in pre-season training loads reduce training injury rates in rugby league players. *Br J Sports Med.* 2004;38:743–9.
- Gabbett TJ, Domrow N. Relationships between training load, injury, and fitness in sub-elite collision sport athletes. *J Sports Sci.* 2007;25:1507–19. <http://dx.doi.org/10.1080/02640410701215066>.
- Gabbett TJ. The training-injury prevention paradox: should athletes be training smarter and harder? *Br J Sports Med Published Online First.* 2016. <http://dx.doi.org/10.1136/bjsports-2015-095788>.
- van Mechelen W, Hlobil H, Kemper HC. Incidence, severity, aetiology and prevention of sports injuries. A review of concepts. *Sports Med.* 2012;14:82–99.
- Fuller CW. Managing the risk of injury in sport. *Clin J Sport Med.* 2007;17:182–7. <http://dx.doi.org/10.1097/JSM.0b013e31805930b0>.
- Hägglund M, Waldén M, Til L, Pruna R. The importance of epidemiological research in sports medicine. *Apunts Medicina de l'Esport.* 2010;45:57–9. <http://dx.doi.org/10.1016/j.apunts.2010.02.006>.
- Fort Vanmeerhaeghe A, Romero Rodriguez D. Análisis de los factores de riesgo neuromusculares de las lesiones deportivas. *Apunts Med Esport.* 2013. <http://dx.doi.org/10.1016/j.apunts.2013.05.003>
- Hägglund M, Waldén M, Bahr R, Ekstrand J. Methods for epidemiological study of injuries to professional football players: developing the UEFA model. *Br J Sports Med.* 2005;39:340–6. <http://dx.doi.org/10.1136/bjism.2005.018267>.
- Moreira N, Vagetti G, de Oliveira V, Campos W. Asociación entre lesión y calidad de vida en deportistas: una revisión sistemática 1980–2013. *Apunts Med Esport.* 2014;49:123–38.
- Finch CF, Orchard JW, Twomey DM, Saleem MS, Ekegren CL, Lloyd DG, et al. Coding OSICS sports injury diagnoses in epidemiological studies: does the background of the coder matter. *Br J Sports Med.* 2014;48:552–6.
- Hoover DL, VanWye WR, Judge LW. Periodization and physical therapy: bridging the gap between training and rehabilitation. *Phys Therapy Sport.* 2015. <http://dx.doi.org/10.1016/j.ptsp.2015.08.003>
- Árnason Á. ¿Cuál es la Evidencia Científica en los Programas de Prevención de la Lesión Muscular? *Apunts Med Esport.* 2009;8:174–8.
- Bangsbo J, Mohr M, Poulsen A, Perez-Gomez J, Krstrup P. Training and testing the elite athlete. *J Exer Sci Fitness.* 2006;4:1–18.
- Matheson G, Shultz R, Bido J, Mitten M, Meeuwisse W, Shrier I. Return-to-play decisions: are they the team physician's responsibility? *Clin J Sport Med.* 2011;21:25–30.
- Ardern CL, Glasgow P, Schneiders A, Witvrouw E, Clarsen B, Cools A, et al. Consensus statement on return to sport from the First World Congress in Sports Physical Therapy, Bern. *Br J Sports Med.* 2016;50:853–64.
- Szudino JG, Mezêna B, Soncin R, Ferreira JC, Couto BP, Szmuchrowski LA. Pre vertical jump performance to regulate the training volume. *Int J Sports Med.* 2012;33:101–7. <http://dx.doi.org/10.1055/s-0031-1286293>.
- Gabbett TJ. The development and application of an injury prediction model for noncontact, soft-tissue injuries in elite collision sport athletes. *J Strength Conditioning Res.* 2010;24:2593–603.
- Gabbett TJ, Ullah S. Relationships between running loads and soft-tissue injury in elite team sport athletes. *J Strength Conditioning Res.* 2012;26:953–60.
- Hagglund M, Waldén M, Magnusson H, Kristenson K, Bengtsson H, Ekstrand J. Injuries affect team performance negatively in professional football: an 11-year follow-up of the UEFA Champions League injury study. *Br J Sports Med.* 2013;47:738–42. <http://dx.doi.org/10.1136/bjsports-2013-092215>.
- Raysmith B, Drew MK. Performance success or failure is influenced by weeks lost to injury and illness in elite Australian Track and Field athletes: a 5-year prospective study. *J Sci Med Sport.* 2016;19:778–83. <http://dx.doi.org/10.1016/j.jsams.2015.12.515>.
- Sapega AA, Minkoff J, Nicholas JA, Valsamis M. Sport-specific performance factor profiling: fencing as a prototype. *Am J Sports Med.* 1978;6:232–5.
- Schlinkman B. Norms for high school football players derived from cybex data reduction computer. *J Orthop Sports Phys Ther.* 1984;5:243–5.
- Gisselman AS, Baxter GD, Wright A, Hedemus E, Tumilty S. Musculoskeletal overuse injuries and heart rate variability: is there a link? *Med Hypotheses.* 2016;87:1–7.
- Caparrós T, Alentorn-Geli E, Myer GD, Capdevila L, Samuelsson K, Hamilton B, Rodas G. The relationship of practice exposure and injury rate on game performance and season success in professional male basketball. *J Sports Sci Med.* 2016;15:397–402. eCollection 2016.
- Moen MH, Reurink G, Weir A, Tol JL, Maas M, Goudswaard GJ. Predicting return to play after hamstring injuries. *Br J Sports Med.* 2014;48:1358–63.
- Klouche S, Lefevre N, Herman S, Gerometta A, Bohu Y. Return to sport after rotator cuff tear repair: a systematic review and meta-analysis. *Am J Sports Med.* 2016;44:1877–87.
- Zellers JA, Carmont MR, Grävare Silbernagel K. Return to play post-Achilles tendon rupture: a systematic review and meta-analysis of rate and measures of return to play. *Br J Sports Med.* 2016. Jun 3. pii: bjsports-2016-096106.
- Zwolski C, Schmitt LC, Quatman-Yates C, Thomas S, Hewett TE, Paterno MV. The influence of quadriceps strength asymmetry on patient-reported function at time of return to sport after anterior cruciate ligament reconstruction. *Am J Sports Med.* 2015;43:2242–9.
- Reurink G, Brilman EG, de Vos R-J, Maas M, Moen MH, Weir A, et al. Magnetic resonance imaging in acute hamstring injury: can we provide a return to play prognosis? *Sports Med.* 2015;45:133–46.
- Roi GS, Creta D, Nanni G, Marcacci M, Zaffagnini S, Snyder-Mackler L, et al. Return to official Italian first division soccer games within 90 days after anterior cruciate ligament reconstruction: a case report. *J Orthop Sports Phys Ther.* 2005;35:52–61.
- Short SM, Anloague PA, Strack DS. Rehabilitation and return to sport following surgical repair of the rectus abdominis and adductor longus in a professional basketball player: a case report. *J Orthop Sports Phys Ther.* 2016;3:1–32.
- Bizzini M, Hancock D, Impicizzeri F. Suggestions from the field for return to sports participation following anterior cruciate ligament reconstruction: soccer. *J Orthop Sports Phys Ther.* 2012;42:304–12.
- Waters E. Suggestions from the field for return to sports participation following anterior cruciate ligament reconstruction: basketball. *J Orthop Sports Phys Ther.* 2012;42:326–36.
- Verstegen M, Falsone S, Orr R, Smith S. Suggestions from the field for return to sports participation following anterior cruciate

- ate ligament reconstruction: American football. *J Orthop Sports Phys Ther.* 2012;42:337–44.
38. Sclafani MP, Davis CC. Return to play progression for rugby following injury to the lower extremity: a clinical commentary and review of the literature. *Int J Sports Phys Ther.* 2016;11:302–20.
 39. Ekstrand J, Hägglund M, Waldén M. Epidemiology of muscle injuries. In *professional football players have residual isokinetic deficit.* Br J Sports Med. 2014. Feb 3. pii: bjsports-2013-093016.
 40. Tol J, Hamilton B, Eirale C, Muxart P, Jacobsen P, Whiteley R. At return to play following hamstring injury the majority of professional football players have residual isokinetic deficit. *Br J Sports Med.* 2014. Feb 3. pii: bjsports-2013-093016.
 41. Pedret C, Badius R. Lesions musculares en el deporte Actualización de un artículo del Dr. Cabot, publicado en *Apuntes de Medicina Deportiva en 1965.* Apuntes Med Esport. 2015;50:111–20.
 42. Frisch A, Urhausen A, Seil R, Croisier JL, Windal T, Theisen D. Association between Preseason functional tests and Injury in youth football: a prospective follow-up. *Scand J Med Sci Sports.* 2011;21:468–76, <http://dx.doi.org/10.1111/j.1600-0838.2011.01369.x>.
 43. Gabbett TJ, Stein JG, Kemp JG, Lorenzen C. Relationships between tests of physical qualities and physical match performance in elite rugby league players. *J Strength Conditioning Res.* 2013;27:1539–45.
 44. Drinkwater EJ, Moore NR, Bird SP. Effects of changing from full range of motion to partial range of motion on squad kinetics. *J Strength Conditioning Res/Natl Strength Conditioning Assoc.* 2012;26:890–6.
 45. Drinkwater EJ, Pyne DB, McKenna MJ. Design and interpretation of anthropometric and fitness testing of basketball players. *Sports Med.* 2008;38:565–78.
 46. Futbol Club Barcelona Serveis Mèdics. Guia de Pràctica Clínica de les Lesions Musculars. Epidemiologia, Diagnòstic, Tractament i Prevenció. *Apuntes de Med Esport.* 2009;164:179–204.
 47. Futbol Club Barcelona. Serveis Mèdics., Guia de Pràctica Clínica de les, Tendinopaties, Diagnòstic, Tractament i, Prevenció. *Apuntes de Med Esport.* 2012;62:1–24, <http://dx.doi.org/10.1016/j.apunts.2012.09.001>.
 48. Bizzini M, Hancock D, Impellizzeri F. Suggestions from the field for return to sports participation following anterior cruciate ligament reconstruction: Soccer. *J Orthop Sports Phys Ther.* 2012 Apr;42:304–12.
 49. Verhoshanski Y, Chornonson G. Jump exercises in sprint training. *Track Field Q.* 1909;9:1967–9.
 50. Roig Pull M, Ranson C. Eccentric muscle actions: implications for injury prevention and rehabilitation. *Phys Therapy Sport.* 2007;8:88–97.
 51. Ekstrom R, Osborn R, Goehner H, Moen A, Ommen B, Mefferd M, et al. Electromyographic normalization procedures for determining exercise intensity of closed chain exercises for strengthening the quadriceps femoris muscles. *J Strength Conditioning Res.* 2012;26:766–71.
 52. De Mey K, Danneels L, Cagnie B, Borms D, T'jonck Z, Van Damme E, et al. Shoulder muscle activation levels during four closed kinetic chain exercises with and without record slings. *J Strength Conditioning Res.* 2014;28:1626–35.
 53. Sherry M, Best T. A comparison of 2 rehabilitation programs in the treatment of acute hamstring strain. *J Orthop Sports Phys Ther.* 2004;34(3.).
 54. LaStayo P, Wolf J, Lewek M, Sneyder-Mackler L, Reich T, Lindstedt L. Eccentric muscle contractions: their contribution to injury, prevention, rehabilitation and sport. *J Orthop Sports Phys Ther.* 2003;33.
 55. Järvinen T, Järvinen T, Kääriäinen M, Kalimo H, Järvinen M. Muscle injuries. *Biology and treatment.* *Am J Sports Med.* 2005;33, <http://dx.doi.org/10.1177/0363546505274714>.
 56. Ribera N. Planificació a largo plazo en los deportes de equipo. *Apuntes de Seirul.lo. RED 2009, Tomo XXIII, num 4.*
 57. Orchard J, Best TM, Verrall GM. Return to play following muscle strains. *Clin J Sport Med.* 2005;15:436–41.
 58. Pruna R. Return to play: ¿Hacia dónde vamos? Esto no es un juego de adivinanzas. *Apuntes Med Esport.* 2016;51:109–12.
 59. Bangsbo J, Iaia FM, Krstrup P. The Yo-Yo intermittent recovery test intermittent sports. *Sports Med (Auckland NZ).* 2008;38:37–51.
 60. Caparrós T, Padullés JM, Rodas G, Capdevila LI. Can the strength test performance and injury rate at professional basketball? *Apuntes Educació Física Esports.* 2014;118:48–58.
 61. Drew MK, Finch CF. The relationship between training load and injury illness and soreness: a systematic and literature review. *Sports Med.* 2016;46:861–83.
 62. Rebelo A, Brito J, Seabra A, Oliveira J, Drust B, Krstrup P. A new tool to measure training load in soccer training and match play. *Int J Sports Med.* 2012;33:297–304, <http://dx.doi.org/10.1055/s-0031-1297952>.
 63. Gray AJ, Jenkins DG. Match analysis and the physiological demands of international football. *Sports Med (Auckland NZ).* 2010;40:347–60, <http://dx.doi.org/10.2165/11531400-000000000-00000>.
 64. Živ G, Lidor R. Vertical jump in female and male basketball players – a review of observational and experimental studies. *J Sci Med Sport/Sports Med Australia.* 2010;13:332–9, <http://dx.doi.org/10.1016/j.jsams.2009.02.009>.
 65. Scanlan AT, Wen N, Tucker P, Borges N, Dalbo V. The relationships between internal and external training load models during basketball training. *J Strength Conditioning Res.* 2014;28:2397–405.
 66. Casamichana D, Castellano J, Calleja J, San Román J, Castagna C. Relationship between indicators of training load in soccer players. *J Strength Conditioning Res.* 2012. Publish Ahead of Print DOI: 10.1519/JSC.0b013e3182548af1.
 67. Buchheit M, Gray M, Morin JB. Assessing stride variables and vertical stiffness with the GPS-embedded accelerometers: preliminary insights for the monitoring of neuromuscular fatigue on the field. *J Sports Sci Med.* 2015;14:698–701.
 68. Scanlan AT, Wen N, Tucker P, Borges N, Dalbo V. Training mode's influence on the relationships between training-load models during basketball conditioning international. *J Sports Physiol Perform.* 2014;9:851–6, <http://dx.doi.org/10.1123/ijspp.2013-0410>.
 69. Moreno J, Parrado E, Capdevila LI. Variabilidad de la frecuencia cardíaca y perfiles psicofisiológicos en deportes de equipo de alto rendimiento. *Rev Psicol Deporte.* 2013;22:345–52. ISSN: 1132-239X.